

# VARIATIONAL PRINCIPLES IN MATHEMATICAL PHYSICS, GEOMETRY, AND ECONOMICS

Alexandru Kristály, Vicențiu D. Rădulescu and  
Csaba György Varga

Variational Principles in Mathematical  
Physics, Geometry, and Economics

Qualitative Analysis of Nonlinear Equations and Unilateral  
Problems

Alexandru Kristály, Vicențiu D. Rădulescu, Csaba Varga



# Contents

<i>Foreword</i>	<i>page</i> vii
<i>Preface</i>	ix

## **P A R T I: Variational Principles in Mathematical Physics**

<b>1</b>	<b>1 Variational Principles</b>	<b>2</b>
1.1	Minimization techniques and Ekeland variational principle	2
1.2	Borwein-Preiss variational principle	8
1.3	Minimax principles	12
1.4	Ricceri's variational results	20
1.5	$H^1$ versus $C^1$ local minimizers	30
1.6	Szulkin type functionals	35
1.7	Pohozaev's fibering method	41
1.8	Historical comments	42
<b>2</b>	<b>Variational Inequalities</b>	<b>45</b>
2.1	Introduction	45
2.2	Variational inequalities on unbounded strips	46
2.3	Area-type variational inequalities	59
2.4	Historical notes and comments	83
<b>3</b>	<b>Nonlinear Eigenvalue Problems</b>	<b>86</b>
3.1	Weighted Sobolev spaces	87
3.2	Eigenvalue problems	91
3.3	Superlinear case	99
3.4	Sublinear case	111
3.5	Comments and further perspectives	123
<b>4</b>	<b>Elliptic Systems of Gradient Type</b>	<b>125</b>

4.1	Introduction	125
4.2	Formulation of the problems	126
4.3	Systems with superlinear potential	128
4.4	Systems with sublinear potential	136
4.5	Shift solutions for gradient systems	144
4.6	Comments and historical notes	154
<b>5</b>	<b>Systems with Arbitrary Growth Nonlinearities</b>	<b>156</b>
5.1	Introduction	156
5.2	Elliptic systems with mountain pass geometry	159
5.3	Elliptic systems with oscillatory terms	164
5.4	Comments and perspectives	172
<b>6</b>	<b>Scalar Field Systems</b>	<b>173</b>
6.1	Introduction	173
6.2	Multiple solutions of a double eigenvalue problem	174
6.3	Scalar field systems with nonlinear oscillatory terms	184
6.4	Applications	190
6.5	Comments and historical notes	194
<b>7</b>	<b>Competition Phenomena in Dirichlet Problems</b>	<b>195</b>
7.1	Introduction	196
7.2	Effects of the competition	197
7.3	A general location property	203
7.4	Nonlinearities with oscillation near the origin	206
7.5	Nonlinearities with oscillation at infinity	212
7.6	Perturbation from symmetry	219
7.7	Historical notes, comments	223
<b>8</b>	<b>Problems to Part I</b>	<b>225</b>
<b>P A R T II: Variational Principles in Geometry</b>		<b>230</b>
<b>9</b>	<b>Sublinear Problems on Riemannian Manifolds</b>	<b>231</b>
9.1	Introduction	231
9.2	Existence of two solutions	232
9.3	Existence of many global minima	239
9.4	Historical notes and comments	242
<b>10</b>	<b>Asymptotically Critical Problems on Spheres</b>	<b>244</b>
10.1	Introduction	244
10.2	Group-theoretical argument	246
10.3	Arbitrarily small solutions	251
10.4	Arbitrarily large solutions	259

10.5	Historical notes, comments and perspectives	264
<b>11</b>	<b>Equations with Critical Exponent</b>	266
11.1	Introduction	266
11.2	Subcritical case	268
11.3	Critical case	271
11.4	Comments and historical notes	278
<b>12</b>	<b>Problems to Part II</b>	280
<b>P A R T III: Variational Principles in Economics</b>		285
<b>13</b>	<b>Mathematical Preliminaries</b>	285
13.1	Metrics, geodesics, flag curvature	285
13.2	Busemann-type inequalities on Finsler manifolds	290
13.3	Variational inequalities	296
<b>14</b>	<b>Minimization of Cost-functions on Manifolds</b>	298
14.1	Introduction	298
14.2	A necessary condition	301
14.3	Existence and uniqueness results	302
14.4	Examples on the Finslerian-Poincaré disc	306
14.5	Comments and further perspectives	308
<b>15</b>	<b>Best Approximation Problems on Manifolds</b>	311
15.1	Introduction	311
15.2	Existence of projections	312
15.3	Geometric properties of projections	314
15.4	Geodesic convexity and Chebyshev sets	317
15.5	Optimal connection of two submanifolds	321
15.6	Remarks and perspectives	327
<b>16</b>	<b>A Variational Approach to Nash Equilibria</b>	329
16.1	Introduction	329
16.2	Nash equilibria and variational inequalities	330
16.3	Nash equilibria for set-valued maps	333
16.4	Lack of convexity: a Riemannian approach	339
16.5	Historical comments and perspectives	345
<b>17</b>	<b>Problems to Part III</b>	347
<b>Appendices</b>		351
<b>Appendix A Elements of convex analysis</b>		353
A.1	Convex sets and convex functions	353
A.2	Convex analysis in Banach spaces	358

<b>Appendix B</b>	<b>Function spaces</b>	360
B.1	Lebesgue spaces	360
B.2	Sobolev spaces	361
B.3	Compact embedding results	363
B.4	Sobolev spaces on Riemann manifolds	367
<b>Appendix C</b>	<b>Category and genus</b>	370
<b>Appendix D</b>	<b>Clarke and Degiovanni gradients</b>	373
D.1	Locally Lipschitz functionals	373
D.2	Continuous or lower semi-continuous functionals	375
<b>Appendix E</b>	<b>Elements of set-valued analysis</b>	381
	<i>References</i>	384
	<i>Index</i>	401

# Foreword

The use of variational principles has a long and fruitful history in mathematics and physics, both in solving problems and shaping theories, and it has been introduced recently in economics. The corresponding literature is enormous and several monographs are already classical. The present book 'Variational Principles in Mathematical Physics, Geometry and Economics', by Kristály, Rădulescu and Varga, is original in several ways.

In Part I, devoted to variational principles in mathematical physics, unavoidable classical topics like Ekeland variational principle, mountain pass lemma and Ljusternik-Schnirelmann category are supplemented with more recent methods and results of Ricceri, Brezis-Nirenberg, Szulkin and Pohozaev. The chosen applications cover variational inequalities on unbounded strips and for area-type functionals, nonlinear eigenvalue problems for quasilinear elliptic equations, and a substantial study of systems of elliptic partial differential equations. This is a challenging topics of growing importance, with many applications in natural and human sciences, like demography.

Part II shows the importance of variational problems in geometry. Classical questions like geodesics or minimal surfaces are not considered, but instead the authors concentrate on a less standard problem, namely the transformation of classical questions related to the Emden-Fowler equation into problems defined on some four-dimensional sphere. The combination of the calculus of variations with group theory provides interesting results. The case of equations with critical exponents, which is of special importance in geometrical problems since Yamabe's work, is also treated.

Part III deals with variational principles in economics. Some choice is also necessary in this area, and the authors first study the minimization

of cost-functions on manifolds, with special attention on the Finslerian-Poincaré disc. Then they consider best approximation problems on manifolds before approaching Nash equilibria through variational inequalities.

The high level of mathematical sophistication requested in all three parts could be an obstacle for potential readers more interested in applications. However several Appendices recall in a precise way the basic concepts and results of convex analysis, functional analysis, topology and set-valued analysis. Because the present in science depends upon its past and shapes its future, historical and bibliographical notes are complemented by perspectives. Some exercises are proposed as complements to the covered topics.

Among the wide recent literature on critical point theory and its applications, the authors had to make a selection. Their choice has of course been influenced by their own tastes and contributions. It is a happy one, because of the interest and beauty of selected topics, because of their potential for applications, and because of the fact that most of them have not been covered in existing monographs. Hence I believe that the book of Kristály, Rădulescu and Varga will be appreciated by all scientists interested in variational methods and in their applications.

Jean Mawhin  
Académie Royale de Belgique

# Preface

For since the fabric of the  
universe is most perfect and  
the work of a most wise  
Creator, nothing at all takes  
place in the universe in which  
some rule of maximum or  
minimum does not appear.

---

Leonhard Euler (1707–1783)

The understanding of the nature is impossible without an understanding of the partial differential equations and variational principles that govern a large part of physics. That is why, it is not surprising that nonlinear partial differential equations first arose from an interplay of physics and geometry. The roots of the calculus of variations go back to the 17th century. Indeed, Johann Bernoulli raised as a challenge the “brachistochrone problem” in 1696. The same year, when he heard of this problem, Sir Isaac Newton found that he could not sleep until he had solved it. Having done so, he published the solution anonymously. Bernoulli, however, knew at once that the author of the solution was Newton and, cf. [296], in a famous remark asserted that he “recognized the Lion by the print of its paw”.

However, the modern calculus of variations appeared in the middle of the 19th century, as a basic tool in the qualitative analysis of models arising in physics. Indeed, *it was Riemann who aroused great interest in them [problems of the calculus of variations] by proving many interesting results in function theory by assuming Dirichlet’s principle* (Charles B. Morrey Jr., [219]). The characterization of phenomena by means of variational principles has been a cornerstone in the transition from classical to contemporary physics. Since the middle part of the

twentieth century, the use of variational principles has developed into a range of tools for the study of nonlinear partial differential equations and many problems arising in applications. Cf. Ioffe and Tikhomirov [145], the term “variational principle” refers essentially to a group of results showing that a lower semi-continuous, lower bounded function on a complete metric space possesses arbitrarily small perturbations such that the perturbed function will have an absolute (and even strict) minimum.

Very often, important equations and systems (Yang-Mills equations, Einstein equations, Ginzburg-Landau equations, etc.) describing phenomena in applied sciences arise from the minimization of energy functionals like

$$E(u) = \int_{\Omega} f(x, u(x), \nabla u(x)) \, dx.$$

The class  $\mathcal{C}$  of admissible functions  $u = (u^1, \dots, u^n) : \Omega \subset \mathbb{R}^N \rightarrow \mathbb{R}^n$  may be constrained, for instance, by boundary conditions, while the function  $f = f(x, u, p) : \Omega \times \mathbb{R}^{nN} \rightarrow \mathbb{R}$  is assumed to be sufficiently smooth and verifying natural growth conditions. Formally, the variational problem

$$\min_{u \in \mathcal{C}} E(u) \tag{0.1}$$

gives rise to the nonlinear elliptic system of partial differential equations

$$-\sum_{j=1}^n \frac{\partial}{\partial x^j} f_{p_j^i}(x, u(x), \nabla u(x)) + f_{u^i}(x, u(x), \nabla u(x)) = 0, \tag{0.2}$$

for all  $1 \leq i \leq n$ . The simplest example corresponding to

$$f(x, u, p) = \frac{1}{2} |p|^2$$

implies that problem (0.1) is associated to the minimization of the Dirichlet integral

$$E(u) = \frac{1}{2} \int_{\Omega} |\nabla u|^2 \, dx$$

and (0.2) reduces to the Laplace equation

$$\Delta u = 0.$$

A more sophisticated example corresponds to the function

$$f(x, u, p) = \sum_{i,j=1}^n \sum_{s,t=1}^N g_{ij}(u) \gamma^{st}(x) p_s^i p_t^j,$$

where  $\gamma = (\gamma_{st})_{1 \leq s,t \leq N}$  is an invertible matrix with inverse  $\gamma^{-1} =$

$(\gamma^{st})_{1 \leq s, t \leq N}$ , while  $g = (g_{ij})_{1 \leq i, j \leq n}$  is a uniformly positive definite matrix. In this case, problem (0.1) yields a generalization of the Dirichlet integral on suitable manifolds and (0.2) becomes an equation of the type

$$-\Delta_{\mathcal{M}} u = \sum_{i, j=1}^n \sum_{s, t=1}^N \gamma^{st} \Gamma_{ji}^i u_{x^s}^j u_{x^t}^k.$$

The differential operator  $\Delta_{\mathcal{M}}$  denotes the Laplace–Beltrami operator and  $\Gamma_{ji}^i$  are the Christoffel symbols.

This monograph is an original attempt to develop the modern theory of the calculus of variations from the points of view of several disciplines. This theory is one of the twin pillars on which nonlinear functional analysis is built. The authors of this volume are fully aware of the limited achievements of this volume as compared with the task of understanding the force of variational principles in the description of many processes arising in various applications. Even though necessarily limited, the results in this book benefit from many years of work by the authors and from interdisciplinary exchanges between them and other researchers in this field.

One of the main objectives of this book is to let physicists, geometers, engineers, and economists know about some basic mathematical tools from which they might benefit. We would also like to help mathematicians learn what applied calculus of variations is about, so that they can focus their research on problems of real interest to physics, economics, engineering, as well as geometry or other fields of mathematics. We have tried to make the mathematical part accessible to the physicist and economist, and the physical part accessible to the mathematician, without sacrificing rigor in either case. The mathematical technicalities are kept to a minimum within the book, enabling the discussion to be understood by a broad audience. Each problem we develop in this book has its own difficulties. That is why we intend to develop some standard and appropriate methods that are useful and that can be extended to other problems. However, we do our best to restrict the prerequisites to the essential knowledge. We define as few concepts as possible and give only basic theorems that are useful for our topic. The authors use a first-principles approach, developing only the minimum background necessary to justify mathematical concepts and placing mathematical developments in context. The only prerequisite for this volume is a standard graduate course in partial differential equations, drawing especially from linear elliptic equations to elementary variational methods, with a

special emphasis on the maximum principle (weak and strong variants). This volume may be used for self-study by advanced graduate students and as a valuable reference for researchers in pure and applied mathematics and related fields. Nevertheless, both the presentation style and the choice of the material make the present book accessible to all newcomers to this modern research field which lies at the interface between pure and applied mathematics.

Each chapter gives full details of the mathematical proofs and subtleties. The book also contains many exercises, some included to clarify simple points of exposition, others to introduce new ideas and techniques, and a few containing relatively deep mathematical results. Each chapter concludes with historical notes. Five appendices illustrate some basic mathematical tools applied in this book: elements of convex analysis, function spaces, category and genus, Clarke and Degiovanni gradients, and elements of set-valued analysis. These auxiliary chapters deal with some analytical methods used in this volume, but also include some complements. This unique presentation should ensure a volume of interest to mathematicians, engineers, economists, and physicists. Although the text is geared toward graduate students at a variety of levels, many of the book's applications will be of interest even to experts in the field.

We are very grateful to Diana Gillooly, Editor for Mathematics, for her efficient and enthusiastic help, as well as for numerous suggestions related to previous versions of this book. Our special thanks go also to Clare Dennison, Assistant Editor for Mathematics and Computer Science, and to the other members of the editorial technical staff of Cambridge University Press for the excellent quality of their work.

Our vision throughout this volume is closely inspired by the following prophetic words of Henri Poincaré on the role of partial differential equations in the development of other fields of mathematics and in applications: *A wide variety of physically significant problems arising in very different areas (such as electricity, hydrodynamics, heat, magnetism, optics, elasticity, etc...) have a family resemblance and should be treated by common methods.* (Henri Poincaré, [249]).

Alexandru Kristály, Vicențiu D. Rădulescu, Csaba Varga

July 2009

`alexandrukristaly@yahoo.com`

`vicentiu.radulescu@imar.ro`

`varga_gy_csaba@yahoo.com`

P A R T I

Variational Principles in Mathematical  
Physics

# 1

## Variational Principles

A man is like a fraction whose numerator is what he is and whose denominator is what he thinks of himself. The larger the denominator the smaller the fraction.

---

Leo Tolstoy (1828–1910)

Variational principles are very powerful techniques at the interplay between nonlinear analysis, calculus of variations, and mathematical physics. They have been inspired and have deep applications in modern research fields such as geometrical analysis, constructive quantum field theory, gauge theory, superconductivity, etc.

In this chapter we shortly recall the main variational principles which will be used in the sequel, as Ekeland and Borwein-Preiss variational principles, minimax- and minimization-type principles (mountain pass theorem, Ricceri-type multiplicity theorems, Brézis-Nirenberg minimization technique), the principle of symmetric criticality for non-smooth Szulkin-type functionals, as well as the Pohozaev's fibering method.

### 1.1 Minimization techniques and Ekeland variational principle

Many phenomena arising in applications (such as geodesics or minimal surfaces) can be understood in terms of the minimization of an energy functional over an appropriate class of objects. For the problems of mathematical physics, phase transitions, elastic instability, and diffraction of

the light are among the phenomena that can be studied from this point of view.

A central problem in many nonlinear phenomena is if a bounded from below and lower semi-continuous functional  $f$  attains its infimum. A simple function when the above statement clearly fails is  $f : \mathbb{R} \rightarrow \mathbb{R}$  defined by  $f(s) = e^{-s}$ . Nevertheless, further assumptions either on  $f$  or on its domain may give a satisfactory answer. In the sequel, we give two useful forms of the well-known Weierstrass theorem.

**Theorem 1.1** [Minimization; compact case] *Let  $X$  be a compact topological space and  $f : X \rightarrow \mathbb{R} \cup \{+\infty\}$  a lower semi-continuous functional. Then  $f$  is bounded from below and its infimum is attained on  $X$ .*

*Proof* The set  $X$  can be covered by the open family of sets  $S_n := \{u \in X : f(u) > -n\}$ ,  $n \in \mathbb{N}$ . Since  $X$  is compact, there exists a finite number of sets  $S_{n_0}, \dots, S_{n_l}$  which also cover  $X$ . Consequently,  $f(u) > -\max\{n_0, \dots, n_l\}$  for all  $u \in X$ .

Let  $s = \inf_X f > -\infty$ . Arguing by contradiction, we assume that  $s$  is not achieved which means in particular that  $X = \bigcup_{n=1}^{\infty} \{u \in X : f(u) > s+1/n\}$ . Due to the compactness of  $X$ , there exists a number  $n_0 \in \mathbb{N}$  such that  $X = \bigcup_{n=1}^{n_0} \{u \in X : f(u) > s+1/n\}$ . In particular,  $f(u) > s+1/n_1$  for all  $u \in X$  which is in contradiction with  $s = \inf_X f > -\infty$ .  $\square$

The following result is a very useful tool in the study of various partial differential equations where no compactness is assumed on the domain of the functional.

**Theorem 1.2** [Minimization; noncompact case] *Let  $X$  be a reflexive Banach space,  $M$  be a weakly closed, bounded subset of  $X$ , and  $f : M \rightarrow \mathbb{R}$  be a sequentially weak lower semi-continuous function. Then  $f$  is bounded from below and its infimum is attained on  $M$ .*

*Proof* We argue by contradiction, that is, we assume that  $f$  is not bounded from below on  $M$ . Then, for every  $n \in \mathbb{N}$  there exists  $u_n \in M$  such that  $f(u_n) < -n$ . Since  $M$  is bounded, the sequence  $\{u_n\} \subset M$  is so. Due to the reflexivity of  $X$ , one may subtract a subsequence  $\{u_{n_k}\}$  of  $\{u_n\}$  which weakly converges to an element  $\tilde{x} \in X$ . Since  $M$  is weakly closed,  $\tilde{x} \in M$ . Since  $f : M \rightarrow \mathbb{R}$  is sequentially weak lower semi-continuous, we obtain that  $f(\tilde{x}) \leq \liminf_{k \rightarrow \infty} f(u_{n_k}) = -\infty$ , contradiction. Therefore,  $f$  is bounded from below.

Let  $\{u_n\} \subset M$  be a minimizing sequence of  $f$  over  $M$ , that is,  $\lim_{n \rightarrow \infty} f(u_n) = \inf_M f > -\infty$ . As before, there is a subsequence  $\{u_{n_k}\}$  of  $\{u_n\}$  which

weakly converges to an element  $\bar{x} \in M$ . Due to the sequentially weak lower semi-continuity of  $f$ , we have that  $f(\bar{x}) \leq \liminf_{k \rightarrow \infty} f(u_{n_k}) = \inf_M f$ , which concludes the proof.  $\square$

For any bounded from below, lower semi-continuous functional  $f$ , Ekeland's variational principle provides a minimizing sequence whose elements minimize an appropriate sequence of perturbations of  $f$  which converges locally uniformly to  $f$ . Roughly speaking, Ekeland's variational principle states that there exist points which are almost points of minima and where the "gradient" is small. In particular, it is not always possible to minimize a nonnegative continuous function on a complete metric space. Ekeland's variational principle is a very basic tool that is effective in numerous situations, which led to many new results and strengthened a series of known results in various fields of analysis, geometry, the Hamilton-Jacobi theory, extremal problems, the Ljusternik-Schnirelmann theory, etc.

Its precise statement is as follows.

**Theorem 1.3** [Ekeland's variational principle] *Let  $(X, d)$  be a complete metric space and let  $f : X \rightarrow \mathbb{R} \cup \{\infty\}$  be a lower semi-continuous, bounded from below functional with  $D(f) = \{u \in X : f(u) < \infty\} \neq \emptyset$ . Then for every  $\varepsilon > 0$ ,  $\lambda > 0$ , and  $u \in X$  such that*

$$f(u) \leq \inf_X f + \varepsilon$$

*there exists an element  $v \in X$  such that*

- a)  $f(v) \leq f(u)$ ;
- b)  $d(v, u) \leq \frac{1}{\lambda}$ ;
- c)  $f(w) > f(v) - \varepsilon \lambda d(w, v)$  for each  $w \in X \setminus \{v\}$ .

*Proof* It is sufficient to prove our assertion for  $\lambda = 1$ . The general case is obtained by replacing  $d$  by an equivalent metric  $\lambda d$ . We define the relation on  $X$ :

$$w \leq v \iff f(w) + \varepsilon d(v, w) \leq f(v).$$

It is easy to see that this relation defines a partial ordering on  $X$ . We now construct inductively a sequence  $\{u_n\} \subset X$  as follows:  $u_0 = u$ , and assuming that  $u_n$  has been defined, we set

$$S_n = \{w \in X : w \leq u_n\}$$

and choose  $u_{n+1} \in S_n$  so that

$$f(u_{n+1}) \leq \inf_{S_n} f + \frac{1}{n+1}.$$

Since  $u_{n+1} \leq u_n$  then  $S_{n+1} \subset S_n$  and by the lower semi-continuity of  $f$ ,  $S_n$  is closed. We now show that  $\text{diam} S_n \rightarrow 0$ . Indeed, if  $w \in S_{n+1}$ , then  $w \leq u_{n+1} \leq u_n$  and consequently

$$\varepsilon d(w, u_{n+1}) \leq f(u_{n+1}) - f(w) \leq \inf_{S_n} f + \frac{1}{n+1} - \inf_{S_n} f = \frac{1}{n+1}.$$

This estimate implies that

$$\text{diam} S_{n+1} \leq \frac{2}{\varepsilon(n+1)}$$

and our claim follows. The fact that  $X$  is complete implies that  $\bigcap_{n \geq 0} S_n = \{v\}$  for some  $v \in X$ . In particular,  $v \in S_0$ , that is,  $v \leq u_0 = u$  and hence

$$f(v) \leq f(u) - \varepsilon d(u, v) \leq f(u)$$

and moreover

$$d(u, v) \leq \frac{1}{\varepsilon}(f(u) - f(v)) \leq \frac{1}{\varepsilon}(\inf_X f + \varepsilon - \inf_X f) = 1.$$

Now, let  $w \neq v$ . To complete the proof we must show that  $w \leq v$  implies  $w = v$ . If  $w \leq v$ , then  $w \leq u_n$  for each integer  $n \geq 0$ , that is  $w \in \bigcap_{n \geq 0} S_n = \{v\}$ . So,  $w \not\leq v$ , which is actually c).  $\square$

In  $\mathbb{R}^N$  with the Euclidean metric, properties a) and c) in the statement of Ekeland's variational principle are completely intuitive as Figure 1.1 shows. Indeed, assuming that  $\lambda = 1$ , let us consider a cone lying below the graph of  $f$ , with slope  $+1$ , and vertex projecting onto  $u$ . We move up this cone until it first touches the graph of  $f$  at some point  $(v, f(v))$ . Then the point  $v$  satisfies both a) and c).

In the particular case  $X = \mathbb{R}^N$  we can give the following simple alternative proof to Ekeland's variational principle, due to Hiriart-Urruty, [141]. Indeed, consider the perturbed functional

$$g(w) := f(w) + \varepsilon \lambda \|w - u\|, \quad w \in \mathbb{R}^N.$$

Since  $f$  is lower semi-continuous and bounded from below, then  $g$  is lower semi-continuous and  $\lim_{\|w\| \rightarrow \infty} g(w) = +\infty$ . Therefore there exists  $v \in \mathbb{R}^N$  minimizing  $g$  on  $\mathbb{R}^N$  such that for all  $w \in \mathbb{R}^N$

$$f(v) + \varepsilon \lambda \|v - u\| \leq f(w) + \varepsilon \lambda \|w - u\|. \quad (1.1)$$




Figure 1.1 Geometric illustration of Ekeland's variational principle.

By letting  $w = u$  we find

$$f(v) + \varepsilon\lambda \|v - u\| \leq f(u)$$

and a) follows. Now, since  $f(u) \leq \inf_{\mathbb{R}^N} f + \varepsilon$ , we also deduce that  $\|v - u\| \leq 1/\lambda$ .

We infer from relation (1.1) that for any  $w$ ,

$$f(v) \leq f(w) + \varepsilon\lambda [\|w - u\| - \|v - u\|] \leq f(w) + \varepsilon\lambda \|w - u\|,$$

which is the desired inequality c).

Taking  $\lambda = \frac{1}{\sqrt{\varepsilon}}$  in the above theorem we obtain the following property.

**Corollary 1.1** *Let  $(X, d)$  be a complete metric space and let  $f : X \rightarrow \mathbb{R} \cup \{\infty\}$  be a lower semi-continuous, bounded from below and  $D(f) = \{u \in X : f(u) < \infty\} \neq \emptyset$ . Then for every  $\varepsilon > 0$  and every  $u \in X$  such that*

$$f(u) \leq \inf_X f + \varepsilon$$

*there exists an element  $u_\varepsilon \in X$  such that*

- a)  $f(u_\varepsilon) \leq f(u)$ ;
- b)  $d(u_\varepsilon, u) \leq \sqrt{\varepsilon}$ ;
- c)  $f(w) > f(u_\varepsilon) - \sqrt{\varepsilon}d(w, u_\varepsilon)$  for each  $w \in X \setminus \{u_\varepsilon\}$ .

Let  $(X, \|\cdot\|)$  be a real Banach space,  $X^*$  its topological dual endowed with its natural norm, denoted for simplicity also by  $\|\cdot\|$ . We denote by  $\langle \cdot, \cdot \rangle$  the duality mapping between  $X$  and  $X^*$ , that is,  $\langle x^*, u \rangle = x^*(u)$  for every  $x^* \in X^*, u \in X$ . Theorem 1.3 readily implies the following property, which asserts the existence of *almost critical points*. In other words, Ekeland's variational principle can be viewed as a generalization of the Fermat theorem which establishes that interior extrema points of a smooth functional are, necessarily, critical points of this functional.

**Corollary 1.2** *Let  $X$  be a Banach space and let  $f : X \rightarrow \mathbb{R}$  be a lower semi-continuous functional which is bounded from below. Assume that  $f$  is Gâteaux differentiable at every point of  $X$ . Then for every  $\varepsilon > 0$  there exists an element  $u_\varepsilon \in X$  such that*

- (i).  $f(u_\varepsilon) \leq \inf_X f + \varepsilon$ ;
- (ii).  $\|f'(u_\varepsilon)\| \leq \varepsilon$ .

Letting  $\varepsilon = 1/n, n \in \mathbb{N}$ , Corollary 1.2 gives rise a minimizing sequence for the infimum of a given function which is bounded from below. Note however that such a sequence need not converge to any point. Indeed, let  $f : \mathbb{R} \rightarrow \mathbb{R}$  defined by  $f(s) = e^{-s}$ . Then,  $\inf_{\mathbb{R}} f = 0$ , and any minimizing sequence fulfilling (a) and (b) from Corollary 1.2 tends to  $+\infty$ . The following definition is dedicated to handle such situations.

**Definition 1.1** (a) A function  $f \in C^1(X, \mathbb{R})$  satisfies the Palais-Smale condition at level  $c \in \mathbb{R}$  (shortly,  $(PS)_c$ -condition) if every sequence  $\{u_n\} \subset X$  such that  $\lim_{n \rightarrow \infty} f(u_n) = c$  and  $\lim_{n \rightarrow \infty} \|f'(u_n)\| = 0$ , possesses a convergent subsequence.

(b) A function  $f \in C^1(X, \mathbb{R})$  satisfies the Palais-Smale condition (shortly,  $(PS)$ -condition) if it satisfies the Palais-Smale condition at every level  $c \in \mathbb{R}$ .

Combining this compactness condition with Corollary 1.2, we obtain the following result.

**Theorem 1.4** *Let  $X$  be a Banach space and a function  $f \in C^1(X, \mathbb{R})$  which is bounded from below. If  $f$  satisfies the  $(PS)_c$ -condition at level  $c = \inf_X f$ , then  $c$  is a critical value of  $f$ , that is, there exists a point  $u_0 \in X$  such that  $f(u_0) = c$  and  $u_0$  is a critical point of  $f$ , that is,  $f'(u_0) = 0$ .*

## 1.2 Borwein-Preiss variational principle

The Borwein-Preiss variational principle [45] is an important tool in infinite dimensional nonsmooth analysis. This basic result is strongly related with Stegall's variational principle [282], *smooth bumps* on Banach spaces, Smulyan's test describing the relationship between Fréchet differentiability and the strong extremum, properties of continuous convex functions on separable Asplund spaces, variational characterizations of Banach spaces, the Bishop-Phelps theorem or Phelps' lemma [243]. The generalized version we present here is due to Loewen and Wang [198] and enables us to deduce the standard form of the Borwein-Preiss variational principle, as well as other related results.

Let  $X$  be a Banach space and assume that  $\rho : X \rightarrow [0, \infty)$  is a continuous function satisfying

$$\rho(0) = 0 \quad \text{and} \quad \rho_M := \sup\{\|x\|; \rho(x) < 1\} < +\infty. \quad (1.2)$$

An example of function with these properties is  $\rho(x) = \|x\|^p$  with  $p > 0$ .

Given the families of real numbers  $\mu_n \in (0, 1)$  and vectors  $e_n \in X$  ( $n \geq 0$ ), we associate to  $\rho$  the *penalty function*  $\rho_\infty$  defined for all  $x \in X$ ,

$$\rho_\infty(x) = \sum_{n=0}^{\infty} \rho_n(x - e_n), \quad \text{where} \quad \rho_n(x) := \mu_n \rho((n+1)x). \quad (1.3)$$

**Definition 1.2** For the function  $f : X \rightarrow \mathbb{R} \cup \{+\infty\}$ , a point  $x_0 \in X$  is a strong minimizer if  $f(x_0) = \inf_X f$  and every minimizing sequence  $(z_n)$  of  $f$  satisfies  $\|z_n - x_0\| \rightarrow 0$  as  $n \rightarrow \infty$ .

We observe that any strong minimizer of  $f$  is, in fact, a strict minimizer, that is  $f(x) > f(x_0)$  for all  $x \in X \setminus \{x_0\}$ . The converse is true, as shown by  $f(x) = x^2 e^x$ ,  $x \in \mathbb{R}$ ,  $x_0 = 0$ .

The generalized version of the Borwein-Preiss variational principle due to Loewen and Wang is the following.

**Theorem 1.5** *Let  $f : X \rightarrow \mathbb{R} \cup \{+\infty\}$  be a lower semi-continuous function. Assume that  $x_0 \in X$  and  $\varepsilon > 0$  satisfy*

$$f(x_0) < \varepsilon + \inf_X f.$$

*Let  $(\mu_n)_{n \geq 0}$  be a decreasing sequence in  $(0, 1)$  such that the series  $\sum_{n=0}^{\infty} \mu_n$  is convergent. Then for any continuous function  $\rho$  satisfying (1.2), there exists a sequence  $(e_n)_{n \geq 0}$  in  $X$  converging to  $e$  such that*

- (i)  $\rho(x_0 - e) < 1$ ;
- (ii)  $f(e) + \varepsilon\rho_\infty(e) \leq f(x_0)$ ;
- (iii)  $e$  is a strong minimizer of  $f + \varepsilon\rho_\infty$ . In particular,  $e$  is a strict minimizer of  $f + \varepsilon\rho_\infty$ , that is,

$$f(e) + \varepsilon\rho_\infty(e) < f(x) + \varepsilon\rho_\infty(x) \quad \text{for all } x \in X \setminus \{e\}.$$

*Proof* Define the sequence  $(f_n)_{n \geq 0}$  such that  $f_0 = f$  and for any  $n \geq 0$ ,

$$f_{n+1}(x) := f_n(x) + \varepsilon\rho_n(x - e_n).$$

Then  $f_n \leq f_{n+1}$  and  $f_n$  is lower semi-continuous.

Set  $e_0 = x_0$ . We observe that for any  $n \geq 0$ ,

$$\inf_X f_{n+1} \leq f_{n+1}(e_n) = f_n(e_n). \quad (1.4)$$

If this inequality is strict, then there exists  $e_{n+1} \in X$  such that

$$f_{n+1}(e_{n+1}) \leq \frac{\mu_{n+1}}{2} f_n(e_n) + \left(1 - \frac{\mu_{n+1}}{2}\right) \inf_X f_{n+1} \leq f_n(e_n). \quad (1.5)$$

If equality holds in relation (1.4) then (1.5) also holds, but for  $e_{n+1}$  replaced with  $e_n$ . Consequently, there exists a sequence  $(e_n)_{n \geq 0}$  in  $X$  such that relation (1.5) holds true.

Set

$$D_n := \left\{ x \in X; f_{n+1}(x) \leq f_{n+1}(e_{n+1}) + \frac{\varepsilon\mu_n}{2} \right\}.$$

Then  $D_n$  is not empty, since  $e_{n+1} \in D_n$ . By the lower semi-continuity of functions  $f_n$  we also deduce that  $D_n$  is a closed set. Since  $\mu_{n+1} \in (0, 1)$ , relation (1.5) implies

$$\begin{aligned} f_{n+1}(e_{n+1}) - \inf_X f_{n+1} &\leq \frac{\mu_{n+1}}{2} [f_n(e_n) - \inf_X f_{n+1}] \\ &\leq f_n(e_n) - \inf_X f_n. \end{aligned} \quad (1.6)$$

We also observe that

$$f_0(e_0) - \inf_X f_0 = f(x_0) - \inf_X f < \varepsilon.$$

Next, we prove that

$$\text{the sequence } (D_n)_{n \geq 0} \text{ is decreasing} \quad (1.7)$$

and

$$\text{diam}(D_n) \rightarrow 0 \quad \text{as } n \rightarrow \infty. \quad (1.8)$$

In order to prove (1.7), assume that  $x \in D_n$ ,  $n \geq 1$ . Since the sequence  $(\mu_n)_{n \geq 0}$  is decreasing, relation (1.5) implies

$$f_n(x) \leq f_{n+1}(x) \leq f_{n+1}(e_{n+1}) + \frac{\varepsilon \mu_n}{2} \leq f_n(e_n) + \frac{\varepsilon \mu_{n-1}}{2},$$

hence  $x \in D_{n-1}$ .

Since  $f_n \geq f_{n-1}$ , relations (1.5) and (1.6) imply

$$\begin{aligned} f_n(e_n) - \inf_X f_n &\leq \frac{\mu_n}{2} [f_{n-1}(e_{n-1}) - \inf_X f_n] \\ &\leq \frac{\mu_n}{2} [f_{n-1}(e_{n-1}) - \inf_X f_{n-1}] < \frac{\varepsilon \mu_n}{2}. \end{aligned} \quad (1.9)$$

For any  $x \in D_n$ , combining relation (1.9) and the definitions of  $f_{n+1}$  and  $D_n$  we obtain

$$\begin{aligned} \varepsilon \mu_n \rho((n+1)(x - e_n)) &\leq f_{n+1}(e_{n+1}) - f_n(x) + \frac{\varepsilon \mu_n}{2} \\ &\leq f_{n+1}(e_{n+1}) - \inf_X f_n + \frac{\varepsilon \mu_n}{2} \\ &\leq f_n(e_n) - \inf_X f_n + \frac{\varepsilon \mu_n}{2} < \varepsilon \mu_n. \end{aligned} \quad (1.10)$$

Therefore  $\rho((n+1)(x - e_n)) < 1$ . So, by (1.2),

$$(n+1)\|x - e_n\| \leq \rho_M,$$

which shows that  $\text{diam}(D_n) \leq 2\rho_M/(n+1)$ . This implies (1.8).

Since  $D_n$  is a closed set for any  $n \geq 1$ , then (1.7) and (1.8) imply that  $\bigcap_{n=1}^{\infty} D_n$  contains a single point, denoted by  $e$ . Then  $e_n \rightarrow e$  as  $n \rightarrow \infty$ . Thus, using  $\rho((n+1)(x - e_n)) < 1$  for all  $n \geq 0$  and  $x \in X$ , we deduce that  $\rho(x_0 - e) < 1$ .

Since the sequence  $(f_n(e_n))_{n \geq 0}$  is nonincreasing and  $f_0(e_0) = f(x_0)$  it follows that, in order to prove (ii), it is enough to deduce that

$$f(e) + \varepsilon \rho_{\infty}(e) \leq f_n(e_n). \quad (1.11)$$

For this purpose we define the nonempty closed sets

$$C_n := \{x \in X; f_{n+1}(x) \leq f_{n+1}(e_{n+1})\}.$$

Since  $f_n \leq f_{n+1}$  and  $f_n(e_n) \geq f_{n+1}(e_{n+1})$  for all  $n$ , it follows that the sequence  $(C_n)_{n \geq 0}$  is nested and  $C_n \subset D_n$  for all  $n$ . Therefore  $\bigcap_{n=0}^{\infty} C_n = \{e\}$  and

$$f_m(e) \leq f_m(e_m) \leq f_n(e_n) \leq f(x_0) \quad \text{provided that } m > n. \quad (1.12)$$

Taking  $m \rightarrow \infty$  we obtain (1.11).

It remains to argue that  $e$  is a strong minimizer of  $f_\varepsilon := f + \varepsilon\rho_\infty$ . Since

$$f_\varepsilon(x) \leq \inf_X f_\varepsilon + \frac{\varepsilon\mu_n}{2},$$

relation (1.12) yields

$$f_{n+1}(x) \leq f_\varepsilon(x) \leq f_\varepsilon(e) + \frac{\varepsilon\mu_n}{2} \leq f_{n+1}(e_{n+1}) + \frac{\varepsilon\mu_n}{2}.$$

Setting

$$A_n := \left\{ x \in X; f_\varepsilon(x) \leq \frac{\varepsilon\mu_n}{2} + \inf_X f \right\},$$

the above relation shows that  $A_n \subset D_n$ . So, by (1.8), we deduce that  $\text{diam}(A_n) \rightarrow 0$  as  $n \rightarrow \infty$  which shows that  $e$  is a strong minimizer of  $f_\varepsilon := f + \varepsilon\rho_\infty$ .  $\square$

Assume that  $p \geq 1$  and  $\lambda > 0$ . Taking

$$\rho(x) = \frac{\|x\|^p}{\lambda^p} \quad \text{and} \quad \mu_n = \frac{1}{2^{n+1}(n+1)}$$

we obtain the initial smooth version of the Borwein-Preiss variational principle. Roughly speaking, it asserts that the Lipschitz perturbations obtained in Ekeland's variational principle can be replaced by *superlinear perturbations* in a certain class of admissible functions.

**Theorem 1.6** *Given  $f : X \rightarrow \mathbb{R} \cup \{+\infty\}$  lower semi-continuous function,  $x_0 \in X$ ,  $\varepsilon > 0$ ,  $\lambda > 0$ , and  $p \geq 1$ , suppose*

$$f(x_0) < \varepsilon + \inf_X f.$$

*Then there exists a sequence  $(\mu_n)_{n \geq 0}$  with  $\mu_n \geq 0$ ,  $\sum_{n=0}^{\infty} \mu_n = 1$ , and a point  $e$  in  $X$ , expressible as the limit of some sequence  $(e_n)$ , such that for all  $x \in X$ ,*

$$f(x) + \frac{\varepsilon}{\lambda^p} \sum_{n=0}^{\infty} \mu_n \|x - e_n\|^p \geq f(e) + \frac{\varepsilon}{\lambda^p} \sum_{n=0}^{\infty} \mu_n \|e - e_n\|^p.$$

*Moreover,  $\|x_0 - e\| < \lambda$  and  $f(e) \leq \varepsilon + \inf_X f$ .*

We have seen in Corollary 1.2 of Ekeland's variational principle that any smooth bounded from below functional on a Banach space admits a sequence of "almost critical points". The next consequence of Borwein-Preiss' variational principle asserts that, in the framework of Hilbert spaces, such a functional admits a sequence of *stable* "almost critical points".

**Corollary 1.3** *Let  $f$  be a real-valued  $C^2$ -functional that is bounded from below on a Hilbert space  $X$ . Assume that  $(u_n)$  is a minimizing sequence of  $f$ . Then there exists a minimizing sequence  $(v_n)$  of  $f$  such that the following properties hold true:*

- (i)  $\lim_{n \rightarrow \infty} \|u_n - v_n\| = 0$ ;
- (ii)  $\lim_{n \rightarrow \infty} \|f'(v_n)\| = 0$ ;
- (iii)  $\liminf_{n \rightarrow \infty} \langle f''(v_n)w, w \rangle \geq 0$  for any  $w \in X$ .

### 1.3 Minimax principles

In this section we are interested in some powerful techniques for finding solutions of some classes of stationary nonlinear boundary value problems. These solutions are viewed as critical points of a natural functional, often called the *energy* associated to the system. The critical points obtained in this section by means of topological techniques are generally *nonstable* critical points which are neither maxima nor minima of the energy functional.

#### 1.3.1 Mountain pass type results

In many nonlinear problems we are interested in finding solutions as stationary points of some associated “energy” functionals. Often such a mapping is unbounded from above and below, so that it has no maximum or minimum. This forces us to look for saddle points, which are obtained by minimax arguments. In such a case one maximizes a functional  $f$  over a closed set  $A$  belonging to some family  $\Gamma$  of sets and then one minimizes with respect to the set  $A$  in the family. Thus we define

$$c = \inf_{A \in \Gamma} \sup_{u \in A} f(u) \tag{1.13}$$

and one tries to prove, under various hypotheses, that this number  $c$  is a critical value of  $f$ , hence there is a point  $u$  such that  $f(u) = c$  and  $f'(u) = 0$ . Indeed, it seems intuitively obvious that  $c$  defined in (1.13) is a critical value of  $f$ . However, this is not true in general, as showed by the following example in the plane: let  $f(x, y) = x^2 - (x - 1)^3 y^2$ . Then  $(0, 0)$  is the only critical point of  $f$  but  $c$  is not a critical value. Indeed, looking for sets  $A$  lying in a small neighborhood of the origin, then  $c > 0$ . This example shows that the heart of the matter is to find appropriate conditions on  $f$  and on the family  $\Gamma$ .

One of the most important minimax results is the so-called *mountain pass theorem* whose geometrical interpretation will be roughly described in the sequel. Denote by  $f$  the function which measures the altitude of a mountain terrain and assume that there are two points in the horizontal plane  $L_1$  and  $L_2$ , representing the coordinates of two locations such that  $f(L_1)$  and  $f(L_2)$  are the deepest points of two separated valleys. Roughly speaking, our aim is to walk along an optimal path on the mountain from the point  $(L_1, f(L_1))$  to  $(L_2, f(L_2))$  spending the least amount of energy by passing the mountain ridge between the two valleys. Walking on a path  $(\gamma, f(\gamma))$  from  $(L_1, f(L_1))$  to  $(L_2, f(L_2))$  such that the maximal altitude along  $\gamma$  is the smallest among all such continuous paths connecting  $(L_1, f(L_1))$  and  $(L_2, f(L_2))$ , we reach a point  $L$  on  $\gamma$  passing the ridge of the mountain which is called a *mountain pass point*. As pointed out by Brezis and Browder [48], the mountain pass theorem “extends ideas already present in Poincaré and Birkhoff”.

In the sequel, we give a first formulation of the mountain pass theorem.

**Theorem 1.7** [Mountain pass theorem; positive altitude] *Let  $X$  be a Banach space,  $f : X \rightarrow \mathbb{R}$  be a function of class  $C^1$  such that*

$$\inf_{\|u - e_0\| = \rho} f(u) \geq \alpha > \max\{f(e_0), f(e_1)\}$$

*for some  $\alpha \in \mathbb{R}$  and  $e_0 \neq e_1 \in X$  with  $0 < \rho < \|e_0 - e_1\|$ . If  $f$  satisfies the  $(PS)_c$ -condition at level*

$$c = \inf_{\gamma \in \Gamma} \max_{t \in [0,1]} f(\gamma(t)),$$

*where*

$$\Gamma = \{\gamma \in C([0, 1], X) : \gamma(0) = e_0, \gamma(1) = e_1\},$$

*then  $c$  is a critical value of  $f$  with  $c \geq \alpha$ .*

There are two different ways to prove Theorem 1.7; namely, via Ekeland’s variational principle, or by using a sort of deformation lemma. We present its proof by means of the latter argument. We refer to Figure 1.3.1 for a geometric illustration of Theorem 1.7.

*Proof of Theorem 1.7.* We may assume that  $e_0 = 0$  and let  $e := e_1$ . Since  $f(u) \geq \alpha$  for every  $u \in X$  with  $\|u\| = \rho$  and  $\rho < \|e\|$ , the definition of the number  $c$  shows that  $\alpha \leq c$ .

It remains to prove that  $c$  is a critical value of  $f$ . Arguing by contradiction, assume  $K_c = \emptyset$ . Thus, on account of Remark D.1, we may choose  $\mathcal{O} = \emptyset$  in Theorem D.1, see Appendix D, and  $\bar{\varepsilon} > 0$  such that




Figure 1.2 Mountain pass landscape between “villages”  $e_0$  and  $e_1$ .

$\bar{\varepsilon} < \min\{\alpha - f(0), \alpha - f(e)\}$ . Consequently, there exist  $\varepsilon > 0$  and a continuous map  $\eta : X \times [0, 1] \rightarrow X$  verifying the properties (i)-(iv) from Theorem D.1.

From the definition of the number  $c$ , there exists  $\gamma \in \Gamma$  such that

$$\max_{t \in [0, 1]} f(\gamma(t)) \leq c + \varepsilon. \quad (1.14)$$

Let  $\gamma_1 : [0, 1] \rightarrow X$  defined by  $\gamma_1(t) = \eta(\gamma(t), 1)$ ,  $\forall t \in [0, 1]$ .

We prove that  $\gamma_1 \in \Gamma$ . The choice of  $\bar{\varepsilon} > 0$  gives that  $\max\{f(0), f(e)\} < \alpha - \bar{\varepsilon} \leq c - \bar{\varepsilon}$ , thus  $0, e \notin f^{-1}(]c - \bar{\varepsilon}, c + \bar{\varepsilon}[)$ . Consequently, due to Theorem D.1 (iii), we have  $\gamma_1(0) = \eta(\gamma(0), 1) = \eta(0, 1) = 0$  and  $\gamma_1(1) = \eta(\gamma(1), 1) = \eta(e, 1) = e$ . Therefore,  $\gamma_1 \in \Gamma$ .

Note that (1.14) means that  $\gamma(t) \in f^{c+\varepsilon} = \{u \in X : f(u) \leq c + \varepsilon\}$  for all  $t \in [0, 1]$ . Then by Theorem D.1 (iv) and the definition of  $\gamma_1$  we have

$$c \leq \max_{t \in [0, 1]} f(\gamma_1(t)) = \max_{t \in [0, 1]} f(\eta(\gamma(t), 1)) \leq c - \varepsilon,$$

contradiction.

**Remark 1.1** Using Theorem D.1, we can provide an alternative proof to Theorem 1.4. Indeed, if we suppose that  $K_c = \emptyset$ , with  $c = \inf_X f > -\infty$ , one may deform continuously the level set  $f^{c+\varepsilon}$  (for  $\varepsilon > 0$  small

enough) into a subset of  $f^{c-\varepsilon}$ , see Theorem D.1 (iv). But,  $f^{c-\varepsilon} = \emptyset$ , contradiction.

**Remark 1.2** Note that the choice of  $0 < \bar{\varepsilon} < \min\{\alpha - f(0), \alpha - f(e)\}$  with  $\inf_{\|u\|=\rho} f(u) \geq \alpha$  is crucial in the proof of Theorem 1.7. Actually, it means that the ridge of the mountain between the two valleys has a positive altitude. However, a more involved proof allows us to handle the so-called “zero altitude” case. More precisely, the following result holds.

**Theorem 1.8** [Mountain pass theorem; zero altitude] *Let  $X$  be a Banach space,  $f : X \rightarrow \mathbb{R}$  be a function of class  $C^1$  such that*

$$\inf_{\|u-e_0\|=\rho} f(u) \geq \max\{f(e_0), f(e_1)\}$$

for some  $e_0 \neq e_1 \in X$  with  $0 < \rho < \|e_0 - e_1\|$ . If  $f$  satisfies the  $(PS)_c$ -condition at level

$$c = \inf_{\gamma \in \Gamma} \max_{t \in [0,1]} f(\gamma(t)),$$

where

$$\Gamma = \{\gamma \in C([0, 1], X) : \gamma(0) = e_0, \gamma(1) = e_1\},$$

then  $c$  is a critical value of  $f$  with  $c \geq \max\{f(e_0), f(e_1)\}$ .

**Remark 1.3** In both Theorems 1.7 and 1.8 the Palais-Smale condition may be replaced by a weaker one, called the *Cerami compactness condition*. More precisely, a function  $f \in C^1(X, \mathbb{R})$  satisfies the *Cerami condition at level  $c \in \mathbb{R}$*  (shortly,  $(C)_c$ -condition) if every sequence  $\{u_n\} \subset X$  such that  $\lim_{n \rightarrow \infty} f(u_n) = c$  and  $\lim_{n \rightarrow \infty} (1 + \|u_n\|)\|f'(u_n)\| = 0$ , possesses a convergent subsequence.

### 1.3.2 Minimax results via Ljusternik-Schnirelmann category

A very useful method to find multiple critical points for a given functional is related to the notion of *Ljusternik-Schnirelmann category*. In the case of variational problems on finite dimensional manifolds, this notion is useful to find a lower bound for the number of critical points in terms of topological invariants. In this subsection we present a general form of this approach. We first present some basic properties of Finsler-Banach manifolds.

Let  $M$  be a Banach manifold of class  $C^1$  and let  $TM$  be the usual

tangent bundle,  $T_pM$  the tangent space at the point  $p \in M$ . A Finsler structure on the Banach manifold  $M$  can be introduced as follows.

**Definition 1.1** A Finsler structure on  $TM$  is a continuous function  $\|\cdot\| : TM \rightarrow [0, \infty)$  such that

- (a) for each  $p \in M$  the restriction  $\|\cdot\|_p = \|\cdot\|_{T_pM}$  is a norm on  $T_pM$ .
- (b) for each  $p_0 \in M$  and  $k > 1$ , there is a neighborhood  $U \subset M$  of  $p_0$  such that

$$\frac{1}{k}\|\cdot\|_p \leq \|\cdot\|_{p_0} \leq k\|\cdot\|_p$$

for all  $p \in U$ .

The Banach manifold  $M$  of class  $C^1$  endowed with a Finsler structure is called Banach-Finsler manifold  $M$  of class  $C^1$ .

**Definition 1.2** Let  $M$  be a Banach-Finsler manifold of class  $C^1$  and  $\sigma : [a, b] \rightarrow M$  a  $C^1$ -path. The length of  $\sigma$  is defined by  $l(\sigma) = \int_a^b \|\sigma'(t)\| dt$ . If  $p$  and  $q$  are in the same component of  $M$  we define the distance from  $p$  to  $q$  by

$$\rho(p, q) = \inf\{l(\sigma) : \sigma \text{ is a } C^1 \text{ - path from } p \text{ to } q\}.$$

**Definition 1.3** A Banach-Finsler manifold  $M$  of class  $C^1$  is said to be complete if the metric space  $(M, \rho)$  is complete.

**Theorem 1.9** If  $M$  is a Banach-Finsler manifold of class  $C^1$  then the function  $\rho : M \times M \rightarrow \mathbb{R}$  (from Definition 1.2) is a metric function for each component of  $M$ . Moreover, this metric is consistent with the topology of  $M$ .

**Theorem 1.10** If  $M$  is a Banach-Finsler manifold of class  $C^1$  and  $N$  is a  $C^1$  submanifold of  $M$ , then  $\|\cdot\|_{T(N)}$  is a Finsler structure for  $N$  (called the Finsler structure induced from  $M$ ). If  $M$  is complete and  $N$  is a closed  $C^1$  submanifold of  $M$ , then  $N$  is a complete Banach-Finsler manifold in the induced Finsler structure.

On a Banach-Finsler manifold  $M$  the cotangent bundle  $TM^*$  has a dual Finsler structure defined by

$$\|y\| = \sup\{\langle y, v \rangle : v \in T_pM, \|v\|_p = 1\},$$

where  $y \in T_pM^*$  and  $\langle \cdot, \cdot \rangle$  denotes the duality pairing between  $T_pM^*$  and  $T_pM$ . Consequently, for a functional  $f : M \rightarrow \mathbb{R}$  of class  $C^1$ , the

application  $p \mapsto \|f'(p)\|$  is well-defined and continuous. These facts allow to introduce the following definition.

**Definition 1.3** Let  $M$  be a Banach-Finsler manifold of class  $C^1$ .

(a) A function  $f \in C^1(M, \mathbb{R})$  satisfies the Palais-Smale condition at level  $c \in \mathbb{R}$  (shortly,  $(PS)_c$ -condition) if every sequence  $\{u_n\} \subset M$  such that  $\lim_{n \rightarrow \infty} f(u_n) = c$  and  $\lim_{n \rightarrow \infty} \|f'(u_n)\| = 0$ , possesses a convergent subsequence.

(b) A function  $f \in C^1(M, \mathbb{R})$  satisfies the Palais-Smale condition (shortly,  $(PS)$ -condition) if it satisfies the Palais-Smale condition at every level  $c \in \mathbb{R}$ .

Let  $M$  be a Banach-Finsler manifold of class  $C^1$ . For  $h \geq 1$ , we introduce the set

$$\Gamma_h = \{A \subseteq M : \text{cat}_M(A) \geq h, A \text{ compact}\}.$$

Here,  $\text{cat}_M(A)$  denotes the *Ljusternik-Schnirelmann category* of the set  $A$  relative to  $M$ , see Appendix C.

If  $h \leq \text{cat}_M(M) = \text{cat}(M)$ , then  $\Gamma_h \neq \emptyset$ . Moreover, if  $\text{cat}(M) = +\infty$ , we may take any positive integer  $h$  in the above definition. If  $f : M \rightarrow \mathbb{R}$  is a functional of class  $C^1$ , we denote by  $K$  the set of all critical points of  $f$  on  $M$ , that is,  $K = \{u \in M : f'(u) = 0\}$  and  $K_c = K \cap f^{-1}(c)$ . Finally, for every  $h \leq \text{cat}(M)$  we define the number

$$c_h = \inf_{A \in \Gamma_h} \max_{u \in A} f(u).$$

**Proposition 1.1** Let  $f : X \rightarrow \mathbb{R}$  be a functional of class  $C^1$ . Using the above notations, we have

- a)  $c_1 = \inf_M f$ .
- b)  $c_h \leq c_{h+1}$ .
- c)  $c_h < \infty$  for every  $h \leq \text{cat}(M)$ .
- d) If  $c := c_h = c_{h+m-1}$  for some  $h, m \geq 1$ , and  $f$  satisfies the  $(PS)_c$ -condition, then  $\text{cat}_M(K_c) \geq m$ . In particular,  $K_c \neq \emptyset$ .

*Proof* Properties a)-c) are obvious. To prove d), we argue by contradiction. Thus, we assume that  $\text{cat}_M(K_c) \leq m-1$ . Since  $M$  is an ANR, there exists a neighborhood  $\mathcal{O}$  of  $K_c$  such that  $\text{cat}_M(\overline{\mathcal{O}}) = \text{cat}_M(K_c) \leq m-1$ . By using Theorem D.1, there exists a continuous map  $\eta : X \times [0, 1] \rightarrow X$  and  $\varepsilon > 0$  such that  $\eta(f^{c+\varepsilon} \setminus \mathcal{O}, 1) \subset f^{c-\varepsilon}$  and  $\eta(u, 0) = u$  for all  $u \in X$ . Let  $A_1 \in \Gamma_{h+m-1}$  such that  $\max_{u \in A_1} f(u) \leq c + \varepsilon$ . Considering the set  $A_2 = \overline{A_1} \setminus \mathcal{O}$ , we obtain that

$$\text{cat}_M(A_2) \geq \text{cat}_M(A_1) - \text{cat}_M(\overline{\mathcal{O}}) \geq h + m - 1 - (m - 1) = h.$$

Therefore,  $A_2 \in \Gamma_h$ . Moreover,  $\max_{u \in A_2} f(u) \leq \max_{u \in A_1} f(u) \leq c + \varepsilon$  and  $A_2 \cap \mathcal{O} = \emptyset$ . In conclusion, we have  $A_2 \subset f^{c+\varepsilon} \setminus \mathcal{O}$ . Due to Theorem D.1 (iv),  $\eta(A_2, 1) \subset f^{c-\varepsilon}$ . Moreover,  $\text{cat}_M(\eta(A_2, 1)) \geq \text{cat}_M(A_2) \geq h$ , thus  $\eta(A_2, 1) \in \Gamma_h$ . But  $c = \inf_{A \in \Gamma_h} \max_{u \in A} f(u) \leq \max f(\eta(A_2, 1)) \leq c - \varepsilon$ , contradiction.  $\square$

The main result of the present subsection is the following theorem.

**Theorem 1.11** *Let  $M$  be a complete Banach-Finsler manifold of class  $C^1$ , and  $f : M \rightarrow \mathbb{R}$  a functional of class  $C^1$  which is bounded from below on  $M$ . If  $f$  satisfies the (PS)-condition then  $f$  has at least  $\text{cat}(M)$  critical points.*

*Proof* Since  $f$  is bounded from below, every  $c_h$  is finite,  $h = 1, \dots, \text{cat}(M)$ , see Proposition 1.1 c). To prove the statement, it is enough to show that

$$\text{card}(K \cap f^{c_h}) \geq h, \quad h = 1, \dots, \text{cat}(M). \quad (1.15)$$

We proceed by induction. For  $h = 1$ , relation (1.15) is obvious since the global minimum belongs to  $K$ . Now, we assume that (1.15) holds for  $h = 1, \dots, k$ . We will prove that (1.15) also holds for  $k + 1$ . There are two cases:

I)  $c_k \neq c_{k+1}$ . Due to Proposition 1.1 d), we have that  $K_{c_{k+1}} \neq \emptyset$ . Consequently, each element of the set  $K_{c_{k+1}}$  clearly differs from those of the set  $K \cap f^{c_k}$ . Therefore,  $K \cap f^{c_{k+1}}$  contains at least  $k + 1$  points.

II)  $c_k = c_{k+1} := c$ . Let  $m$  be the least positive integer such that  $c_m = c_{k+1}$ . Due to Proposition 1.1 d), we have  $\text{cat}_M(K_c) \geq k + 1 - m + 1 = k - m + 2$ . In particular,  $\text{card}(K_c) = \text{card}(K_{c_{k+1}}) \geq k - m + 2$ . Two distinguish again two subcases:

IIa)  $m = 1$ . The claim easily follows.

IIb)  $m > 1$ . Since  $m \leq k$ , then  $\text{card}(K \cap f^{c_{m-1}}) \geq m - 1$ . Consequently,

$$\begin{aligned} \text{card}(K \cap f^{c_{k+1}}) &\geq \text{card}(K \cap f^{c_{m-1}}) + \text{card}(K_{c_{k+1}}) \\ &\geq (m - 1) + (k - m + 2) \\ &= k + 1, \end{aligned}$$

which complete the proof of (1.15), thus the theorem.  $\square$

Let  $X$  be an infinite dimensional, separable real Banach space and  $S = \{u \in X : \|u\| = 1\}$  its unit sphere. Important questions appear when the study of some classes of nonlinear elliptic problems reduce to finding critical points of a given functional on  $S$ . Note that  $\text{cat}(S) = 1$ ;

therefore, Theorem 1.11 does not give multiple critical points on  $S$ . However, exploiting the symmetric property of  $S$ , by means of the *Krasnoselski genus*, see Appendix C, a multiplicity result may be given, similar to Theorem 1.11. In order to state this result, we put ourselves to a general framework.

Let  $X$  be a real Banach space, and we denote the family of sets

$$\mathcal{A} = \{A \subset X \setminus \{0\} : A = -A, A \text{ is closed}\}.$$

Assume that  $M = G^{-1}(0) \in \mathcal{A}$  is a submanifold of  $X$ , where  $G : X \rightarrow \mathbb{R}$  is of class  $C^1$  with  $G'(u) \neq 0$  for every  $u \in M$ .

For any  $h \leq \gamma_0(M) := \sup\{\gamma(K) : K \subseteq M, K \in \mathcal{A}, K \text{ compact}\}$ , we introduce the set

$$\mathcal{A}_h = \{A \subseteq M : A \in \mathcal{A}, \gamma(A) \geq h, A \text{ compact}\}.$$

Here,  $\gamma(A)$  denotes the Krasnoselski genus of the set  $A$ , see Appendix C. Finally, if  $f : X \rightarrow \mathbb{R}$  is an even function, we denote

$$\tilde{c}_h = \inf_{A \in \mathcal{A}_h} \max_{u \in A} f(u).$$

**Proposition 1.2** *Let  $f : X \rightarrow \mathbb{R}$  be an even functional of class  $C^1$ .*

*Using the above notations, we have*

- a)  $\tilde{c}_1 > -\infty$  whenever  $f$  is bounded from below on  $M$ .
- b)  $\tilde{c}_h \leq \tilde{c}_{h+1}$ .
- c)  $\tilde{c}_h < \infty$  for every  $h \leq \gamma_0(M)$ .
- d) If  $c := \tilde{c}_h = \tilde{c}_{h+m-1}$  for some  $h, m \geq 1$ , and  $f$  satisfies the  $(PS)_c$ -condition on  $M$ , then  $\gamma(K_c) \geq m$ . In particular,  $K_c \neq \emptyset$ .

The following result may be stated.

**Theorem 1.12** *Let  $M = G^{-1}(0) \subset X$  be a submanifold of a real Banach space, where  $G : X \rightarrow \mathbb{R}$  is of class  $C^1$  with  $G'(u) \neq 0$  for every  $u \in M$ . Assume that  $M \in \mathcal{A}$  and let  $f : X \rightarrow \mathbb{R}$  be a functional of class  $C^1$  which is bounded from below on  $M$  and even. If  $\gamma(M) = +\infty$  and  $f$  satisfies the  $(PS)$ -condition on  $M$  then  $f$  has infinitely many critical points  $\{u_k\} \subset M$  with  $\lim_{k \rightarrow \infty} f(u_k) = \sup_M f$ .*

We conclude this section by stating the following version of the Lagrange multiplier rule.

**Theorem 1.13** *Let  $(X, \|\cdot\|)$  be a real Banach space,  $\Omega \subset X$  be an open subset,  $f : \Omega \rightarrow \mathbb{R}$  be a differentiable function, and  $G : \Omega \rightarrow \mathbb{R}^n$  be*

of class  $C^1$ . Let also  $a \in \mathbb{R}^n$  be such that  $M = G^{-1}(a)$  is not empty. If  $x_0 \in M$  is a solution of the minimization problem

$$f(x_0) = \min_{x \in M} f(x)$$

and if  $DG(x_0)$  is surjective, then there exists  $(\lambda_1, \dots, \lambda_n) \in \mathbb{R}^n$  such that

$$Df(x_0) = \sum_{i=1}^n \lambda_i DG_i(x_0),$$

where  $G = (G_1, \dots, G_n)$ .

## 1.4 Ricceri's variational results

The main part of the present section is dedicated to Ricceri's recent multiplicity results. First, several three critical points results with one/two parameter are stated. In the second subsection a general variational principle of Ricceri is presented. Finally, in the last subsection a new kind of multiplicity result of Ricceri is given which guarantees the existence of  $k \geq 2$  distinct critical points for a fixed functional.

### 1.4.1 Three critical point results

It is a simple exercise to show that a function  $f : \mathbb{R} \rightarrow \mathbb{R}$  of class  $C^1$  having two local minima has necessarily a third critical point. However, once we are dealing with functions defined on a multi-dimensional space, the problem becomes much deeper. The well-known form of the three critical point theorem is due to Pucci-Serrin [250, 251] and it can be formulated as follows.

**Theorem 1.14** *A function  $f : X \rightarrow \mathbb{R}$  of class  $C^1$  satisfies the (PS)-condition and it has two local minima. Then  $f$  has at least three distinct critical points.*

*Proof* Without loss of generality we may assume that 0 and  $e \in X \setminus \{0\}$  are the two local minima of  $f$  and  $f(e) \leq f(0) = 0$ . We face two cases:

I) If there exist constants  $\alpha, \rho > 0$  such that  $\|e\| > \rho$  and  $\inf_{\|u\|=\rho} f(u) \geq \alpha$ , then the existence of a critical point of  $f$  at a minimax level  $c$  with  $c \geq \alpha$  is guaranteed by Theorem 1.7. Consequently, this critical point certainly differs from 0 and  $e$ .

II) Assume now that such constants do not exist as in I). Since 0 is

a local minima of  $f$ , we may choose  $r < \|e\|$  such that  $f(u) \geq 0$  for every  $u \in X$  with  $\|u\| \leq r$ . We apply Ekeland's variational principle, see Theorem 1.3 with  $\varepsilon = \frac{1}{n^2}$ ,  $\lambda = n$  and  $f$  restricted to the closed ball  $B[0, r] = \{u \in X : \|u\| \leq r\}$ . Let us fix  $0 < \rho < r$  small enough. Since  $\inf_{\|u\|=\rho} f(u) = 0$ , there exist  $z_n \in X$  with  $\|z_n\| = \rho$  and  $u_n \in B[0, r]$  such that

$$0 \leq f(u_n) \leq f(z_n) \leq \frac{1}{n^2}, \quad \|u_n - z_n\| \leq \frac{1}{n}$$

and

$$f(w) - f(u_n) \geq -\frac{1}{n}\|w - u_n\|$$

for all  $w \in B[0, r]$ . Since  $\rho < r$ , for  $n$  enough large, we have  $\|u_n\| < r$ . Fix  $v \in X$  arbitrarily and let  $w = u_n + tv$ . If  $t > 0$  is small enough then  $w \in B[0, r]$ . Therefore, with this choice, the last inequality gives that  $\|f'(u_n)\| \leq \frac{1}{n}$ . Since  $0 \leq f(u_n) \leq \frac{1}{n^2}$  and  $f$  satisfies the  $(PS)$ -condition, we may assume that  $u_n \rightarrow u \in X$  which is a critical point of  $f$ . Since  $|\|u_n\| - \rho| = |\|u_n\| - \|z_n\|| \leq \frac{1}{n}$ , we actually have that  $\|u\| = \rho$ , thus  $0 \neq u \neq e$ .  $\square$

Let  $f : X \times I \rightarrow \mathbb{R}$  be a function such that  $f(\cdot, \lambda)$  is of class  $C^1$  for every  $\lambda \in I \subset \mathbb{R}$ . In view of Theorem 1.14, the main problem Ricceri dealt with is the *stability* of the three critical points of  $f(\cdot, \lambda)$  with respect to the parameters  $\lambda \in I$ . Ricceri's main three critical point theorem is the following.

**Theorem 1.15** *Let  $X$  be a separable and reflexive real Banach space,  $I \subseteq \mathbb{R}$  be an interval, and  $f : X \times I \rightarrow \mathbb{R}$  be a function satisfying the following conditions:*

- ( $\alpha_1$ ) for each  $u \in X$ , the function  $f(u, \cdot)$  is continuous and concave;
- ( $\beta_1$ ) for each  $\lambda \in I$ , the function  $f(\cdot, \lambda)$  is sequentially weakly lower semi-continuous and Gâteaux differentiable, and  $\lim_{\|u\| \rightarrow +\infty} f(u, \lambda) = +\infty$ ;
- ( $\gamma_1$ ) there exists a continuous concave function  $h : I \rightarrow \mathbb{R}$  such that

$$\sup_{\lambda \in I} \inf_{u \in X} (f(u, \lambda) + h(\lambda)) < \inf_{u \in X} \sup_{\lambda \in I} (f(u, \lambda) + h(\lambda)).$$

Then, there exist an open interval  $J \subseteq I$  and a positive real number  $\rho$ , such that, for each  $\lambda \in J$ , the equation

$$f'_u(u, \lambda) = 0$$

has at least two solutions in  $X$  whose norms are less than  $\rho$ . If, in

addition, the function  $f$  is (strongly) continuous in  $X \times I$ , and, for each  $\lambda \in I$ , the function  $f(\cdot, \lambda)$  is of class  $C^1$  and satisfies the (PS)-condition, then the above conclusion holds with "three" instead of "two".

The proof of Theorem 1.15 is quite involved combining deep arguments from nonlinear analysis as a topological minimax result of Saint Raymond [268], a general selector result of Kuratowski and Ryll-Nardzewski [185], [147], and the mountain pass theorem. Instead of its proof, we give a useful consequence of Theorem 1.15.

**Corollary 1.4** *Let  $X$  be a separable and reflexive real Banach space,  $\Phi : X \rightarrow \mathbb{R}$  a continuously Gâteaux differentiable and sequentially weakly lower semi-continuous functional whose Gâteaux derivative admits a continuous inverse on  $X^*$ ,  $\Psi : X \rightarrow \mathbb{R}$  a continuously Gâteaux differentiable functional whose Gâteaux derivative is compact, and  $I \subseteq \mathbb{R}$  an interval. Assume that*

$$\lim_{\|u\| \rightarrow +\infty} (\Phi(u) - \lambda\Psi(u)) = +\infty$$

for all  $\lambda \in I$ , and that there exists a continuous concave function  $h : I \rightarrow \mathbb{R}$  such that

$$\sup_{\lambda \in I} \inf_{u \in X} (\Phi(u) - \lambda\Psi(u) + h(\lambda)) < \inf_{u \in X} \sup_{\lambda \in I} (\Phi(u) - \lambda\Psi(u) + h(\lambda)).$$

Then, there exist an open interval  $J \subseteq I$  and a positive real number  $\rho$  such that, for each  $\lambda \in J$ , the equation

$$\Phi'(u) - \lambda\Psi'(u) = 0$$

has at least three solutions in  $X$  whose norms are less than  $\rho$ .

*Proof* We may apply Theorem 1.15 to the function  $f : X \times I \rightarrow \mathbb{R}$  defined by  $f(u, \lambda) = \Phi(u) - \lambda\Psi(u)$  for each  $(u, \lambda) \in X \times I$ . In particular, the fact that  $\Psi'$  is compact implies that  $\Psi$  is sequentially weakly continuous. Moreover, our assumptions ensure that, for each  $\lambda \in I$ , the function  $f(\cdot, \lambda)$  satisfies the (PS)-condition.  $\square$

In the sequel we point out a useful result concerning the *location* of parameters in Corollary 1.4. Actually, this location result is an improved version of Theorem 1.15 given by Bonanno [42].

**Theorem 1.16** *Let  $X$  be a separable and reflexive real Banach space, and let  $\Phi, \Psi : X \rightarrow \mathbb{R}$  be two continuously Gâteaux differentiable functionals. Assume that there exists  $u_0 \in X$  such that  $\Phi(u_0) = \Psi(u_0) = 0$*

and  $\Phi(u) \geq 0$  for every  $u \in X$  and that there exists  $u_1 \in X$ ,  $r > 0$  such that

- (i)  $r < \Phi(u_1)$ ;
- (ii)  $\sup_{\Phi(u) < r} \Psi(u) < r \frac{\Psi(u_1)}{\Phi(u_1)}$ .

Further, put

$$\bar{a} = \frac{\zeta r}{r \frac{\Psi(u_1)}{\Phi(u_1)} - \sup_{\Phi(u) < r} \Psi(u)},$$

with  $\zeta > 1$ , and assume that the functional  $\Phi - \lambda\Psi$  is sequentially weakly lower semi-continuous, satisfies the (PS)-condition and

- (iii)  $\lim_{\|u\| \rightarrow +\infty} (\Phi(u) - \lambda\Psi(u)) = +\infty$  for every  $\lambda \in [0, \bar{a}]$ .

Then, there exists an open interval  $\Lambda \subset [0, \bar{a}]$  and a positive real number  $\rho$  such that, for each  $\lambda \in \Lambda$ , the equation

$$\Phi'(u) - \lambda\Psi'(u) = 0$$

admits at least three solutions in  $X$  whose norms are less than  $\rho$ .

*Proof* We prove the minimax inequality from Corollary 1.4 with  $I = [0, \bar{a}]$  and the function  $h$  suitable chosen. Fix  $\zeta > 1$ . Due to (ii), there exists  $\sigma \in \mathbb{R}$  such that

$$\sup_{\Phi(u) < r} \Psi(u) + \frac{r \frac{\Psi(u_1)}{\Phi(u_1)} - \sup_{\Phi(u) < r} \Psi(u)}{\zeta} < \sigma < r \frac{\Psi(u_1)}{\Phi(u_1)}. \quad (1.16)$$

Due to (i) and (1.16) we obtain that  $0 < \sigma < \Psi(u_1)$ ; thus, introducing the notation  $A := \sigma \frac{\Phi(u_1)}{\Psi(u_1)}$ , we clearly have

$$\frac{A - \Phi(u_1)}{\sigma - \Psi(u_1)} = \frac{A}{\sigma}. \quad (1.17)$$

Let  $\lambda \in \mathbb{R}$ . Due to (1.17), one has either  $\lambda \geq \frac{A - \Phi(u_1)}{\sigma - \Psi(u_1)}$  or  $\lambda < \frac{A}{\sigma}$ . When  $\lambda \geq \frac{A - \Phi(u_1)}{\sigma - \Psi(u_1)}$  we have that

$$\inf_{u \in X} (\Phi(u) + \lambda(\sigma - \Psi(u))) \leq \Phi(u_1) + \lambda(\sigma - \Psi(u_1)) \leq A.$$

When  $\lambda < \frac{A}{\sigma}$ , we have

$$\inf_{u \in X} (\Phi(u) + \lambda(\sigma - \Psi(u))) \leq \Phi(u_0) + \lambda(\sigma - \Psi(u_0)) = \lambda\sigma < A.$$

Consequently,

$$\sup_{\lambda \in [0, \bar{a}]} \inf_{u \in X} (\Phi(u) + \lambda(\sigma - \Psi(u))) \leq \sup_{\lambda \in \mathbb{R}} \inf_{u \in X} (\Phi(u) + \lambda(\sigma - \Psi(u))) \leq A. \quad (1.18)$$

Set  $k = [r\Psi(u_1)/\Phi(u_1) - \sup_{\Phi(u) < r} \Psi(u)]/\zeta > 0$ . On account of (1.16), one has  $\sup_{\Phi(u) < r} \Psi(u) < \sigma - k$ . In particular, we have

$$\inf_{\Psi(u) > \sigma - k} \Phi(u) \geq r. \quad (1.19)$$

By (1.16) again, we have  $r > \sigma \frac{\Phi(u_1)}{\Psi(u_1)} = A$ . Consequently, from (1.19) one has

$$\inf_{\Psi(u) > \sigma - k} \Phi(u) > A.$$

Moreover, we also have

$$\bar{a}k = r > A.$$

After combining these facts, we obtain that

$$\begin{aligned} \inf_{u \in X} \sup_{\lambda \in [0, \bar{a}]} (\Phi(u) + \lambda(\sigma - \Psi(u))) &= \inf_{u \in X} (\Phi(u) + \bar{a} \max\{0, \sigma - \Psi(u)\}) \\ &\geq \min \left\{ \inf_{\Psi(u) > \sigma - k} \Phi(u); \inf_{\Psi(u) \leq \sigma - k} \Phi(u) + \bar{a}k \right\} > A. \end{aligned}$$

Combining this estimate with (1.18), the minimax inequality from Corollary 1.4 holds with the choices  $I = [0, \bar{a}]$  and  $h(\lambda) = \lambda\sigma$ . The proof is complete.  $\square$

We conclude the present subsection with a double eigenvalue problem. More precisely, we are interested in the minimal number of critical points of the functional  $\Phi - \lambda\Psi$  (from Theorem 1.16) which is perturbed by an arbitrarily functional of class  $C^1$ . To state this result, we are going to apply a pure topological result due to Ricceri [266].

**Theorem 1.17** *Let  $X$  be a reflexive real Banach space,  $I \subseteq \mathbb{R}$  an interval, and  $\varphi : X \times I \rightarrow \mathbb{R}$  a function such that  $\varphi(u, \cdot)$  is concave in  $I$  for all  $u \in X$ , while  $\varphi(\cdot, \lambda)$  is continuous, coercive and sequentially weakly lower semi-continuous in  $X$  for all  $\lambda \in I$ . Further, assume that*

$$\sup_I \inf_X \varphi < \inf_X \sup_I \varphi.$$

*Then, for each  $\eta > \sup_I \inf_X \varphi$ , there exists a non-empty open set  $A \subseteq I$  with the following property: for every  $\lambda \in A$  and every sequentially weakly*

lower semi-continuous functional  $\Upsilon : X \rightarrow \mathbb{R}$ , there exists  $\delta > 0$  such that for each  $\mu \in [0, \delta]$ , the functional  $\varphi(\cdot, \lambda) + \mu\Upsilon(\cdot)$  has at least two local minima lying in the set  $\{u \in X : \varphi(u, \lambda) < \eta\}$ .

Now we prove another variant of Corollary 1.4 which appears in Ricceri [267].

**Theorem 1.18** *Let  $X$  be a separable and reflexive real Banach space,  $I \subseteq \mathbb{R}$  an interval,  $\Phi : X \rightarrow \mathbb{R}$  a sequentially weakly lower semi-continuous functional of class  $C^1$  whose derivative admits a continuous inverse on  $X^*$  and  $\Psi : X \rightarrow \mathbb{R}$  a functional of class  $C^1$  with compact derivative. Assume that*

$$\lim_{\|u\| \rightarrow +\infty} (\Phi(u) - \lambda\Psi(u)) = +\infty$$

for all  $\lambda \in I$ , and that there exists  $\sigma \in \mathbb{R}$  such that

$$\sup_{\lambda \in I} \inf_{u \in X} (\Phi(u) + \lambda(\sigma - \Psi(u))) < \inf_{u \in X} \sup_{\lambda \in I} (\Phi(u) + \lambda(\sigma - \Psi(u))). \quad (1.20)$$

Then, there exist a non-empty open set  $A \subseteq I$  and a positive real number  $\rho$  with the following property: for every  $\lambda \in A$  and every functional  $\Upsilon : X \rightarrow \mathbb{R}$  of class  $C^1$  with compact derivative, there exists  $\delta > 0$  such that, for each  $\mu \in [0, \delta]$ , the equation

$$\Phi'(u) - \lambda\Psi'(u) + \mu\Upsilon'(u) = 0$$

has at least three solutions in  $X$  whose norms are less than  $\rho$ .

*Proof* Note that any functional on  $X$  of class  $C^1$  with compact derivative is sequentially weakly continuous; in particular, it is bounded on each bounded subset of  $X$ , due to the reflexivity of  $X$ . Consequently, the function  $\varphi : X \times I \rightarrow \mathbb{R}$  defined by

$$\varphi(u, \lambda) = \Phi(u) + \lambda(\sigma - \Psi(u))$$

satisfies all the hypotheses of Theorem 1.17. Fix  $\eta > \sup_I \inf_X \varphi$  and consider a non-empty open set  $A$  with the property stated in Theorem 1.17. Fix also a compact interval  $[a, b] \subset A$ . We have

$$\bigcup_{\lambda \in [a, b]} \{u \in X : \varphi(u, \lambda) < \eta\}$$

$$\subseteq \{u \in X : \Phi(u) - a\Psi(u) < \eta - a\sigma\} \cup \{u \in X : \Phi(u) - b\Psi(u) < \eta - b\sigma\}.$$

Note that the set on the right-hand side is bounded, due to the coercivity assumption. Consequently, there is some  $\tilde{\eta} > 0$  such that

$$\bigcup_{\lambda \in [a, b]} \{u \in X : \varphi(u, \lambda) < \eta\} \subseteq B_{\tilde{\eta}}, \quad (1.21)$$

where  $B_{\tilde{\eta}} = \{u \in X : \|u\| < \tilde{\eta}\}$ . Now, set

$$c^* = \sup_{B_{\tilde{\eta}}} \Phi + \max\{|a|, |b|\} \sup_{B_{\tilde{\eta}}} |\Psi|$$

and fix  $\rho > \tilde{\eta}$  such that

$$\bigcup_{\lambda \in [a, b]} \{u \in X : \Phi(u) - \lambda\Psi(u) \leq c^* + 2\} \subseteq B_{\rho}. \quad (1.22)$$

Now, let  $\Upsilon : X \rightarrow \mathbb{R}$  be any functional of class  $C^1$  with compact derivative. Let us choose a bounded function  $g : \mathbb{R} \rightarrow \mathbb{R}$  of class  $C^1$  such that  $g(t) = t$  for all  $t \in [-\sup_{B_{\rho}} |\Upsilon|, \sup_{B_{\rho}} |\Upsilon|]$ . Put  $\tilde{\Psi}(u) = g(\Psi(u))$  for all  $u \in X$ . So,  $\tilde{\Psi}$  is a bounded functional on  $X$  of class  $C^1$  such that

$$\tilde{\Upsilon}(u) = \Upsilon(u) \text{ for all } x \in B_{\rho}. \quad (1.23)$$

For any set  $Y \subset X$ , we have

$$\tilde{\Upsilon}'(Y) \subseteq g'(\Upsilon(Y))\Upsilon'(Y)$$

and hence it is clear that the derivative of  $\tilde{\Upsilon}$  is compact. Now, fix  $\lambda \in [a, b]$ . Then, taking (1.21) into account, there exists  $\tilde{\delta} > 0$  such that, for each  $\mu \in [0, \tilde{\delta}]$ , the functional  $\Phi - \lambda\Psi + \mu\tilde{\Upsilon}$  has two local minima, say  $e_0, e_1$  belonging to  $B_{\tilde{\eta}}$ . Further, set

$$\delta = \min \left\{ \tilde{\delta}, \frac{1}{1 + \sup_{\mathbb{R}} |g|} \right\}$$

and fix  $\mu \in [0, \delta]$ . Note that the functional  $\Phi - \lambda\Psi + \mu\tilde{\Upsilon}$  is coercive; moreover, an easy argument shows that this functional satisfies also the  $(PS)$ -condition. Consequently, if we denote by  $\Gamma$  the set of all continuous paths  $\gamma : [0, 1] \rightarrow X$  such that  $\gamma(0) = e_0, \gamma(1) = e_1$ , and set

$$c_{\lambda, \mu} = \inf_{\gamma \in \Gamma} \max_{t \in [0, 1]} (\Phi(\gamma(t)) - \lambda\Psi(\gamma(t)) + \mu\tilde{\Upsilon}(\gamma(t))),$$

by the mountain pass theorem for zero altitude (see Theorem 1.8), there exists  $e \in X$  such that

$$\Phi'(e) - \lambda\Psi'(e) + \mu\tilde{\Upsilon}'(e) = 0 \quad \text{and} \quad \Phi(e) - \lambda\Psi(e) + \mu\tilde{\Upsilon}(e) = c_{\lambda, \mu}.$$

Now, choosing in particular  $\gamma(t) = e_0 + t(e_1 - e_0)$ , we observe that

$$\begin{aligned} c_{\lambda, \mu} &\leq \max_{t \in [0, 1]} (\Phi(e_0 + t(e_1 - e_0)) \\ &\quad - \lambda \Psi(e_0 + t(e_1 - e_0)) + \mu \tilde{\Upsilon}(e_0 + t(e_1 - e_0))) \\ &\leq \sup_{B_{\tilde{\eta}}} \Phi + \max\{|a|, |b|\} \sup_{B_{\tilde{\eta}}} |\Psi| + \delta \sup_{\mathbb{R}} |g| \leq c^* + 1. \end{aligned}$$

Consequently, we have

$$\Phi(e) - \lambda \Psi(e) \leq c^* + 2.$$

By (1.22), it then follows that  $e \in B_\rho$  and so, by (1.23), one has  $\tilde{\Upsilon}'(e_i) = \Upsilon'(e_i)$  for  $i = 0, 1$  and  $\tilde{\Upsilon}'(e) = \Upsilon'(e)$ . In conclusion,  $e_0, e_1$  and  $e$  are three distinct solutions of the equation  $\Phi'(u) - \lambda \Psi'(u) + \mu \Upsilon'(u) = 0$  lying in  $B_\rho$ . The proof is complete.  $\square$

This subsection we end with a minmax result of B. Ricceri [261], which is very useful in applications.

**Theorem 1.19** *Let  $X$  be a topological space,  $\Lambda$  a real interval, and  $f : X \times \Lambda \rightarrow \mathbb{R}$  a function satisfying the following conditions:*

- (i) *For every  $x \in X$ , the function  $f(x, \cdot)$  is quasi-concave and continuous.*
- (ii) *For every  $\lambda \in \Lambda$ , the function  $f(\cdot, \lambda)$  is lower semicontinuous and each of its local minima is a global minimum.*
- (iii) *There exist  $\rho_0 > \sup_{\Lambda} \inf_X f$  and  $\lambda_0 \in \Lambda$  such that  $\{x \in X : f(x, \lambda_0) \leq \rho_0\}$  is compact.*

Then,

$$\sup_{\Lambda} \inf_X f = \inf_X \sup_{\Lambda} f.$$

#### 1.4.2 A general variational principle

The general variational principle of Ricceri gives alternatives for the multiplicity of critical points of certain functions depending on a parameter. It can be stated in a following way, see Ricceri [260].

**Theorem 1.20** *Let  $X$  be a reflexive real Banach space,  $\Phi, \Psi : X \rightarrow \mathbb{R}$  be two sequentially weakly lower semi-continuous, continuously Gâteaux differentiable functionals. Assume that  $\Psi$  is strongly continuous and coercive. For each  $s > \inf_X \Psi$ , set*

$$\varphi(s) := \inf_{u \in \Psi^s} \frac{\Phi(u) - \inf_{cl_w \Psi^s} \Phi}{s - \Psi(u)}, \quad (1.24)$$

where  $\Psi_s := \{u \in X : \Psi(u) < s\}$  and  $cl_w \Psi_s$  is the closure of  $\Psi_s$  in the weak topology of  $X$ . Furthermore, set

$$\delta := \liminf_{s \rightarrow (\inf_X \Psi)^+} \varphi(s), \quad \gamma := \liminf_{s \rightarrow +\infty} \varphi(s). \quad (1.25)$$

Then, the following conclusions hold.

- (A) If  $\delta < +\infty$  then, for every  $\lambda > \delta$ , either
- (A1) there is a global minimum of  $\Psi$  which is a local minimum of  $\Phi + \lambda\Psi$ , or
  - (A2) there is a sequence  $\{u_k\}$  of pairwise distinct critical points of  $\Phi + \lambda\Psi$ , with  $\lim_{k \rightarrow \infty} \Psi(u_k) = \inf_X \Psi$ , weakly converging to a global minimum of  $\Psi$ .
- (B) If  $\gamma < +\infty$  then, for every  $\lambda > \gamma$ , either
- (B1)  $\Phi + \lambda\Psi$  possesses a global minimum, or
  - (B2) there is a sequence  $\{u_k\}$  of critical points of the functional  $\Phi + \lambda\Psi$  such that  $\lim_{k \rightarrow \infty} \Psi(u_k) = +\infty$ .

In order to prove Theorem 1.20, we recall without proof a topological result of Ricceri [260].

**Theorem 1.21** *Let  $X$  be a topological space, and let  $\Phi, \Psi : X \rightarrow \mathbb{R}$  be two sequentially lower semi-continuous functions. Denote by  $I$  the set of all  $s > \inf_X \Psi$  such that the set  $\Psi_s := \{u \in X : \Psi(u) < s\}$  is contained in some sequentially compact subset of  $X$ . Assume that  $I \neq \emptyset$ . For each  $s \in I$ , denote by  $\mathcal{F}_s$  the family of all sequentially compact subsets of  $X$  containing  $\Psi_s$ , and put*

$$\alpha(s) = \sup_{K \in \mathcal{F}_s} \inf_K \Phi.$$

Then, for each  $s \in I$  and each  $\lambda$  satisfying

$$\lambda > \inf_{u \in \Psi_s} \frac{\Phi(u) - \alpha(s)}{s - \Psi(u)}$$

the restriction of the function  $\Phi + \lambda\Psi$  to  $\Psi_s$  has a global minimum.

*Proof of Theorem 1.20.* Endowing the space  $X$  with the weak topology, it is easy to verify that  $cl_w \Psi_s$  is the smallest sequentially weakly compact subset of  $X$  containing the set  $\Psi_s$ . Due to this observation, we also have that the function  $\alpha$  and the interval  $I$  from Theorem 1.21 have the form  $\alpha(s) = \inf_{u \in cl_w \Psi_s} \Phi(u)$  and  $I = ]\inf_X \Psi, \infty[$ , respectively.

(A) We assume that  $\delta < +\infty$ ; let  $\lambda > \delta$ . We may choose a sequence  $\{s_k\} \subset I$  with the properties that  $\lim_{k \rightarrow \infty} s_k = \inf_X \Psi$  and

$$\lambda > \inf_{u \in \Psi_{s_k}} \frac{\Phi(u) - \alpha(s_k)}{s_k - \Psi(u)}$$

for all  $k \in \mathbb{N}$ . Thanks to Theorem 1.21, for each  $k \in \mathbb{N}$ , the restriction of  $\Phi + \lambda\Psi$  to  $\Psi_{s_k}$  has a global minimum in the weak topology; we denote it by  $u_k$ . Consequently,  $u_k \in \Psi_{s_k}$  is a local minimum of  $\Phi + \lambda\Psi$  in the strong topology too. Moreover, if  $\bar{s} = \max_{k \in \mathbb{N}} s_k$ , we clearly have that  $\{u_k\} \subset \Psi_{\bar{s}}$ . On the other hand, the set  $\Psi_{\bar{s}}$  is contained in a sequentially compact subset of  $X$ . Therefore, the sequence  $\{u_k\}$  admits a subsequence, denoted in the same way, which converges weakly to an element  $u \in X$ . We claim that  $u$  is a global minimum of  $\Psi$ . Indeed, since  $\Psi$  is sequentially weakly lower semi-continuous, we have that

$$\Psi(u) \leq \liminf_{k \rightarrow \infty} \Psi(u_k) \leq \lim_{k \rightarrow \infty} s_k = \inf_X \Psi.$$

Taking a subsequence if necessary, we have that

$$\lim_{k \rightarrow \infty} \Psi(u_k) = \Psi(u) = \inf_X \Psi.$$

Now, if  $u = u_k$  for some  $k \in \mathbb{N}$  then (A1) holds; otherwise, we have (A2).

(B) We assume that  $\gamma < +\infty$ ; let  $\lambda > \gamma$ . We fix a sequence  $\{s_k\} \subset I$  with the properties that  $\lim_{k \rightarrow \infty} s_k = +\infty$  and

$$\lambda > \inf_{u \in \Psi_{s_k}} \frac{\Phi(u) - \alpha(s_k)}{s_k - \Psi(u)}$$

for all  $k \in \mathbb{N}$ . Due to Theorem 1.21, for each  $k \in \mathbb{N}$ , there exists  $u_k \in \Psi_{s_k}$  such that

$$\Phi(u_k) + \lambda\Psi(u_k) = \min_{\Psi_{s_k}}(\Phi + \lambda\Psi). \quad (1.26)$$

We have basically two different cases.

When  $\lim_{k \rightarrow \infty} \Psi(u_k) = \infty$ , the proof is complete, since every set  $\Psi_{s_k}$  is open, thus  $u_k$  are different critical points of  $\Phi + \lambda\Psi$ , that is, (B2) holds.

Now, we suppose that  $\liminf_{k \rightarrow \infty} \Psi(u_k) < \infty$ . Let us fix the number  $r = \max\{\inf_X \Psi, \liminf_{k \rightarrow \infty} \Psi(u_k)\}$ . For  $k \in \mathbb{N}$  large enough, we clearly have that  $\{u_k\} \subset \Psi_{r+1}$ . In particular, since  $\Psi_{r+1}$  is contained in a sequentially compact subset of  $X$ , the sequence  $\{u_k\}$  admits a subsequence, denoted in the same way, converging weakly to an element  $u \in X$ . We claim that  $u$  is a global minimum of  $\Psi$ . To see this, we fix  $v \in X$ . Since  $\{s_k\}$  tends

to  $+\infty$ , there exists  $k_0 \in \mathbb{N}$  such that  $\Psi(v) < s_{k_0}$ , that is,  $v \in \Psi_{s_{k_0}}$ . Due to the sequentially weakly lower semi-continuity of  $\Phi + \lambda\Psi$  (note that  $\lambda \geq 0$ ), relation (1.26), we have that

$$\begin{aligned} \Phi(u) + \lambda\Psi(u) &\leq \liminf_{k \rightarrow \infty} (\Phi(u_k) + \lambda\Psi(u_k)) \leq \Phi(u_{k_0}) + \lambda\Psi(u_{k_0}) \\ &= \min_{\Psi_{s_{k_0}}} (\Phi + \lambda\Psi) \leq \Phi(v) + \lambda\Psi(v). \end{aligned}$$

Consequently,  $u \in X$  is a global minimum of  $\Phi + \lambda\Psi$ , that is, (B1) holds. This completes the proof of Theorem 1.20.

The general variational principle of Ricceri (Theorem 1.20) implies immediately the following assertion: *if  $\Phi$  and  $\Psi$  are two sequentially weakly lower semi-continuous functionals on a reflexive Banach space  $X$  and if  $\Psi$  is also continuous and coercive, then the functional  $\Psi + \lambda\Phi$  has at least one local minimum for each  $\lambda > 0$  small enough.*

A deep topological argument developed by Ricceri [264] shows a more precise conclusion. We present here this result without proof.

**Theorem 1.22** *Let  $X$  be a separable and reflexive real Banach space, and let  $\Phi, \Psi : X \rightarrow \mathbb{R}$  be two sequentially weakly lower semi-continuous and continuously Gâteaux differentiable functionals, with  $\Psi$  coercive. Assume that the functional  $\Psi + \lambda\Phi$  satisfies the (PS)-condition for every  $\lambda > 0$  small enough and that the set of all global minima of  $\Psi$  has at least  $k$  connected components in the weak topology, with  $k \geq 2$ .*

*Then, for every  $s > \inf_X \Psi$ , there exists  $\bar{\lambda} > 0$  such that for every  $\lambda \in ]0, \bar{\lambda}[$ , the functional  $\Psi + \lambda\Phi$  has at least  $k + 1$  critical points,  $k$  of which are lying in the set  $\Psi_s = \{u \in X : \Psi(u) < s\}$ .*

**Remark 1.4** The first  $k$  critical points of  $\Psi + \lambda\Phi$  are actually local minima of the same functional which belong to the level set  $\Psi_s$  while the  $(k + 1)^{\text{th}}$  critical point is of mountain pass type where the assumption of the (PS)-condition is exploited. The mountain pass type point is located by means of the classification result of Ghoussoub-Preiss [130].

## 1.5 $H^1$ versus $C^1$ local minimizers

The result contained in this section establishes a surprising property of local minimizers, due to H. Brezis and L. Nirenberg [52].

Let  $\Omega$  be a bounded open set in  $\mathbb{R}^N$  with smooth boundary. Assume that  $f : \bar{\Omega} \times \mathbb{R} \rightarrow \mathbb{R}$  is a Lipschitz continuous function in  $u$ , uniformly in

$x$  such that for all  $(x, u) \in \Omega \times \mathbb{R}$ ,

$$|f(x, u)| \leq C(1 + |u|^p), \quad (1.27)$$

where  $1 < p \leq (N + 2)/(N - 2)$ . We point out that the exponent  $p$  can attain the critical Sobolev exponent  $(N + 2)/(N - 2)$ . Set  $F(x, u) := \int_0^u f(x, s) ds$  and define on  $H_0^1(\Omega)$  the *energy* functional

$$E(u) := \frac{1}{2} \int_{\Omega} |\nabla u|^2 dx - \int_{\Omega} F(x, u) dx.$$

**Definition 1.4** We say that  $u_0 \in C_0^1(\overline{\Omega})$  is a local minimizer of  $E$  in the  $C^1$  topology if there exists  $r > 0$  such that

$$E(u_0) \leq E(u) \quad \text{for every } u \in C_0^1(\overline{\Omega}) \text{ with } \|u - u_0\|_{C^1} \leq r.$$

We say that  $u_0 \in H_0^1(\overline{\Omega})$  is a local minimizer of  $E$  in the  $H_0^1$  topology if there exists  $\varepsilon_0 > 0$  such that

$$E(u_0) \leq E(u) \quad \text{for every } u \in H_0^1(\overline{\Omega}) \text{ with } \|u - u_0\|_{H^1} \leq \varepsilon_0.$$

The main result of this section is the following.

**Theorem 1.23** *Any local minimizer of  $E$  in the  $C^1$  topology is also a local minimizer of  $E$  in the  $H_0^1$  topology.*

*Proof* We can assume without loss of generality that 0 is a local minimizer of  $E$  in the  $C^1$  topology.

*Subcritical case:*  $p < (N + 2)/(N - 2)$ . Arguing by contradiction, we deduce that for any  $\varepsilon > 0$ , there exists  $u_\varepsilon \in H_0^1(\Omega)$  such that

$$\|u_\varepsilon\|_{H^1} \leq \varepsilon \quad \text{and} \quad E(u_\varepsilon) < E(0). \quad (1.28)$$

This enables us to apply a standard lower semi-continuity argument to deduce that  $\min_{u \in \overline{B_\varepsilon}} E(u)$  is attained by some point  $u_\varepsilon$ , where

$$B_\varepsilon := \{u \in H_0^1(\Omega); \|u\|_{H^1} < \varepsilon\}.$$

We claim that

$$u_\varepsilon \rightarrow 0 \quad \text{in } C^1(\overline{\Omega}).$$

Thus, by (1.28), we contradict the hypothesis that 0 is a local minimizer of  $E$  in the  $C^1$  topology.

In order to prove our claim, let  $\mu_\varepsilon \leq 0$  be a Lagrange multiplier of the corresponding Euler equation for  $u_\varepsilon$ , that is,

$$\langle E'(u_\varepsilon), v \rangle_{H^{-1}, H_0^1} = \mu_\varepsilon \langle u_\varepsilon, v \rangle_{H_0^1} \quad \text{for all } v \in H_0^1(\Omega).$$

Therefore

$$\int_{\Omega} [\nabla u_{\varepsilon} \cdot \nabla v - f(x, u_{\varepsilon})v] dx = \mu_{\varepsilon} \int_{\Omega} \nabla u_{\varepsilon} \cdot \nabla v dx \quad \text{for all } v \in H_0^1(\Omega).$$

This can be rewritten as

$$-(1 - \mu_{\varepsilon})\Delta u_{\varepsilon} = f(x, u_{\varepsilon}) \quad \text{in } \Omega. \quad (1.29)$$

Since  $u_{\varepsilon} \rightarrow 0$  in  $H^1(\Omega)$  it follows that, in order to deduce our claim, it is enough to apply the Arzelà–Ascoli theorem, after observing that

$$\|u_{\varepsilon}\|_{C^{1,\alpha}} \leq C.$$

This uniform estimate follows by a standard bootstrap argument. Indeed, assuming that  $u_{\varepsilon} \in L^{q_1}(\Omega)$  for some  $q_1 \geq 2N/(N-2)$ , then  $f(x, u_{\varepsilon}) \in L^r(\Omega)$ , where  $r = q_1/p$ . Thus, by Schauder elliptic estimates,  $u_{\varepsilon} \in W^{2,r}(\Omega) \subset L^{q_2}(\Omega)$ , with

$$\frac{1}{q_2} = \frac{p}{q_1} - \frac{2}{N},$$

provided that the right hand-side is positive. In the remaining case we have  $u_{\varepsilon} \in L^q(\Omega)$  for any  $q > 1$ . Since  $p < (N+2)/(N-2)$ , we deduce that  $q_2 > q_1$ . Starting with  $q_1 = 2N/(N-2)$  and iterating this process, we obtain an increasing divergent sequence  $(q_m)_{m \geq 1}$ . This shows that  $u_{\varepsilon} \in L^q(\Omega)$  for any  $q > 1$ . Returning now to (1.29) we deduce that  $u_{\varepsilon} \in W^{2,r}(\Omega)$  for any  $r > 1$ . Thus, by Sobolev embeddings,  $u_{\varepsilon} \in C^{1,\alpha}(\overline{\Omega})$  for any  $\alpha \in (0, 1)$ . This concludes the proof in the subcritical case corresponding to  $p < (N+2)/(N-2)$ .

*Critical case:*  $p = (N+2)/(N-2)$ . We argue again by contradiction. Thus, for all  $\varepsilon > 0$ , there exists  $u_{\varepsilon} \in H_0^1(\Omega)$  such that relation (1.28) holds. For any positive integer  $m$  we define the truncation map

$$T_m(t) = \begin{cases} -m & \text{if } t \leq -m \\ t & \text{if } -m \leq t \leq m \\ m & \text{if } t \geq m. \end{cases}$$

For any integer  $m \geq 1$ , define the energy functional

$$E_m(u) := \frac{1}{2} \int_{\Omega} |\nabla u|^2 dx - \int_{\Omega} F_m(x, u) dx,$$

where  $f_m(x, t) := f(x, T_m(t))$  and  $F_m(x, u) := \int_0^u f_m(x, t) dt$ .

We first observe that for all  $u \in H_0^1(\Omega)$ ,  $E_m(u) \rightarrow E(u)$  as  $m \rightarrow \infty$ . Thus,

for each  $\varepsilon > 0$ , there exists an integer  $m(\varepsilon)$  such that  $E_{m(\varepsilon)}(u_\varepsilon) < E(0)$ . Let  $v_\varepsilon \in H_0^1(\Omega)$  be such that

$$E_{m(\varepsilon)}(v_\varepsilon) = \min_{u \in \overline{B}_\varepsilon} E_\varepsilon(u).$$

Therefore

$$E_{m(\varepsilon)}(v_\varepsilon) \leq E_{m(\varepsilon)}(u_\varepsilon) < E(0).$$

We claim that  $v_\varepsilon \in C_0^1(\overline{\Omega})$  and that  $v_\varepsilon \rightarrow 0$  in  $C^1(\overline{\Omega})$ . The, if  $\varepsilon > 0$  is sufficiently small, we have

$$E(v_\varepsilon) = E_{m(\varepsilon)}(v_\varepsilon) < E(0),$$

which contradicts our assumption that 0 is a local minimizer of  $E$  in the  $C^1$  topology.

Returning to the proof of our claim, we observe that  $v_\varepsilon$  satisfies the Euler equation

$$-(1 - \mu_\varepsilon)\Delta v_\varepsilon = f_m(x, v_\varepsilon), \quad (1.30)$$

where

$$|f_m(x, u)| \leq C(1 + |u|^p) \quad (1.31)$$

with  $p = (N + 2)/(N - 2)$  and  $C$  not depending on  $m$ . Since  $v_\varepsilon \rightarrow 0$  in  $H_0^1(\Omega) \subset L^{2N/(N-2)}(\Omega)$ , Theorem IV.9 in [47] implies that there exists  $h \in L^{2N/(N-2)}(\Omega)$  such that, up to a subsequence still denoted by  $(v_\varepsilon)$ , we have  $|v_\varepsilon| \leq h$  in  $\Omega$ . Thus, by relation (1.31),

$$|f_m(x, v_\varepsilon)| \leq C(1 + a|v_\varepsilon|)$$

with  $a = h^{4/(N-2)} \in L^{N/2}(\Omega)$ . Using a bootstrap argument as in the proof of the subcritical case  $p < (N + 2)/(N - 2)$ , we deduce that  $(v_\varepsilon)$  is bounded in  $L^q(\Omega)$ , for any  $1 < q < \infty$ . Returning now to relation (1.30) and using (1.31) we obtain that  $(v_\varepsilon)$  is bounded in  $C^{1,\alpha}(\overline{\Omega})$  for any  $\alpha \in (0, 1)$ . Now, since  $v_\varepsilon \rightarrow 0$  in  $H_0^1(\Omega)$ , the Arzelà–Ascoli theorem implies that  $v_\varepsilon \rightarrow 0$  in  $C^{1,\alpha}(\overline{\Omega})$ . This concludes our proof.  $\square$

### 1.5.1 Application to sub and supersolutions

Consider the nonlinear elliptic problem

$$\begin{cases} -\Delta u = f(x, u) & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (1.32)$$

where  $f : \bar{\Omega} \times \mathbb{R} \rightarrow \mathbb{R}$  is a Lipschitz continuous function in  $u$ , uniformly in  $x$ .

The method of sub and supersolutions is a very useful tool in nonlinear analysis which asserts that if  $\underline{u}, \bar{u} \in C^2(\Omega) \cap C^1(\bar{\Omega})$  satisfy  $\underline{u} \leq \bar{u}$  and

$$\begin{cases} -\Delta \underline{u} - f(x, \underline{u}) \leq 0 \leq -\Delta \bar{u} - f(x, \bar{u}) & \text{in } \Omega \\ \underline{u} \leq 0 \leq \bar{u} & \text{on } \partial\Omega, \end{cases} \quad (1.33)$$

then problem (1.32) has at least a solution. The standard proof of this result relies on the maximum principle. By means of Theorem 1.23 we are able to give a *variational* proof to the following qualitative result which guarantees the existence of a local minimizer between a subsolution and a supersolution.

**Theorem 1.24** *Assume that functions  $\underline{u}, \bar{u} \in C^2(\Omega) \cap C^1(\bar{\Omega})$  with  $\underline{u} \leq \bar{u}$  satisfy (1.33). In addition, we suppose that neither  $\underline{u}$  nor  $\bar{u}$  is a solution of (1.32). Then there exists a solution  $u_0$  of problem (1.32) such that  $\underline{u} < u_0 < \bar{u}$  in  $\Omega$  and  $u_0$  is a local minimum of the associated energy functional  $E$  in  $H_0^1(\Omega)$ , where*

$$E(u) := \frac{1}{2} \int_{\Omega} |\nabla u|^2 dx - \int_{\Omega} F(x, u) dx.$$

*Proof* Define the truncated continuous function

$$f_0(x, t) := \begin{cases} f(x, \underline{u}(x)) & \text{if } t < \underline{u}(x) \\ f(x, t) & \text{if } \underline{u}(x) \leq t \leq \bar{u}(x) \\ f(x, \bar{u}(x)) & \text{if } t > \bar{u}(x). \end{cases}$$

Set  $F_0(x, u) := \int_0^u f_0(x, t) dt$  and

$$E_0(u) := \frac{1}{2} \int_{\Omega} |\nabla u|^2 dx - \int_{\Omega} F_0(x, u) dx.$$

Then  $f_0$  is bounded on  $\bar{\Omega} \times \mathbb{R}$ , hence  $E_0$  is bounded from below on  $H_0^1(\Omega)$ . Thus, by semi-continuity arguments,  $\inf_{u \in H_0^1(\Omega)} E_0(u)$  is achieved at some point  $u_0$  and

$$-\Delta u_0 = f_0(x, u_0) \quad \text{in } \Omega.$$

Bootstrap techniques (as developed in the proof of Theorem 1.23) imply that  $u_0 \in W^{2,p}(\Omega)$  for any  $p < \infty$ . So, by Sobolev embeddings,  $u_0 \in C^{1,\alpha}(\bar{\Omega})$  for all  $\alpha \in (0, 1)$ .

Next, we prove that  $\underline{u} < u_0 < \bar{u}$  in  $\Omega$ . Indeed, multiplying by  $(\underline{u} - u_0)^+$  the inequality

$$-\Delta(\underline{u} - u_0) \leq f(x, \underline{u}) - f(x, u_0)$$

and integrating on  $\Omega$ , we find  $(\underline{u} - u_0)^+ = 0$ . Thus,  $\underline{u} \leq u - 0$  in  $\Omega$ . In fact, the strong maximum principle implies that this inequality is strict. With a similar argument we deduce that  $u_0 < \bar{u}$  in  $\Omega$ . Using now  $\underline{u} < u_0 < \bar{u}$  and applying again the strong maximum principle, we find  $\varepsilon_0$  such that for all  $x \in \Omega$ ,

$$\underline{u}(x) + \varepsilon \operatorname{dist}(x, \partial\Omega) \leq u_0(x) \leq \bar{u}(x) - \varepsilon \operatorname{dist}(x, \partial\Omega).$$

We observe that the mapping  $[\underline{u}(x), \bar{u}(x)] \ni x \mapsto F_0(x, u) - F(x, u)$  is a function depending only on  $x$ . This shows that  $E(u) - E_0(u)$  is constant, provided that  $\|u - u_0\|_{C^1}$  is small enough. Since  $u_0$  is a global minimum of  $E_0$ , we deduce that  $u_0$  is a local minimum of  $E$  on  $C_0^1(\Omega)$ . Using now Theorem 1.23 we conclude that  $u_0$  is also a local minimum of  $E$  in  $H_0^1(\Omega)$ .  $\square$

## 1.6 Szulkin type functionals

Let  $X$  be a real Banach space and  $X^*$  its dual and we denote by  $\langle \cdot, \cdot \rangle$  the duality pair between  $X$  and  $X^*$ . Let  $E : X \rightarrow \mathbb{R}$  be a functional of class  $C^1$  and let  $\zeta : X \rightarrow \mathbb{R} \cup \{+\infty\}$  be a proper (that is,  $\neq +\infty$ ), convex, lower semi-continuous function. Then,  $I = E + \zeta$  is a *Szulkin-type functional*, see Szulkin [286]. An element  $u \in X$  is called a *critical point* of  $I = E + \zeta$  if

$$\langle E'(u), v - u \rangle + \zeta(v) - \zeta(u) \geq 0 \quad \text{for all } v \in X. \quad (1.34)$$

The number  $I(u)$  is a *critical value* of  $I$ .

For  $u \in D(\zeta) = \{u \in X : \zeta(u) < \infty\}$  we consider the set

$$\partial\zeta(u) = \{x^* \in X^* : \zeta(v) - \zeta(u) \geq \langle x^*, v - u \rangle, \quad \forall v \in X\}.$$

The set  $\partial\zeta(u)$  is called the *subdifferential* of  $\zeta$  at  $u$ . Note that an equivalent formulation for (1.34) is

$$0 \in E'(u) + \partial\zeta(u) \quad \text{in } X^*. \quad (1.35)$$

**Proposition 1.3** *Every local minimum point of  $I = E + \zeta$  is a critical point of  $I$  in the sense of (1.34).*

*Proof* Let  $u \in D(\zeta)$  be a local minimum point of  $I = E + \zeta$ . Due to the convexity of  $\zeta$ , for every  $t > 0$  small we have

$$0 \leq I((1-t)u + tv) - I(u) \leq E(u + t(v-u)) - E(u) + t(\zeta(v) - \zeta(u)).$$

Dividing by  $t > 0$  and letting  $t \rightarrow 0^+$ , we obtain (1.34).  $\square$

**Definition 1.5** The functional  $I = E + \zeta$  satisfies the Palais-Smale condition at level  $c \in \mathbb{R}$ , (shortly,  $(PSZ)_c$ -condition) if every sequence  $\{u_n\} \subset X$  such that  $\lim_{n \rightarrow \infty} I(u_n) = c$  and

$$\langle E'(u_n), v - u_n \rangle_X + \zeta(v) - \zeta(u_n) \geq -\varepsilon_n \|v - u_n\| \quad \text{for all } v \in X,$$

where  $\varepsilon_n \rightarrow 0$ , possesses a convergent subsequence.

**Remark 1.5** When  $\zeta = 0$ ,  $(PSZ)_c$ -condition is equivalent to the standard  $(PS)_c$ -condition.

### 1.6.1 Minimax results of Szulkin type

**Theorem 1.25** Let  $X$  be a Banach space,  $I = E + \zeta : X \rightarrow \mathbb{R} \cup \{+\infty\}$  a Szulkin-type functional and we assume that

- (i)  $I(u) \geq \alpha$  for all  $\|u\| = \rho$  with  $\alpha, \rho > 0$ , and  $I(0) = 0$ ;
- (ii) there is  $e \in X$  with  $\|e\| > \rho$  and  $I(e) \leq 0$ .

If  $I$  satisfies the  $(PSZ)_c$ -condition for

$$c = \inf_{\gamma \in \Gamma} \max_{t \in [0,1]} I(\gamma(t)),$$

$$\Gamma = \{\gamma \in C([0,1], X) : \gamma(0) = 0, \gamma(1) = e\},$$

then  $c$  is a critical value of  $I$  and  $c \geq \alpha$ .

**Theorem 1.26** Let  $X$  be a separable and reflexive Banach space, let  $I_1 = E_1 + \zeta_1$  Szulkin-type functionals and  $I_2 = E_2 : X \rightarrow \mathbb{R}$  a function of class  $C^1$  and let  $\Lambda \subseteq \mathbb{R}$  be an interval. Suppose that

- (i).  $E_1$  is weakly sequentially lower semi-continuous and  $E_2$  is weakly sequentially continuous;
- (ii). for every  $\lambda \in \Lambda$  the function  $I_1 + \lambda I_2$  fulfils  $(PSZ)_c$ ,  $c \in \mathbb{R}$ , with

$$\lim_{\|u\| \rightarrow +\infty} (I_1(u) + \lambda I_2(u)) = +\infty;$$

- (iii). there exists a continuous concave function  $h : \Lambda \rightarrow \mathbb{R}$  satisfying

$$\sup_{\lambda \in \Lambda} \inf_{u \in X} (I_1(u) + \lambda I_2(u) + h(\lambda)) < \inf_{u \in X} \sup_{\lambda \in \Lambda} (I_1(u) + \lambda I_2(u) + h(\lambda)).$$

Then there is an open interval  $\Lambda_0 \subseteq \Lambda$ , such that for each  $\lambda \in \Lambda_0$  the function  $I_1 + \lambda I_2$  has at least three critical points in  $X$ .

In the next, we give a variant of Ricceri variational principle for Szulkin type functionals which will be used in the study of the existence of scalar systems.

Suppose now that  $X$  and  $Y$  are real Banach spaces such that  $X$  is compactly embedded in  $Y$ . Let  $E_1: Y \rightarrow \mathbb{R}$  and  $E_2: X \rightarrow \mathbb{R}$  be  $C^1$  functions, and let  $\zeta_1: X \rightarrow ]-\infty, +\infty]$  be convex, proper, and lower semi-continuous. Define the maps  $I_1: X \rightarrow ]-\infty, +\infty]$  and  $I_2: X \rightarrow \mathbb{R}$  as

$$I_1(u) = E_1(u) + \zeta_1(u), \quad I_2(u) = E_2(u), \quad \text{for all } x \in X.$$

Denote by  $D(\zeta_1) := \{u \in X \mid \zeta_1(u) < +\infty\}$  and assume that

$$I_2^{-1}(]-\infty, \rho]) \cap D(\zeta_1) \neq \emptyset, \quad \text{for all } \rho > \inf_X I_2. \quad (1.36)$$

For every  $\rho > \inf_X I_2$  put

$$\varphi(\rho) := \inf_{u \in \Psi^{-1}(]-\infty, \rho])} \frac{\Phi(u) - \inf_{v \in (I_2^{-1}(]-\infty, \rho]))_w} \Phi(v)}{\rho - I_2(u)}, \quad (1.37)$$

where  $(I_2^{-1}(]-\infty, \rho]))_w$  is the weak closure of  $I_2^{-1}(]-\infty, \rho])$ ,

$$\gamma := \liminf_{\rho \rightarrow +\infty} \varphi(\rho), \quad (1.38)$$

$$\delta := \liminf_{\rho \rightarrow (\inf_X I_2)^+} \varphi(\rho). \quad (1.39)$$

**Theorem 1.27** *Suppose that  $X$  is reflexive,  $I_2$  is weakly sequentially lower semi-continuous and coercive, and (1.36) is fulfilled. Then the following assertions hold:*

- (a) *For every  $\rho > \inf_X I_2$  and every  $\lambda > \varphi(\rho)$  the function  $I_1 + \lambda I_2$  has a critical point (more exactly: a local minimum) lying in  $I_2^{-1}(]-\infty, \rho]) \cap D(\zeta_1)$ .*
- (b) *If  $\gamma < +\infty$  then, for each  $\lambda > \gamma$ , either*
  - (b1)  *$I_1 + \lambda I_2$  has a global minimum, or*
  - (b2) *there is a sequence  $\{u_n\}$  of critical points (more exactly: local minima) of  $I_1 + \lambda I_2$  lying in  $D(\zeta_1)$  and such that  $\lim_{n \rightarrow \infty} I_2(u_n) = +\infty$ .*
- (c) *If  $\delta < +\infty$  then, for every  $\lambda > \delta$ , either*
  - (c1)  *$I_1 + \lambda I_2$  has a local minimum which is also a global minimum of  $I_2$ , or*

- (c2) *there is a sequence  $\{u_n\}$  of pairwise distinct critical points (more exactly: local minima) of  $I_1 + \lambda I_2$  lying in  $D(\zeta_1)$  and such that  $\{u_n\}$  converges weakly to a global minimum of  $I_2$  and  $\lim_{n \rightarrow \infty} I_2(u_n) = \inf_X I_2$ .*

### 1.6.2 Principle of symmetric criticality

The principle of symmetric criticality plays a central role in many problems from the differential geometry, physics and in partial differential equations. First it was proved by Palais [235] for functionals of class  $C^1$ . In this subsection we recall this principle for functionals of class  $C^1$  defined on Banach spaces and we state its version for Szulkin type functionals, based on the paper of Kobayashi-Otani [160].

Let  $X$  be a Banach space and let  $X^*$  its dual. The norms of  $X$  and  $X^*$  will be denoted by  $\|\cdot\|$  and  $\|\cdot\|_*$ , respectively. We shall denote by  $\langle \cdot, \cdot \rangle$  the duality pairing between  $X$  and  $X^*$ .

Let  $G$  be a group,  $e$  its identity element, and let  $\pi$  a representation of  $G$  over  $X$ , that is  $\pi(g) \in L(X)$  for each  $g \in G$  (where  $L(X)$  denotes the set of the linear and bounded operator from  $X$  into  $X$ ), and

- a)  $\pi(e)u = u, \forall u \in X$ ;  
b)  $\pi(g_1 g_2)u = \pi(g_1)(\pi(g_2)u), \forall g_1, g_2 \in G$  and  $u \in X$ .

The representation  $\pi_*$  of  $G$  over  $X^*$  is naturally induced by  $\pi$  by the relation

$$\langle \pi_*(g)v^*, u \rangle = \langle v^*, \pi(g^{-1})u \rangle, \forall g \in G, v^* \text{ and } u \in X. \quad (1.40)$$

We often write  $gu$  or  $gv^*$  instead of  $\pi(g)u$  or  $\pi_*(g)v^*$ , respectively.

A function  $h : X \rightarrow \mathbb{R}$  (respectively,  $h : X^* \rightarrow \mathbb{R}$ ) is called  $G$ -invariant if  $h(gu) = h(u)$  (respectively,  $h(gv^*) = h(v^*)$ ) for every  $u \in X$  (respectively,  $v^* \in X^*$ ) and  $g \in G$ . A subset  $M$  of  $X$  is called  $G$ -invariant (respectively,  $M^*$  of  $X^*$ ) if

$$gM = \{gu : u \in M\} \subseteq M \quad (\text{respectively, } gM^* \subseteq M^*) \quad \forall g \in G.$$

The fixed point sets of the group action  $G$  on  $X$  and  $X^*$  (some authors call them  $G$ -symmetric points) are defined as

$$\Sigma = X^G = \{u \in X : gu = u \forall g \in G\},$$

$$\Sigma_* = (X^*)^G = \{v^* \in X^* : gv^* = v^* \forall g \in G\}.$$

Hence, by (1.40), we can see that  $v^* \in X^*$  is symmetric if and only if  $v^*$

is a  $G$ -invariant functional. The sets  $\Sigma$  and  $\Sigma_*$  are closed linear subspaces of  $X$  and  $X^*$ , respectively. So  $\Sigma$  and  $\Sigma_*$  are regarded as Banach spaces with their induced topologies. We introduce the notation

$$C_G^1(X) = \{f : X \rightarrow \mathbb{R} : f \text{ is } G\text{-invariant and of class } C^1\}.$$

We consider the following statement, called as *principle of symmetric criticality* (shortly, (PSC)):

- If  $f \in C_G^1(X)$  and  $(f|_\Sigma)'(u) = 0$ , then  $f'(u) = 0$ .

**Theorem 1.28** (PSC) *is valid if and only if  $\Sigma_* \cap \Sigma^\perp = \{0\}$ , where  $\Sigma^\perp = \{v^* \in X^* : \langle v^*, u \rangle = 0, \forall u \in \Sigma\}$ .*

*Proof* "  $\Rightarrow$  " Suppose that  $\Sigma_* \cap \Sigma^\perp = \{0\}$  and let  $u_0 \in \Sigma$  be a critical point of  $f|_\Sigma$ . We show that  $f'(u_0) = 0$ . Since  $f(u_0) = f|_\Sigma(u_0)$  and  $f(u_0 + v) = f|_\Sigma(u_0 + v)$  for all  $v \in \Sigma$ , we obtain  $\langle f'(u_0), v \rangle = \langle (f|_\Sigma)'(u_0), v \rangle_\Sigma$  for every  $v \in \Sigma$ , where  $\langle \cdot, \cdot \rangle_\Sigma$  denotes the duality pairing between  $\Sigma$  and its dual  $\Sigma^*$ . This implies that  $f'(u_0) \in \Sigma^\perp$ . On the other hand, from the  $G$ -invariance of  $f$  follows that

$$\begin{aligned} \langle f'(gu), v \rangle &= \lim_{t \rightarrow 0} \frac{f(gu + tv) - f(gu)}{t} = \lim_{t \rightarrow 0} \frac{f(u + tg^{-1}v) - f(u)}{t} \\ &= \langle f'(u), g^{-1}v \rangle = \langle gf'(u), v \rangle \end{aligned}$$

for all  $g \in G$  and  $u, v \in X$ . This means that  $f'$  is  $G$ -equivariant, that is,  $f'(gu) = gf'(u)$  for every  $g \in G$  and  $u \in X$ . Since  $u_0 \in \Sigma$ , we obtain  $gf'(u_0) = f'(u_0)$  for all  $g \in G$ , that is,  $f'(u_0) \in \Sigma_*$ . Thus we conclude  $f'(u_0) \in \Sigma_* \cap \Sigma^\perp = \{0\}$ . Therefore  $f'(u_0) = 0$ .

"  $\Leftarrow$  " Suppose that there exists a non-zero element  $v^* \in \Sigma_* \cap \Sigma^\perp$  and define  $f_*(\cdot)$  by  $f_*(u) = \langle v^*, u \rangle$ . It is clear that  $f_* \in C_G^1(X)$  and  $(f_*)'(\cdot) = v^* \neq 0$ , so  $f_*$  has no critical point in  $X$ . On the other hand  $v^* \in \Sigma^\perp$  implies  $v^*|_\Sigma = 0$ , thus  $(f_*|_\Sigma)'(u) = 0$  for every  $u \in \Sigma$  which contradicts (PSC).  $\square$

In the sequel, we are interested to find conditions in order to have  $\Sigma_* \cap \Sigma^\perp = \{0\}$ , that is, (PSC) to be verified. There are two ways to achieve this purpose, the so-called "compact" as well as the "isometric" cases. We are dealing now with the first one whose original form has been given by Palais [235].

**Theorem 1.29** *Let  $G$  be a compact topological group and the representation  $\pi$  of  $G$  over  $X$  is continuous, that is,  $(g, u) \rightarrow gu$  is a continuous function  $G \times X$  into  $X$ . Then (PSC) holds.*

In order to prove this theorem, we recall that for each  $u \in X$ , there exists a unique element  $Au \in X$  such that

$$\langle v^*, Au \rangle = \int_G \langle v^*, gu \rangle dg, \quad \forall v^* \in X^*, \quad (1.41)$$

where  $dg$  is the normalized Haar measure on  $G$ . The mapping  $A : X \rightarrow X$  is called the *averaging operator* on  $G$ .

*Proof of Theorem 1.29.* On account of Theorem 1.28, it is enough to verify the condition  $\Sigma_* \cap \Sigma^\perp = \{0\}$ . Let  $v^* \in \Sigma_* \cap \Sigma^\perp$  fixed and suppose that  $v^* \neq 0$ . Let us define the hyperplane  $H_{v^*} = \{u \in X : \langle v^*, u \rangle = 1\}$  which is a non-empty, closed, convex subset of  $X$ . For any  $u \in H_{v^*}$ , since  $v^* \in \Sigma_*$ , we have

$$\begin{aligned} \langle v^*, Au \rangle &= \int_G \langle v^*, gu \rangle dg = \int_G \langle g^{-1}v^*, u \rangle dg = \int_G \langle v^*, u \rangle dg = \\ &= \langle v^*, u \rangle \int_G dg = \langle v^*, u \rangle = 1. \end{aligned}$$

Note however that  $v^* \in \Sigma^\perp$ , thus  $\langle v^*, Au \rangle = 0$  for any  $u \in H_{v^*}$ , contradiction.

Another possibility for (PSC) to be valid is the following one which will be given without proof.

**Theorem 1.30** *Assume that  $X$  is a reflexive and strictly convex Banach space and  $G$  acts isometrically on  $X$ , that is,  $\|gu\| = \|u\|$  for all  $g \in G$  and  $u \in X$ . Then (PSC) holds.*

We conclude this subsection with a non-smooth version of (PSC). On account of relation (1.35), we are entitled to consider the following form of the principle of symmetric criticality for Szulkin functionals (shortly, (PSCSZ)):

- If  $I = E + \zeta : X \rightarrow \mathbb{R} \cup \{+\infty\}$  is a  $G$ -invariant Szulkin type functional and  $0 \in (E|_\Sigma)'(u) + \partial(\zeta|_\Sigma)(u)$  in  $\Sigma^*$  then  $0 \in E'(u) + \partial\zeta(u)$  in  $X^*$ .

Although a slightly more general version is proven by Kobayashi-Otani [160], we recall the following form of the *principle of symmetric criticality for Szulkin functionals* which will be applied in the next chapters.

**Theorem 1.31** [160, Theorem 3.16] *Let  $X$  be a reflexive Banach space and let  $I = E + \zeta : X \rightarrow \mathbb{R} \cup \{+\infty\}$  be a Szulkin-type functional on*

$X$ . If a compact group  $G$  acts linearly and continuously on  $X$ , and the functionals  $E$  and  $\zeta$  are  $G$ -invariant, then (PSCSZ) holds.

The proof of Theorem 1.31 is quite involved, it combines in an ingenious way various methods and notions from convex and functional analysis. The interested reader may consult the paper of Kobayashi-Otani [160].

## 1.7 Pohozaev's fibering method

The *fibering scheme* was introduced by S.I. Pohozaev in [247], [248]. Although there are various versions of it, we present the so-called *one-parametric fibering method*.

Let  $X$  be a real Banach space,  $f : X \rightarrow \mathbb{R}$  be a functional such that  $f$  is of class  $C^1$  on  $X \setminus \{0\}$ . By means of  $f$  we define the function  $\tilde{f} : (\mathbb{R} \setminus \{0\}) \times X \rightarrow \mathbb{R}$  by

$$\tilde{f}(\lambda, v) = f(\lambda v), \quad (\lambda, v) \in (\mathbb{R} \setminus \{0\}) \times X. \quad (1.42)$$

Let  $S = \{u \in X : \|u\| = 1\}$ .

**Definition 1.6** A point  $(\lambda, v) \in (\mathbb{R} \setminus \{0\}) \times S$  is conditionally critical of the function  $\tilde{f}$  if

$$-\tilde{f}'(\lambda, v) \in N_{(\mathbb{R} \setminus \{0\}) \times S}(\lambda, v),$$

where  $N_{(\mathbb{R} \setminus \{0\}) \times S}(\lambda, v)$  is the normal cone to the set  $(\mathbb{R} \setminus \{0\}) \times S$  at the point  $(\lambda, v)$ .

We have the following result, proved by Pohozaev [248].

**Theorem 1.32** *Let  $X$  be a real Banach space with differentiable norm on  $X \setminus \{0\}$  and let  $(\lambda, v) \in (\mathbb{R} \setminus \{0\}) \times S$  be a conditionally critical point of  $\tilde{f} : (\mathbb{R} \setminus \{0\}) \times X \rightarrow \mathbb{R}$ . Then  $u = \lambda v \in X \setminus \{0\}$  is a critical point of  $f : X \rightarrow \mathbb{R}$ , that is,  $f'(u) = 0$ .*

*Proof* Due to the hypothesis, we have that

$$-\tilde{f}'(\lambda, v) \in N_{(\mathbb{R} \setminus \{0\}) \times S}(\lambda, v).$$

On account of Proposition A.2, see Appendix A, we obtain

$$(-\tilde{f}'_{\lambda}(\lambda, v), -\tilde{f}'_v(\lambda, v)) \in N_{\mathbb{R} \setminus \{0\}}(\lambda) \times N_S(v).$$

Here,  $\tilde{f}'_v$  denotes the differential of  $\tilde{f}$  with respect to  $v$  in the whole space  $X$ , while  $\tilde{f}'_{\lambda}$  is the derivative of  $\tilde{f}$  with respect to the variable  $\lambda$ .

Note that  $N_{\mathbb{R} \setminus \{0\}}(\lambda) = \{0\}$ , thus

$$\tilde{f}'_{\lambda}(\lambda, v) = 0. \quad (1.43)$$

Moreover, since  $T_S(v)$  is a linear space, the condition  $-\tilde{f}'_v(\lambda, v) \in N_S(v)$  reduces to the fact that

$$\langle \tilde{f}'_v(\lambda, v), w \rangle = 0 \text{ for all } w \in T_S(v).$$

Since  $T_S(v) = \text{Ker} \|\cdot\|'(v)$ , see Zeidler [302, Theorem 43.C], there are  $\kappa, \mu \in \mathbb{R}$  such that  $\kappa^2 + \mu^2 \neq 0$  and

$$\kappa \tilde{f}'_v(\lambda, v) = \mu \|\cdot\|'(v). \quad (1.44)$$

From (1.44) it follows that for every  $v \in X$ ,

$$\kappa \langle \tilde{f}'_v(\lambda, v), v \rangle = \mu \langle \|\cdot\|'(v), v \rangle. \quad (1.45)$$

A simple calculation shows that

$$\tilde{f}'_v(\lambda, v) = \lambda f'(\lambda v). \quad (1.46)$$

Moreover, we have

$$\langle \tilde{f}'_v(\lambda, v), v \rangle = \lambda \langle f'(\lambda v), v \rangle = \lambda \tilde{f}'_{\lambda}(\lambda, v).$$

Consequently, from (1.45) and the above relation we obtain that

$$\lambda \kappa \tilde{f}'_{\lambda}(\lambda, v) = \mu \langle \|\cdot\|'(v), v \rangle = \mu.$$

From this equality and (1.43) we get  $\mu = 0$ . Then, we necessarily have  $\kappa \neq 0$ . Combining (1.44) with (1.46), we have that

$$\lambda f'(u) = \tilde{f}'_v(\lambda, v) = 0$$

for  $u = \lambda v$ . Since  $\lambda \neq 0$ , we obtain  $f'(u) = 0$  which concludes the proof.  $\square$

## 1.8 Historical comments

Ekeland's variational principle [102] was established in 1974 and is the nonlinear version of the Bishop–Phelps theorem [243, 244], with its main feature of how to use the norm completeness and a partial ordering to obtain a point where a linear functional achieves its supremum on a closed bounded convex set. A major consequence of Ekeland's variational principle is that even if it is not always possible to minimize a nonnegative  $C^1$

functional  $\Phi$  on a Banach space; however, there is always a minimizing sequence  $(u_n)_{n \geq 1}$  such that  $\Phi'(u_n) \rightarrow 0$  as  $n \rightarrow \infty$ .

Sullivan [285] observed that Ekeland's variational principle characterizes complete metric spaces in the following sense.

**Theorem 1.33** *Let  $(M, d)$  be a metric space. Then  $M$  is complete if and only if the following holds: For every application  $\Phi : M \rightarrow (-\infty, \infty]$ ,  $\Phi \not\equiv \infty$ , which is bounded from below, and for every  $\varepsilon > 0$ , there exists  $z_\varepsilon \in M$  such that*

- (i)  $\Phi(z_\varepsilon) \leq \inf_M \Phi + \varepsilon$ ;
- (ii)  $\Phi(x) > \Phi(z_\varepsilon) - \varepsilon d(x, z_\varepsilon)$ , for any  $x \in M \setminus \{z_\varepsilon\}$ .

Many equivalent results to Ekeland's variational principle were discovered independently. Some of the most important are the following: the maximal point version of Phelps' lemma [244, Lemma 1.2], the Krasnoselskii–Zabreiko theorem on normal solvability of operator equations [301], the Takahashi minimization principle [287], the *drop* theorem of Daneš related to the theory of normal solvability of nonlinear equations [81], and the *flower petal* theorem of Penot [240].

The mountain pass theorem was established by Ambrosetti and Rabinowitz in [8]. Their original proof relies on some deep deformation techniques developed by Palais and Smale [234], [237], who put the main ideas of the Morse theory into the framework of differential topology on infinite dimensional manifolds. In this way, Palais and Smale replaced the finite dimensionality assumption with an appropriate compactness hypothesis. H. Brézis and L. Nirenberg provided in [51] a simpler proof of the mountain pass theorem, by combining Ekeland's variational principle and the pseudogradient lemma.

The main underlying ideas of the fibering methods are the following.

1. Let  $X, Y$  be two real Banach spaces and  $A : X \rightarrow Y$  be a nonlinear operator. Fix an element  $h \in Y$  and consider the equation

$$A(u) = h. \tag{1.47}$$

We also consider the wider Banach space  $\tilde{X}, \tilde{Y}$  such that  $X \subset \tilde{X}$  and  $Y \subset \tilde{Y}$ . Now let  $\tilde{A} : \tilde{X} \rightarrow \tilde{Y}$  be a nonlinear operator such that  $\tilde{A}|_X = A$ . Instead of (1.47) we consider the extended equation

$$\tilde{A}(\tilde{u}) = \tilde{h}. \tag{1.48}$$

2. We equip  $\tilde{X}$  with a *nonlinear structure* associated to the nonlinear operator  $A$ .

For given spaces  $X, Y$  and a nonlinear operator  $A : X \rightarrow Y$ , that is, for the triple  $(X, A, Y)$  we construct a triple  $(\xi, \alpha, \eta)$ , where  $\xi : \tilde{X} \rightarrow X$  and  $\eta : \tilde{Y} \rightarrow Y$  are fibrations and  $\alpha = (\tilde{A}, A) : (\tilde{X}, X) \rightarrow (\tilde{Y}, Y)$  is a morphism of fibrations, that is,  $\eta \circ \tilde{A} = A \circ \xi$ . If we take  $\tilde{X} = \mathbb{R}^k \times X$ , then we obtain the  $k$ -parametric fibering methods. If  $k = 1$  we get the *one-parametric fibering methods*. The basic idea in *one parametric fibering methods* is the representation of solutions for  $A(u) = h$  in the form

$$u = tv, \quad (1.49)$$

where  $t \in \mathbb{R} \setminus \{0\}$  is a real parameter and  $v \in X \setminus \{0\}$  satisfying the fibering constraint

$$H(t, v) = c. \quad (1.50)$$

The function  $H$  will be said *fibering functional*. In particular case, when  $H(t, v) \equiv \|v\|$ , the condition (1.50) becomes  $\|v\| = 1$  and it is called *spherical fibering*.

## 2

# Variational Inequalities

Just as houses are made of  
stones, so is science made of  
facts; but a pile of stones is  
not a house and a collection of  
facts is not necessarily science.

---

Henri Poincaré (1854–1912)

### 2.1 Introduction

The theory of variational inequalities appeared in the middle 1960's, in connection with the notion of subdifferential in the sense of convex analysis, see Definition A.5. One of the first and simplest results in the theory of variational inequalities is the Hartman–Stampacchia theorem which asserts that if  $V$  is a finite dimensional Banach space,  $K \subset V$  is compact and convex,  $A : K \rightarrow V^*$  is continuous, then there exists  $u \in K$  such that for every  $v \in K$ ,

$$\langle Au, v - u \rangle \geq 0. \quad (2.1)$$

We postpone for the end of this chapter an elementary proof to this pioneering result in the theory of variational inequalities.

All the inequality problems treated to the beginning of 1980's were related to convex energy functionals and therefore strictly connected to monotonicity: for instance, only monotone (possibly multivalued) boundary conditions and stress-strain laws could be studied. Nonconvex inequality problems first appeared in [206] in the setting of global analysis and were related to the subdifferential introduced in [84] (see

Marino [205] for a survey of the developments in this direction). A typical feature of nonconvex problems is that, while in the convex case the stationary variational inequalities give rise to minimization problems for the potential or for the energy, in the nonconvex case the problem of the stationarity of the potential emerges and therefore it becomes reasonable to expect results also in the line of critical point theory.

## 2.2 Variational inequalities on unbounded strips

### 2.2.1 Multi-dimensional case

Let  $\Omega = \omega \times \mathbb{R}^l$  be an unbounded strip (or, in other words, a strip-like domain), where  $\omega \subset \mathbb{R}^m$  is open bounded, and  $l \geq 2$ ,  $m \geq 1$ . Let  $f : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$  be a continuous function, which satisfies the following condition:

(f1) there exist  $c_1 > 0$  and  $p \in (2, 2^*)$  such that

$$|f(x, s)| \leq c_1(|s| + |s|^{p-1}), \quad \text{for every } (x, s) \in \Omega \times \mathbb{R}.$$

We have denoted by  $2^* = 2(m+l)(m+l-2)^{-1}$  the critical Sobolev exponent.

Let  $H_0^1(\Omega)$  denote the Sobolev space endowed with the inner product

$$\langle u, v \rangle_{H_0^1} = \int_{\Omega} \nabla u \nabla v \, dx$$

and norm  $\|\cdot\|_{H_0^1} = \sqrt{\langle \cdot, \cdot \rangle_{H_0^1}}$ , while the norm of  $L^\alpha(\Omega)$  will be denoted by  $\|\cdot\|_\alpha$ . Then the embedding  $H_0^1(\Omega) \hookrightarrow L^\alpha(\Omega)$  is continuous, for all  $\alpha \in [2, 2^*]$ . Thus, there exists  $k_\alpha > 0$  such that  $\|u\|_\alpha \leq k_\alpha \|u\|_{H_0^1}$ , for every  $u \in H_0^1(\Omega)$ .

Consider finally the closed convex cone

$$\mathcal{K} = \{u \in H_0^1(\Omega) : u \geq 0 \text{ a.e. in } \Omega\}.$$

The aim of this section is to study the following eigenvalue problem for variational inequality (denoted by (P)):

Find  $(u, \lambda) \in \mathcal{K} \times (0, \infty)$  such that for all  $v \in \mathcal{K}$ ,

$$\int_{\Omega} \nabla u(x) (\nabla v(x) - \nabla u(x)) \, dx - \lambda \int_{\Omega} f(x, u(x)) (v(x) - u(x)) \, dx \geq 0.$$

We say that a function  $h : \Omega \rightarrow \mathbb{R}$  is *axially symmetric*, if  $h(x, y) = h(x, gy)$  for all  $x \in \omega$ ,  $y \in \mathbb{R}^l$  and  $g \in O(l)$ . In particular, we denote by  $H_{0,s}^1(\Omega)$  the closed subspace of axially symmetric functions of  $H_0^1(\Omega)$ . Define  $F : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$  by  $F(x, s) = \int_0^s f(x, t)dt$  and beside of **(f1)**, we make the following assumptions:

- (f2).**  $\lim_{s \rightarrow 0} \frac{|f(x, s)|}{s} = 0$  uniformly for every  $x \in \Omega$ .  
**(f3).** There exist  $q \in ]0, 2[$ ,  $\nu \in [2, 2^*]$ ,  $\alpha \in L^{\nu/(\nu-q)}(\Omega)$ ,  $\beta \in L^1(\Omega)$  such that

$$F(x, s) \leq \alpha(x)|s|^q + \beta(x).$$

- (f4).** There exists  $u_0 \in H_{0,s}^1(\Omega) \cap \mathcal{K}$  such that  $\int_{\Omega} F(x, u_0(x))dx > 0$ .

The main result of this section is the following.

**Theorem 2.1** *Let  $f : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$  be a continuous function which satisfies **(f1)**-**(f4)** and  $F(\cdot, s)$  is axially symmetric for every  $s \in \mathbb{R}$ . Then there is an open interval  $\Lambda_0 \subset (0, \infty)$  such that for every  $\lambda \in \Lambda_0$  there are at least three distinct elements  $u_i^\lambda \in \mathcal{K}$  ( $i \in \{1, 2, 3\}$ ) which are axially symmetric, having the property that  $(u_i^\lambda, \lambda)$  are solutions of (P) for every  $i \in \{1, 2, 3\}$ .*

From now on, we assume that the hypotheses of Theorem 2.1 are fulfilled. Before to prove Theorem 2.1, some preliminary results will be given.

**Remark 2.1** *For every  $\varepsilon > 0$  there exists  $c(\varepsilon) > 0$  such that*

- (i).  $|f(x, s)| \leq \varepsilon|s| + c(\varepsilon)|s|^{p-1}$  for every  $(x, s) \in \Omega \times \mathbb{R}$ .  
(ii).  $|F(x, s)| \leq \varepsilon s^2 + c(\varepsilon)|s|^p$  for every  $(x, s) \in \Omega \times \mathbb{R}$ .

Let us define  $\mathcal{F} : H_0^1(\Omega) \rightarrow \mathbb{R}$  by  $\mathcal{F}(u) = \int_{\Omega} F(x, u(x))dx$ . From condition **(f1)** it follows that  $\mathcal{F}$  is of class  $C^1$  and

$$\mathcal{F}'(u)(v) = \int_{\Omega} f(x, u(x))v(x)dx.$$

We consider the indicator function of the set  $\mathcal{K}$ , that is,  $\zeta_{\mathcal{K}} : H_0^1(\Omega) \rightarrow ]-\infty, \infty]$  defined by

$$\zeta_{\mathcal{K}}(u) = \begin{cases} 0, & \text{if } u \in \mathcal{K} \\ +\infty, & \text{if } u \notin \mathcal{K}. \end{cases}$$

Then  $\zeta_{\mathcal{K}}$  is convex, proper and lower semi-continuous. Moreover, define for  $\lambda > 0$  the function  $I_\lambda : H_0^1(\Omega) \rightarrow ]-\infty, \infty]$  by

$$I_\lambda(u) = \frac{1}{2} \|u\|_{H_0^1}^2 - \lambda \mathcal{F}(u) + \zeta_{\mathcal{K}}(u). \quad (2.2)$$

Then  $I_\lambda$  is a Szulkin-type functional. The relationship between solutions of problem (P) and  $I_\lambda$  is pointed out in what follows.

**Remark 2.2** *If  $u \in H_0^1(\Omega)$  is a critical point of  $I_\lambda$ , then  $(u, \lambda)$  is a solution of (P).*

Let  $G = id_{\mathbb{R}^m} \times O(l) \subset O(m+l)$ . Define the action of  $G$  on  $H_0^1(\Omega)$  by  $gu(x) = u(g^{-1}x)$  for every  $g \in G$ ,  $u \in H_0^1(\Omega)$  and  $x \in \Omega$ . Since  $\mathcal{K}$  is a  $G$ -invariant set, then  $\zeta_{\mathcal{K}}$  is a  $G$ -invariant function. Since  $F(\cdot, s)$  is axially symmetric for every  $s \in \mathbb{R}$ , then  $\mathcal{F}$  is also a  $G$ -invariant function. The norm  $\|\cdot\|_{H_0^1}$  is a  $G$ -invariant function as well. In conclusion, if we consider the set

$$\Sigma = H_{0,s}^1(\Omega) = \{u \in H_0^1(\Omega) : gu = u \text{ for every } g \in G\},$$

then, in view of Theorem 1.31, every critical point of  $I_\lambda|_\Sigma$  becomes as well critical point of  $I_\lambda$ .

We apply Theorem 1.26 for

$$X = \Sigma = H_{0,s}^1(\Omega), \quad E_1 = \frac{1}{2} \|\cdot\|_\Sigma^2, \quad \zeta_1 = \zeta_{\mathcal{K}}|_\Sigma, \quad E_2 = -\mathcal{F}|_\Sigma, \quad \Lambda = [0, \infty[.$$

As usual,  $\|\cdot\|_\Sigma$ ,  $\zeta_{\mathcal{K}}|_\Sigma$  and  $\mathcal{F}|_\Sigma$  denote the restrictions of  $\|\cdot\|_{H_0^1}$ ,  $\zeta_{\mathcal{K}}$  and  $\mathcal{F}$  to  $\Sigma$ , respectively. We also use the notation  $\langle \cdot, \cdot \rangle_\Sigma$  for the restriction of  $\langle \cdot, \cdot \rangle_{H_0^1}$  to  $\Sigma$  and denote by  $\langle \cdot, \cdot \rangle_\Sigma$  the duality mapping restricted to  $\Sigma \times \Sigma^*$ .

Now, we are going to verify the hypotheses (i)-(iii) of Theorem 1.26.

**Step 1.** (Verification of (i)). The weakly sequentially lower semi-continuity of  $E_1$  is standard. We prove that  $E_2$  is weakly sequentially continuous.

Let  $\{u_n\}$  be a sequence from  $\Sigma$  which converges weakly to some  $u \in \Sigma$ . In particular,  $\{u_n\}$  is bounded in  $\Sigma$  and by virtue of Lemma 2.1,  $F(x, s) = o(s^2)$  as  $s \rightarrow 0$ , and  $F(x, s) = o(s^{2^*})$  as  $s \rightarrow +\infty$ , uniformly for every  $x \in \Omega$ . But, from [109, Lemma 4, p. 368], it follows that  $E_2(u_n) \rightarrow E_2(u)$  as  $n \rightarrow \infty$ , hence  $E_2$  is weakly sequentially continuous.

**Step 2.** (Verification of (ii)). Fix  $\lambda \in \Lambda$ . First, we prove that  $I_1 +$

$\lambda I_2 \equiv E_1 + \zeta_1 + \lambda E_2$  is coercive. Indeed, due to **(f3)** and using Hölder's inequality, we have for every  $u \in \Sigma$

$$\begin{aligned} I_1(u) + \lambda I_2(u) &\geq \frac{1}{2} \|u\|_{\Sigma}^2 - \lambda \int_{\Omega} \alpha(x) |u(x)|^q dx - \lambda \int_{\Omega} \beta(x) dx \\ &\geq \frac{1}{2} \|u\|_{\Sigma}^2 - \lambda \|\alpha\|_{\nu/(\nu-q)} k_{\nu}^q \|u\|_{\Sigma}^q - \lambda \|\beta\|_1. \end{aligned}$$

Since  $q < 2$ , then  $\|u\|_{\Sigma} \rightarrow +\infty$ , hence  $I_1(u) + \lambda I_2(u) \rightarrow +\infty$ , as claimed.

Now, we prove that  $I_1 + \lambda I_2$  verifies condition  $(PSZ)_c$ , for all  $c \in \mathbb{R}$ . Let  $\{u_n\} \subset \Sigma$  be a sequence such that

$$I_1(u_n) + \lambda I_2(u_n) \rightarrow c \quad (2.3)$$

and for every  $v \in \Sigma$  we have

$$\langle u_n, v - u_n \rangle_{\Sigma} + \lambda \langle E_2'(u_n), v - u_n \rangle_{\Sigma} + \zeta_1(v) - \zeta_1(u_n) \geq -\varepsilon_n \|v - u_n\|_{\Sigma}, \quad (2.4)$$

for a sequence  $\{\varepsilon_n\}$  in  $[0, +\infty[$  with  $\varepsilon_n \rightarrow 0$ . In particular, (2.3) shows that  $\{u_n\} \subset \mathcal{K}$ . Moreover, the coerciveness of the function  $I_1 + \lambda I_2$  implies that the sequence  $\{u_n\}$  is bounded in  $\Sigma \cap \mathcal{K}$ . Therefore, there exists an element  $u \in \mathcal{K} \cap \Sigma$  such that  $\{u_n\}$  converges weakly to  $u$  in  $\Sigma$ . (Note that  $\mathcal{K}$  is convex and closed, thus, weakly closed.) Moreover, since the embedding  $\Sigma \hookrightarrow L^p(\Omega)$  is compact then, up to a subsequence,  $\{u_n\}$  converges strongly to  $u$  in  $L^p(\Omega)$ . Choosing in particular  $v = u$  in (2.4), we have

$$\|u_n - u\|_{\Sigma}^2 \leq \lambda \langle E_2'(u_n), u - u_n \rangle_{\Sigma} + \langle u, u - u_n \rangle_{\Sigma} + \varepsilon_n \|u - u_n\|_{\Sigma}.$$

The last two terms tend to zero as  $n \rightarrow \infty$ . Thus, in order to prove  $\|u_n - u\|_{\Sigma} \rightarrow 0$ , it is enough to show that the first term in the right hand side tends to zero as well. From Remark 2.1 a), we obtain

$$\begin{aligned} \langle E_2'(u_n), u - u_n \rangle_{\Sigma} &= \int_{\Omega} f(x, u_n(x)) (-u(x) + u_n(x)) dx \\ &\leq \int_{\Omega} [\varepsilon |u_n(x)| + c(\varepsilon) |u_n(x)|^{p-1}] |u_n(x) - u(x)| dx \\ &\leq \varepsilon k_2^2 \|u_n\|_{\Sigma} \|u_n - u\|_{\Sigma} + c(\varepsilon) \|u_n\|_p^{p-1} \|u_n - u\|_p. \end{aligned}$$

Due to the arbitrariness of  $\varepsilon > 0$ , the last term tends to zero, therefore,  $\|u_n - u\|_{\Sigma} \rightarrow 0$  as  $n \rightarrow \infty$ .

**Step 3.** (Verification of (iii)). Let us define the function  $\gamma : [0, \infty[ \rightarrow \mathbb{R}$  by

$$\gamma(s) = \sup\{-E_2(u) : \|u\|_{\Sigma}^2 \leq 2s\}.$$

Due to Remark 2.1 (ii), we have

$$\gamma(s) \leq 2\varepsilon k_2^2 s + 2^p c(\varepsilon) k_p^p s^{\frac{p}{2}}.$$

On the other hand, we know that  $\gamma(s) \geq 0$  for  $s \geq 0$ . Due to the arbitrariness of  $\varepsilon > 0$ , we deduce

$$\lim_{s \rightarrow 0^+} \frac{\gamma(s)}{s} = 0.$$

By **(f4)** it is clear that  $u_0 \neq 0$  ( $E_2(0) = 0$ ). Therefore it is possible to choose a number  $\eta$  such that

$$0 < \eta < -2E_2(u_0)\|u_0\|_{\Sigma}^{-2}.$$

From  $\lim_{s \rightarrow 0^+} \gamma(s)/s = 0$  follows the existence of a number  $s_0 \in ]0, \|u_0\|_{\Sigma}^2/2[$  such that  $\gamma(s_0) < \eta s_0$ . Therefore,

$$\gamma(s_0) < -2E_2(u_0)\|u_0\|_{\Sigma}^{-2} s_0.$$

Choose  $\rho_0 > 0$  such that

$$\gamma(s_0) < \rho_0 < -2E_2(u_0)\|u_0\|_{\Sigma}^{-2} s_0. \quad (2.5)$$

Due to the choice of  $t_0$  and (2.5) we have

$$\rho_0 < -E_2(u_0). \quad (2.6)$$

Define  $h : \Lambda = [0, +\infty[ \rightarrow \mathbb{R}$  by  $h(\lambda) = \rho_0 \lambda$ . We prove that the function  $h$  satisfies the inequality

$$\sup_{\lambda \in \Lambda} \inf_{u \in \Sigma} (I_1(u) + \lambda I_2(u) + \rho_0 \lambda) < \inf_{u \in \Sigma} \sup_{\lambda \in \Lambda} (I_1(u) + \lambda I_2(u) + \rho_0 \lambda).$$

Note, that in the previous inequality we can put  $\Sigma \cap \mathcal{K}$  instead of  $\Sigma$ ; indeed, if  $u \in \Sigma \setminus \mathcal{K}$ , then  $I_1(u) = +\infty$ .

The function

$$\Lambda \ni \lambda \mapsto \inf_{u \in \Sigma \cap \mathcal{K}} [\|u\|_{\Sigma}^2/2 + \lambda(\rho_0 + E_2(u))]$$

is upper semi-continuous on  $\Lambda$ . Relation (2.6) implies that

$$\begin{aligned} \lim_{\lambda \rightarrow +\infty} \inf_{u \in \Sigma \cap \mathcal{K}} [I_1(u) + \lambda I_2(u) + \rho_0 \lambda] &\leq \\ \lim_{\lambda \rightarrow +\infty} [\|u_0\|_{\Sigma}^2/2 + \lambda(\rho_0 + E_2(u_0))] &= -\infty. \end{aligned}$$

Thus, we find an element  $\bar{\lambda} \in \Lambda$  such that

$$\begin{aligned} \sup_{\lambda \in \Lambda} \inf_{u \in \Sigma \cap \mathcal{K}} (I_1(u) + \lambda I_2(u) + \rho_0 \lambda) = \\ \inf_{u \in \Sigma \cap \mathcal{K}} [\|u\|_{\Sigma}^2/2 + \bar{\lambda}(\rho_0 + E_2(u))] . \end{aligned} \quad (2.7)$$

Since  $\gamma(s_0) < \rho_0$ , for all  $u \in \Sigma$  such that  $\|u\|_{\Sigma}^2 \leq 2s_0$ , we have  $E_2(u) > -\rho_0$ . Thus,

$$\begin{aligned} s_0 &\leq \inf\{\|u\|_{\Sigma}^2/2 : E_2(u) \leq -\rho_0\} \\ &\leq \inf\{\|u\|_{\Sigma}^2/2 : u \in \mathcal{K}, E_2(u) \leq -\rho_0\} . \end{aligned} \quad (2.8)$$

On the other hand,

$$\begin{aligned} \inf_{u \in \Sigma \cap \mathcal{K}} \sup_{\lambda \in \Lambda} (I_1(u) + \lambda I_2(u) + \rho_0 \lambda) &= \inf_{u \in \Sigma \cap \mathcal{K}} \left[ \|u\|_{\Sigma}^2/2 + \sup_{\lambda \in \Lambda} (\lambda(\rho_0 + E_2(u))) \right] \\ &= \inf_{u \in \Sigma \cap \mathcal{K}} \{ \|u\|_{\Sigma}^2/2 : E_2(u) \leq -\rho_0 \} . \end{aligned}$$

Therefore, relation (2.8) can be written as

$$s_0 \leq \inf_{u \in \Sigma \cap \mathcal{K}} \sup_{\lambda \in \Lambda} (I_1(u) + \lambda I_2(u) + \rho_0 \lambda). \quad (2.9)$$

There are two distinct cases:

(A) If  $0 \leq \bar{\lambda} < s_0/\rho_0$ , we have

$$\inf_{u \in \Sigma \cap \mathcal{K}} [\|u\|_{\Sigma}^2/2 + \bar{\lambda}(\rho_0 + E_2(u))] \leq \bar{\lambda}(\rho_0 + E_2(0)) = \bar{\lambda}\rho_0 < s_0.$$

Combining this inequality with (2.7) and (2.9) we obtain the desired inequality.

(B) If  $s_0/\rho_0 \leq \bar{\lambda}$ , from  $\rho_0 < -E_2(u_0)$  and (2.5) we find

$$\begin{aligned} \inf_{u \in \Sigma \cap \mathcal{K}} [\|u\|_{\Sigma}^2/2 + \bar{\lambda}(\rho_0 + E_2(u))] &\leq \|u_0\|_{\Sigma}^2/2 + \bar{\lambda}(\rho_0 + E_2(u_0)) \\ &\leq \|u_0\|_{\Sigma}^2/2 + s_0(\rho_0 + E_2(u_0))/\rho_0 < s_0. \end{aligned}$$

Now, we repeat the last part of (A), which concludes Step 3.

*Proof of Theorem 2.1.* Due to the above three steps, Theorem 1.26 implies the existence of an open interval  $\Lambda_0 \subset [0, \infty[$ , such that for each  $\lambda \in \Lambda_0$ , the function  $I_{\lambda}|_{\Sigma} \equiv I_1 + \lambda I_2$  has at least three critical points in  $\Sigma \cap \mathcal{K}$ . It remains to apply Theorem 1.31 and Remark 2.2.  $\square$

### 2.2.2 One-dimensional case

In this section we continue our studies on the strip-like domains, but contrary to the previous section, we consider domains of the form  $\Omega = \omega \times \mathbb{R}$ , where  $\omega \subset \mathbb{R}^m$  ( $m \geq 1$ ) is a bounded open subset.

J.-L. Lions [100] observed that defining the closed convex cone

$$\mathcal{K} = \{u \in H_0^1(\omega \times \mathbb{R}) : u \text{ is nonnegative,} \\ y \mapsto u(x, y) \text{ is nonincreasing for } x \in \omega, y \geq 0, \quad (\text{K}) \\ y \mapsto u(x, y) \text{ is nondecreasing for } x \in \omega, y \leq 0\},$$

then bounded subsets of  $\mathcal{K}$  are relatively compact in  $L^p(\omega \times \mathbb{R})$  whenever  $p \in (2, 2^*)$ . Note that  $2^* = \infty$ , if  $m = 1$ .

The main goal of this section is to give a new approach to treat stationary eigenvalue problems on domains of the type  $\Omega = \omega \times \mathbb{R}$ . The genesis of our method relies on the Szulkin-type functionals. Indeed, since the indicator function of a closed convex subset of a vector space (so, in particular  $\mathcal{K}$  in  $H_0^1(\omega \times \mathbb{R})$ ) is convex, lower semi-continuous and proper, this approach arises in a natural manner. In order to formulate our problem, we consider a continuous function  $f : (\omega \times \mathbb{R}) \times \mathbb{R} \rightarrow \mathbb{R}$  such that

(F<sub>1</sub>)  $f(x, 0) = 0$ , and there exist  $c_1 > 0$  and  $p \in (2, 2^*)$  such that

$$|f(x, s)| \leq c_1(|s| + |s|^{p-1}), \quad \forall (x, s) \in (\omega \times \mathbb{R}) \times \mathbb{R}.$$

Let  $a \in L^1(\omega \times \mathbb{R}) \cap L^\infty(\omega \times \mathbb{R})$  with  $a \geq 0$ ,  $a \not\equiv 0$ , and  $q \in (1, 2)$ . For  $\lambda > 0$ , we denote by (P<sub>λ</sub>) the following *variational inequality problem*:

Find  $u \in \mathcal{K}$  such that for all  $v \in \mathcal{K}$ ,

$$\int_{\omega \times \mathbb{R}} \nabla u \cdot \nabla(v - u) \, dx + \int_{\omega \times \mathbb{R}} f(x, u)(-v + u) \, dx \\ \geq \lambda \int_{\omega \times \mathbb{R}} a(x)|u|^{q-2}u(v - u) \, dx.$$

For the sake of notation, we introduce  $\Omega = \omega \times \mathbb{R}$ .

Define  $F : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$  by  $F(x, s) = \int_0^s f(x, t)dt$  and beside of (F<sub>1</sub>)

we make the following assumptions:

(F<sub>2</sub>)  $\lim_{s \rightarrow 0} \frac{f(x, s)}{s} = 0$  uniformly for every  $x \in \Omega$ ;

(F<sub>3</sub>) There exists  $\nu > 2$  such that

$$\nu F(x, s) - s f(x, s) \leq 0, \quad \forall (x, s) \in \Omega \times \mathbb{R};$$

(F<sub>4</sub>) There exists  $R > 0$  such that

$$\inf\{F(x, s) : (x, |s|) \in \omega \times [R, \infty)\} > 0.$$

**Lemma 2.1** *If the functions  $f, F : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$  satisfy  $(F_1)$ ,  $(F_3)$  and  $(F_4)$  then there exist  $c_2, c_3 > 0$  such that for all  $(x, s) \in \Omega \times \mathbb{R}$ ,*

$$F(x, s) \geq c_2|s|^\nu - c_3s^2.$$

*Proof* For arbitrary fixed  $(x, u) \in \Omega \times \mathbb{R}$  we consider the function  $g : (0, +\infty) \rightarrow \mathbb{R}$  defined by

$$g(s) = s^{-\nu}F(x, su).$$

Clearly,  $g$  is a function of class  $\mathcal{C}^1$  and we have

$$g'(s) = -\nu s^{-\nu-1}F(x, su) + s^{-\nu}uf(x, su), \quad s > 0.$$

Let  $s > 1$  and by mean value theorem, there exist  $\tau = \tau(x, u) \in (1, s)$  such that  $g(s) - g(1) = g'(\tau)(s - 1)$ . Therefore,

$$g'(\tau) = -\nu\tau^{-\nu-1}F(x, \tau u) + \tau^{-\nu}uf(x, \tau u).$$

Therefore

$$g(s) - g(1) = -\tau^{-\nu-1}[\nu F(x, \tau u) - \tau uf(x, \tau u)](s - 1).$$

By  $(F_3)$  one has  $g(s) \geq g(1)$ , hence  $F(x, su) \geq s^\nu F(x, u)$ , for every  $s \geq 1$ . Let  $c_R = \inf\{F(x, s) : (x, |s|) \in \omega \times [R, \infty)\}$ , which is a strictly positive number, due to  $(F_4)$ . Combining the above facts we obtain

$$F(x, s) \geq \frac{c_R}{R^\nu}|s|^\nu, \quad \forall (x, s) \in \Omega \times \mathbb{R} \text{ with } |s| \geq R. \quad (2.10)$$

On the other hand, by  $((F_1))$  we have  $|F(x, s)| \leq c_1(s^2 + |s|^p)$  for every  $(x, s) \in \Omega \times \mathbb{R}$ . In particular, we have

$$-F(x, s) \leq c_1(s^2 + |s|^p) \leq c_1(1 + R^{p-2} + R^{\nu-2})s^2 - c_1|s|^\nu$$

for every  $(x, s) \in \Omega \times \mathbb{R}$  with  $|s| \leq R$ . Combining the above inequality with (2.10), the desired inequality yields if one chooses  $c_2 = \min\{c_1, c_R/R^\nu\}$  and  $c_3 = c_1(1 + R^{p-2} + R^{\nu-2})$ .  $\square$

**Remark 2.3** In particular, from Lemma 2.1 we observe that  $2 < \nu < p$ .

To investigate the existence of solutions of  $(P_\lambda)$  we construct a functional  $I_\lambda : H_0^1(\Omega) \rightarrow \mathbb{R}$  associated to  $(P_\lambda)$  defined by

$$I_\lambda(u) = \frac{1}{2} \int_\Omega |\nabla u|^2 - \int_\Omega F(x, u(x)) dx - \frac{\lambda}{q} \int_\Omega a(x)|u|^q + \zeta_{\mathcal{K}}(u),$$

where  $\zeta_{\mathcal{K}}$  is the indicator function of the set  $\mathcal{K}$ .

If we consider the function  $\mathcal{F} : H_0^1(\Omega) \rightarrow \mathbb{R}$  defined by

$$\mathcal{F}(u) = \int_{\Omega} F(x, u(x)) dx,$$

then  $\mathcal{F}$  is of class  $\mathcal{C}^1$  and

$$\langle \mathcal{F}'(u), v \rangle_{H_0^1} = \int_{\Omega} f(x, u(x)) v(x) dx, \quad \forall u, v \in H_0^1(\Omega).$$

Then the functionals  $A_1, A_2 : H_0^1(\Omega) \rightarrow \mathbb{R}$  defined by  $A_1(u) = \|u\|_{H_0^1}^2$  and  $A_2(u) = \int_{\Omega} a(x)|u|^q dx$  are of class  $\mathcal{C}^1$ , with derivatives

$$\langle A_1'(u), v \rangle_{H_0^1} = 2\langle u, v \rangle_{H_0^1}$$

and

$$\langle A_2'(u), v \rangle_{H_0^1} = q \int_{\Omega} a(x)|u|^{q-2} uv dx.$$

Therefore the functional

$$E_{\lambda}(u) = \frac{1}{2} \|u\|_{H_0^1}^2 - \frac{\lambda}{q} \int_{\Omega} a(x)|u|^q - \mathcal{F}(u)$$

defined on  $H_0^1(\Omega)$  is of class  $\mathcal{C}^1$ . On the other hand, the indicator function of the set  $\mathcal{K}$  defined by

$$\zeta_{\mathcal{K}}(u) = \begin{cases} 0, & \text{if } u \in \mathcal{K}, \\ +\infty, & \text{if } u \notin \mathcal{K}, \end{cases}$$

is convex, proper, and lower semi-continuous. In conclusion,  $I_{\lambda} = E_{\lambda} + \zeta_{\mathcal{K}}$  is a Szulkin-type functional.

**Proposition 2.1** *Fix  $\lambda > 0$  arbitrary. Then every critical point  $u \in H_0^1(\Omega)$  of  $I_{\lambda} = E_{\lambda} + \zeta_{\mathcal{K}}$  (in the sense of Szulkin) is a solution of  $(P_{\lambda})$ .*

*Proof* Since  $u \in H_0^1(\Omega)$  is a critical point of  $I_{\lambda} = E_{\lambda} + \zeta_{\mathcal{K}}$ , one has

$$\langle E_{\lambda}'(u), v - u \rangle_{H_0^1} + \zeta_{\mathcal{K}}(v) - \zeta_{\mathcal{K}}(u) \geq 0, \quad \forall v \in H_0^1(\Omega).$$

We deduce immediately that  $u \in \mathcal{K}$ . Otherwise, we would have  $\zeta_{\mathcal{K}}(u) = +\infty$  which led us to a contradiction, letting for instance  $v = 0 \in \mathcal{K}$  in the above inequality. Now, we fix  $v \in \mathcal{K}$  arbitrary and we obtain the desired inequality.  $\square$

**Remark 2.4** We observe that  $0 \in \mathcal{K}$  is a trivial solution of  $(P_{\lambda})$  for every  $\lambda \in \mathbb{R}$ .

**Proposition 2.2** *If the conditions  $(F_1) - (F_3)$  hold, then  $I_{\lambda} = E_{\lambda} + \zeta_{\mathcal{K}}$  satisfies  $(PSZ)_c$ -condition for every  $c \in \mathbb{R}$  and  $\lambda > 0$ .*

*Proof* Let  $\lambda > 0$  and  $c \in \mathbb{R}$  be some fixed numbers and let  $\{u_n\}$  be a sequence from  $H_0^1(\Omega)$  such that for all  $v \in H_0^1(\Omega)$ ,

$$I_\lambda(u_n) = E_\lambda(u_n) + \zeta_{\mathcal{K}}(u_n) \rightarrow c; \quad (2.11)$$

$$\langle E'_\lambda(u_n), v - u_n \rangle_{H_0^1} + \zeta_{\mathcal{K}}(v) - \zeta_{\mathcal{K}}(u_n) \geq -\varepsilon_n \|v - u_n\|_{H_0^1}, \quad (2.12)$$

for a sequence  $\{\varepsilon_n\}$  in  $[0, \infty)$  with  $\varepsilon_n \rightarrow 0$ . By (2.11) one concludes that the sequence  $\{u_n\}$  belongs entirely to  $\mathcal{K}$ . Setting  $v = 2u_n$  in (2.12), we obtain

$$\langle E'_\lambda(u_n), u_n \rangle_{H_0^1} \geq -\varepsilon_n \|u_n\|_{H_0^1}.$$

From the above inequality we derive

$$\|u_n\|_{H_0^1}^2 - \lambda \int_{\Omega} a(x)|u_n|^q - \int_{\Omega} f(x, u_n(x))u_n(x)dx \geq -\varepsilon_n \|u_n\|_{H_0^1}. \quad (2.13)$$

By (2.11) we deduce that, for large  $n$ ,

$$c + 1 \geq \frac{1}{2} \|u_n\|_{H_0^1}^2 - \frac{\lambda}{q} \int_{\Omega} a(x)|u_n|^q - \int_{\Omega} F(x, u_n(x))dx \quad (2.14)$$

Multiplying (2.13) by  $\nu^{-1}$  and adding this one to (2.14), by Hölder's inequality we have for large  $n \in \mathbb{N}$

$$\begin{aligned} c + 1 + \frac{1}{\nu} \|u_n\|_{H_0^1} &\geq \left(\frac{1}{2} - \frac{1}{\nu}\right) \|u_n\|_{H_0^1}^2 - \lambda \left(\frac{1}{q} - \frac{1}{\nu}\right) \int_{\Omega} a(x)|u_n|^q \\ &\quad - \frac{1}{\nu} \int_{\Omega} [\nu F(x, u_n(x)) - u_n(x)f(x, u_n(x))]dx \\ &\stackrel{(F_3)}{\geq} \left(\frac{1}{2} - \frac{1}{\nu}\right) \|u_n\|_{H_0^1}^2 - \lambda \left(\frac{1}{q} - \frac{1}{\nu}\right) \|a\|_{\nu/(\nu-q)} \|u_n\|_{\nu}^q \\ &\geq \left(\frac{1}{2} - \frac{1}{\nu}\right) \|u_n\|_{H_0^1}^2 - \lambda \left(\frac{1}{q} - \frac{1}{\nu}\right) \|a\|_{\nu/(\nu-q)} k_{\nu}^q \|u_n\|_{H_0^1}^q. \end{aligned}$$

In the above inequalities we used the Remark 2.3 and the hypothesis  $a \in L^1(\Omega) \cap L^\infty(\Omega)$  thus, in particular,  $a \in L^{\nu/(\nu-q)}(\Omega)$ . Since  $q < 2 < \nu$ , from the above estimation we derive that the sequence  $\{u_n\}$  is bounded in  $\mathcal{K}$ . Therefore,  $\{u_n\}$  is relatively compact in  $L^p(\Omega)$ ,  $p \in (2, 2^*)$ . Up to a subsequence, we can suppose that

$$u_n \rightharpoonup u \text{ weakly in } H_0^1(\Omega); \quad (2.15)$$

$$u_n \rightarrow u \text{ strongly in } L^\mu(\Omega), \mu \in (2, 2^*). \quad (2.16)$$

Since  $\mathcal{K}$  is (weakly) closed then  $u \in \mathcal{K}$ . Setting  $v = u$  in (2.12), we have

$$\begin{aligned} & \langle u_n, u - u_n \rangle_{H_0^1} + \int_{\Omega} f(x, u_n(x))(u_n(x) - u(x)) dx \\ & - \lambda \int_{\Omega} a(x) |u_n|^{q-2} u_n (u - u_n) \geq -\varepsilon_n \|u - u_n\|_{H_0^1}. \end{aligned}$$

Therefore, in view of Remark 2.1 (i) we derive

$$\begin{aligned} \|u - u_n\|_{H_0^1}^2 & \leq \langle u, u - u_n \rangle_{H_0^1} + \int_{\Omega} f(x, u_n(x))(u_n(x) - u(x)) dx \\ & \quad - \lambda \int_{\Omega} a(x) |u_n|^{q-2} u_n (u - u_n) + \varepsilon_n \|u - u_n\|_{H_0^1} \\ & \leq \langle u, u - u_n \rangle_{H_0^1} + \lambda \|a\|_{\nu/(p-q)} \|u_n\|_{\nu}^{q-1} \|u - u_n\|_{\nu} \\ & \quad + \varepsilon_n \|u - u_n\|_{H_0^1} + \varepsilon \|u_n\|_{H_0^1} \|u_n - u\|_{H_0^1} \\ & \quad + c(\varepsilon) \|u_n\|_p^{p-1} \|u_n - u\|_p, \end{aligned}$$

where  $\varepsilon > 0$  is arbitrary small. Taking into account relations (2.15) and (2.16), the facts that  $\nu, p \in (2, 2^*)$ , the arbitrariness of  $\varepsilon > 0$  and  $\varepsilon_n \rightarrow 0^+$ , one has that  $\{u_n\}$  converges strongly to  $u$  in  $H_0^1(\Omega)$ . This completes the proof.  $\square$

**Proposition 2.3** *If the conditions  $(F_1) - (F_4)$  are verified, then there exists a  $\lambda_0 > 0$  such that for every  $\lambda \in (0, \lambda_0)$  the function  $I_{\lambda}$  has mountain pass geometry, that is, the following assertions are true:*

- (i). *there exist constants  $\alpha_{\lambda} > 0$  and  $\rho_{\lambda} > 0$  such that  $I_{\lambda}(u) \geq \alpha_{\lambda}$  for all  $\|u\|_{H_0^1} = \rho_{\lambda}$ ;*
- (ii). *there exists  $e_{\lambda} \in H_0^1(\Omega)$  with  $\|e_{\lambda}\|_{H_0^1} > \rho_{\lambda}$  and  $I_{\lambda}(e_{\lambda}) \leq 0$ .*

*Proof* (i) Due to Remark 2.1 (ii), for every  $\varepsilon > 0$  there exists  $c(\varepsilon) > 0$  such that  $\mathcal{F}(u) \leq \varepsilon \|u\|_{H_0^1}^2 + c(\varepsilon) \|u\|_p^p$  for every  $u \in H_0^1(\Omega)$ . It suffices to restrict our attention to elements  $u$  which belong to  $\mathcal{K}$ ; otherwise  $I_{\lambda}(u)$  will be  $+\infty$ , hence (i) holds trivially. Fix  $\varepsilon_0 \in (0, \frac{1}{2})$ . One has

$$I_{\lambda}(u) \geq \left( \frac{1}{2} - \varepsilon_0 \right) \|u\|_{H_0^1}^2 - k_p^p c(\varepsilon_0) \|u\|_{H_0^1}^p \quad (2.17)$$

$$- \frac{\lambda k_p^q}{q} \|a\|_{p/(p-q)} \|u\|_{H_0^1}^q \quad (2.18)$$

$$= (A - B \|u\|_{H_0^1}^{p-2} - \lambda C \|u\|_{H_0^1}^{q-2}) \|u\|_{H_0^1}^2, \quad (2.19)$$

where  $A = (\frac{1}{2} - \varepsilon_0) > 0$ ,  $B = k_p^p c(\varepsilon_0) > 0$  and  $C = k_p^q \|a\|_{p/(p-q)}/q > 0$ .

For every  $\lambda > 0$ , let us define a function  $g_\lambda : (0, \infty) \rightarrow \mathbb{R}$  by

$$g_\lambda(s) = A - Bs^{p-2} - \lambda Cs^{q-2}.$$

Clearly,  $g'_\lambda(s_\lambda) = 0$  if and only if  $s_\lambda = (\lambda \frac{2-q}{p-2} \frac{C}{B})^{\frac{1}{p-q}}$ . Moreover,  $g_\lambda(s_\lambda) = A - D\lambda^{\frac{p-2}{p-q}}$ , where  $D = D(p, q, B, C) > 0$ . Choosing  $\lambda_0 > 0$  such that  $g_{\lambda_0}(s_{\lambda_0}) > 0$ , we deduce that  $g_\lambda(s_\lambda) > 0$  for all  $\lambda \in (0, \lambda_0)$ . Therefore, for every  $\lambda \in (0, \lambda_0)$ , setting  $\rho_\lambda = s_\lambda$  and  $\alpha_\lambda = g_\lambda(s_\lambda)s_\lambda^2$ , the assertion from (i) holds true.

(ii) By Lemma 2.1 we have  $\mathcal{F}(u) \geq c_2\|u\|_\nu^\nu - c_3\|u\|_2^2$  for every  $u \in H_0^1(\Omega)$ . Let us fix  $u \in \mathcal{K}$ . Then we have

$$I_\lambda(u) \leq \left(\frac{1}{2} + c_3k_2^2\right) \|u\|_{H_0^1}^2 - c_2\|u\|_\nu^\nu + \frac{\lambda}{q} \|a\|_{\nu/(\nu-q)} k_\nu^q \|u\|_{H_0^1}^q. \quad (2.20)$$

Fix  $u_0 \in \mathcal{K} \setminus \{0\}$ . Letting  $u = su_0$  ( $s > 0$ ) in (2.20), we have  $I_\lambda(su_0) \rightarrow -\infty$  as  $s \rightarrow +\infty$ , since  $\nu > 2 > q$ . Thus, for every  $\lambda \in (0, \lambda_0)$ , it is possible to set  $s = s_\lambda$  so large that for  $e_\lambda = s_\lambda u_0$ , we have  $\|e_\lambda\|_{H_0^1} > \rho_\lambda$  and  $I_\lambda(e_\lambda) \leq 0$ . This ends the proof of the proposition.  $\square$

The main result of this section is the following.

**Theorem 2.2** *Let  $f : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$  be a function which satisfies  $(F_1)$ - $(F_4)$ . Then there exists  $\lambda_0 > 0$  such that  $(P_\lambda)$  has at least two nontrivial, distinct solutions  $u_\lambda^1, u_\lambda^2 \in \mathcal{K}$  whenever  $\lambda \in (0, \lambda_0)$ .*

*Proof* In the first step we prove the existence of the first nontrivial solution of  $(P_\lambda)$ . By Proposition 2.2, the functional  $I_\lambda$  satisfies  $(PSZ)_c$ -condition for every  $c \in \mathbb{R}$  and clearly  $I_\lambda(0) = 0$  for every  $\lambda > 0$ . Let us fix  $\lambda \in (0, \lambda_0)$ ,  $\lambda_0$  being from Proposition 2.3. It follows that there are constants  $\alpha_\lambda, \rho_\lambda > 0$  and  $e_\lambda \in H_0^1(\Omega)$  such that  $I_\lambda$  fulfills the properties (i) and (ii) from Theorem 1.25. Therefore, the number  $c_\lambda^1 = \inf_{\gamma \in \Gamma} \sup_{t \in [0,1]} \mathcal{J}_\lambda(\gamma(t))$ , where  $\Gamma = \{\gamma \in C([0,1], H_0^1(\Omega)) : \gamma(0) = 0, \gamma(1) = e_\lambda\}$ , is a critical value of  $I_\lambda$  with  $c_\lambda^1 \geq \alpha_\lambda > 0$ . It is clear that the critical point  $u_\lambda^1 \in H_0^1(\Omega)$  which corresponds to  $c_\lambda^1$  cannot be trivial since  $I_\lambda(u_\lambda^1) = c_\lambda^1 > 0 = I_\lambda(0)$ . It remains to apply Proposition 2.1 which concludes that  $u_\lambda^1$  is actually an element of  $\mathcal{K}$  and it is a solution of  $(P_\lambda)$ .

In the next step we prove the existence of the second solution of the problem  $(P_\lambda)$ . For this let us fix  $\lambda \in (0, \lambda_0)$  arbitrary,  $\lambda_0$  being from the first step. By Proposition 2.3, there exists  $\rho_\lambda > 0$  such that

$$\inf_{\|u\|_{H_0^1} = \rho_\lambda} I_\lambda(u) > 0. \quad (2.21)$$

On the other hand, since  $a \geq 0$ ,  $a \not\equiv 0$ , there exists  $u_0 \in \mathcal{K}$  such that  $\int_{\Omega} a(x)|u_0(x)|^q dx > 0$ . Thus, for  $t > 0$  small we have

$$I_{\lambda}(tu_0) \leq t^2 \left( \frac{1}{2} + c_3 k_2^2 \right) \|u_0\|_{H_0^1}^2 - c_2 t^{\nu} \|u_0\|_{\nu}^{\nu} - \frac{\lambda}{q} t^q \int_{\Omega} a(x)|u_0(x)|^q dx < 0.$$

For  $r > 0$ , let us denote by  $B_r = \{u \in H_0^1(\Omega) : \|u\|_{H_0^1} \leq r\}$  and  $S_r = \{u \in H_0^1(\Omega) : \|u\|_{H_0^1} = r\}$ . With these notations, relation (2.21) and the above inequality can be summarized as

$$c_{\lambda}^2 = \inf_{u \in B_{\rho_{\lambda}}} I_{\lambda}(u) < 0 < \inf_{u \in S_{\rho_{\lambda}}} I_{\lambda}(u). \quad (2.22)$$

We point out that  $c_{\lambda}^2$  is finite, due to (2.17). Moreover, we will show that  $c_{\lambda}^2$  is another critical point of  $I_{\lambda}$ . To this end, let  $n \in \mathbb{N} \setminus \{0\}$  such that

$$\frac{1}{n} < \inf_{u \in S_{\rho_{\lambda}}} I_{\lambda}(u) - \inf_{u \in B_{\rho_{\lambda}}} I_{\lambda}(u). \quad (2.23)$$

By Ekeland's variational principle applied to the lower semi-continuous functional  $I_{\lambda}|_{B_{\rho_{\lambda}}}$ , which is bounded below (see (2.22)), there is  $u_{\lambda,n} \in B_{\rho_{\lambda}}$  such that

$$I_{\lambda}(u_{\lambda,n}) \leq \inf_{u \in B_{\rho_{\lambda}}} I_{\lambda}(u) + \frac{1}{n}; \quad (2.24)$$

$$I_{\lambda}(w) \geq I_{\lambda}(u_{\lambda,n}) - \frac{1}{n} \|w - u_{\lambda,n}\|_{H_0^1}, \quad \forall w \in B_{\rho_{\lambda}}. \quad (2.25)$$

By (2.23) and (2.24) we have  $I_{\lambda}(u_{\lambda,n}) < \inf_{u \in S_{\rho_{\lambda}}} I_{\lambda}(u)$ , hence  $\|u_{\lambda,n}\|_{H_0^1} < \rho_{\lambda}$ .

Fix an element  $v \in H_0^1(\Omega)$ . It is possible to choose  $t > 0$  so small that  $w = u_{\lambda,n} + t(v - u_{\lambda,n}) \in B_{\rho_{\lambda}}$ . Putting this element into (2.25), using the convexity of  $\zeta_{\mathcal{K}}$  and dividing by  $t > 0$ , one concludes

$$\frac{E_{\lambda}(u_{\lambda,n} + t(v - u_{\lambda,n})) - E_{\lambda}(u_{\lambda,n})}{t} + \zeta_{\mathcal{K}}(v) - \zeta_{\mathcal{K}}(u_{\lambda,n}) \geq -\frac{1}{n} \|v - u_{\lambda,n}\|_{H_0^1}.$$

Letting  $t \rightarrow 0^+$ , we derive

$$\langle E'_{\lambda}(u_{\lambda,n}), v - u_{\lambda,n} \rangle_{H_0^1(\Omega)} + \zeta_{\mathcal{K}}(v) - \zeta_{\mathcal{K}}(u_{\lambda,n}) \geq -\frac{1}{n} \|v - u_{\lambda,n}\|_{H_0^1}. \quad (2.26)$$

By (2.22) and (2.24) we obtain that

$$I_{\lambda}(u_{\lambda,n}) = E_{\lambda}(u_{\lambda,n}) + \zeta_{\mathcal{K}}(u_{\lambda,n}) \rightarrow c_{\lambda}^2 \quad (2.27)$$

as  $n \rightarrow \infty$ . Since  $v$  was arbitrary fixed in (2.26), the sequence  $\{u_{\lambda,n}\}$  fulfills (2.11) and (2.12), respectively. Hence, it is possible to prove in a similar manner as in Proposition 2.2 that  $\{u_{\lambda,n}\}$  contains a convergent

subsequence; denote it again by  $\{u_{\lambda,n}\}$  and its limit point by  $u_\lambda^2$ . It is clear that  $u_\lambda^2$  belongs to  $B_{\rho_\lambda}$ . By the lower semi-continuity of  $\zeta_{\mathcal{K}}$  we have  $\zeta_{\mathcal{K}}(u_\lambda^2) \leq \liminf_{n \rightarrow \infty} \zeta_{\mathcal{K}}(u_{\lambda,n})$ . Combining this inequality with  $\lim_{n \rightarrow \infty} \langle E'_\lambda(u_{\lambda,n}), v - u_{\lambda,n} \rangle_{H_0^1} = \langle E'_\lambda(u_\lambda^2), v - u_\lambda^2 \rangle_{H_0^1}$  and (2.26) we have

$$\langle E'_\lambda(u_\lambda^2), v - u_\lambda^2 \rangle_{H_0^1} + \zeta_{\mathcal{K}}(v) - \zeta_{\mathcal{K}}(u_\lambda^2) \geq 0, \quad \forall v \in H_0^1(\Omega),$$

i.e.  $u_\lambda^2$  is a critical point of  $I_\lambda$ . Moreover,

$$c_\lambda^2 \stackrel{(2.22)}{=} \inf_{u \in B_{\rho_\lambda}} I_\lambda(u) \leq I_\lambda(u_\lambda^2) \leq \liminf_{n \rightarrow \infty} I_\lambda(u_{\lambda,n}) \stackrel{(2.27)}{=} c_\lambda^2,$$

i.e.  $I_\lambda(u_\lambda^2) = c_\lambda^2$ . Since  $c_\lambda^2 < 0$ , it follows that  $u_\lambda^2$  is not trivial. We apply again Proposition 2.1, concluding that  $u_\lambda^2$  is a solution of  $(P_\lambda)$  which differs from  $u_\lambda^1$ . This completes the proof of Theorem 2.2.  $\square$

### 2.3 Area-type variational inequalities

In this section we are interested to obtain existence and multiplicity results for hemivariational inequalities associated with energies which come from the relaxation of functionals

$$f(u) = \int_{\Omega} \sqrt{1 + |\nabla u|^2} dx + \int_{\Omega} G(x, u) dx,$$

where  $u \in W_0^{1,1}(\Omega; \mathbb{R}^N)$ ,  $\Omega$  open in  $\mathbb{R}^n$ ,  $n \geq 2$ . The first feature is that the functional  $f$  does not satisfy the Palais-Smale condition in the space of *functions of bounded variation*

$$BV(\Omega; \mathbb{R}^N) := \left\{ u \in L^1(\Omega); \frac{\partial u}{\partial x_i} \text{ is a measure, for all } i = 1, 2, \dots, n \right\},$$

which is the natural domain of  $f$ . Therefore we extend  $f$  to the function space  $L^{n/(n-1)}(\Omega; \mathbb{R}^N)$  with value  $+\infty$  outside  $BV(\Omega; \mathbb{R}^N)$ . This larger space is better behaved for the compactness properties, but the nonsmoothness of the functional increases. The second feature is that the assumptions we impose on  $G$  imply the second term of  $f$  to be continuous on  $L^{n/(n-1)}(\Omega; \mathbb{R}^N)$ , but not locally Lipschitz. More precisely, the function  $\{s \mapsto G(x, s)\}$  is supposed to be locally Lipschitz for a.e.  $x \in \Omega$ , but the growth conditions we impose do not ensure the corresponding property for the integral on  $L^{n/(n-1)}(\Omega; \mathbb{R}^N)$ .

After recalling the main tools we need in the sequel, we establish some general results for a class of lower semicontinuous functionals  $f : L^p(\Omega; \mathbb{R}^N) \rightarrow \mathbb{R} \cup \{+\infty\}$ . Next, we show that the area-type integrals

fall into the class considered in this section. By the way, we also prove a relation between the convergence in the so-called intermediate topologies of  $BV(\Omega; \mathbb{R}^N)$  and the convergence in  $L^{n/(n-1)}(\Omega; \mathbb{R}^N)$ . Finally, we obtain multiplicity results of Clark and Ambrosetti-Rabinowitz type.

We start with some notions and properties on nonsmooth analysis. We refer to Appendix D for basic definitions and related results.

**Definition 2.1** Let  $c \in \mathbb{R}$ . We say that  $f$  satisfies condition  $(epi)_c$ , if there exists  $\varepsilon > 0$  such that

$$\inf \{ |d\mathcal{G}_f|(u, \lambda) : f(u) < \lambda, |\lambda - c| < \varepsilon \} > 0.$$

The next two results are useful in dealing with condition  $(epi)_c$ .

**Proposition 2.4** Let  $(u, \lambda) \in \text{epi}(f)$ . Assume that there exist positive numbers  $\varrho, \sigma, \delta, \varepsilon$  and a continuous map

$$\mathcal{H} : \{w \in B_\delta(u) : f(w) < \lambda + \delta\} \times [0, \delta] \rightarrow X$$

satisfying

$$d(\mathcal{H}(w, t), w) \leq \varrho t, \quad f(\mathcal{H}(w, t)) \leq \max\{f(w) - \sigma t, \lambda - \varepsilon\}$$

whenever  $w \in B_\delta(u)$ ,  $f(w) < \lambda + \delta$  and  $t \in [0, \delta]$ .

Under these assumptions,

$$|d\mathcal{G}_f|(u, \lambda) \geq \frac{\sigma}{\sqrt{\varrho^2 + \sigma^2}}.$$

If, moreover,  $X$  is a normed space,  $f$  is even,  $u = 0$  and  $\mathcal{H}(-w, t) = -\mathcal{H}(w, t)$ , then

$$|d_{\mathbb{Z}_2}\mathcal{G}_f|(0, \lambda) \geq \frac{\sigma}{\sqrt{\varrho^2 + \sigma^2}}.$$

*Proof* Let  $\delta' \in ]0, \delta]$  be such that  $\delta' + \sigma\delta' \leq \varepsilon$  and let

$$\mathcal{K} : (B_{\delta'}(u, \lambda) \cap \text{epi}(f)) \times [0, \delta'] \rightarrow \text{epi}(f)$$

be defined by  $\mathcal{K}((w, \mu), t) = (\mathcal{H}(w, t), \mu - \sigma t)$ . If  $(w, \mu) \in B_{\delta'}(u, \lambda) \cap \text{epi}(f)$  and  $t \in [0, \delta']$ , we have

$$\lambda - \varepsilon \leq \lambda - \delta' - \sigma\delta' < \mu - \sigma t, \quad f(w) - \sigma t \leq \mu - \sigma t,$$

hence

$$f(\mathcal{K}((w, \mu), t)) \leq \max\{f(w) - \sigma t, \lambda - \varepsilon\} \leq \mu - \sigma t.$$

Therefore  $\mathcal{K}$  actually takes its values in  $\text{epi}(f)$ . Furthermore, it is

$$d(\mathcal{K}((w, \mu), t), (w, \mu)) \leq \sqrt{\varrho^2 + \sigma^2} t,$$

$$\mathcal{G}_f(\mathcal{K}((w, \mu), t)) = \mu - \sigma t = \mathcal{G}_f(w, \mu) - \sigma t.$$

Taking into account Definition D.6, the first assertion follows.

In the symmetric case,  $\mathcal{K}$  automatically satisfies the further condition required in Definition D.14.  $\square$

**Corollary 2.1** *Let  $(u, \lambda) \in \text{epi}(f)$  with  $f(u) < \lambda$ . Assume that for every  $\varrho > 0$  there exist  $\delta > 0$  and a continuous map*

$$\mathcal{H} : \{w \in B_\delta(u) : f(w) < \lambda + \delta\} \times [0, \delta] \rightarrow X$$

*satisfying*

$$d(\mathcal{H}(w, t), w) \leq \varrho t, \quad f(\mathcal{H}(w, t)) \leq f(w) + t(f(u) - f(w) + \varrho)$$

*whenever  $w \in B_\delta(u)$ ,  $f(w) < \lambda + \delta$  and  $t \in [0, \delta]$ .*

*Then we have  $|d\mathcal{G}_f|(u, \lambda) = 1$ . If moreover  $X$  is a normed space,  $f$  is even,  $u = 0$  and  $\mathcal{H}(-w, t) = -\mathcal{H}(w, t)$ , then  $|d_{\mathbb{Z}_2}\mathcal{G}_f|(0, \lambda) = 1$ .*

*Proof* Let  $\varepsilon > 0$  with  $\lambda - 2\varepsilon > f(u)$ , let  $0 < \varrho < \lambda - f(u) - 2\varepsilon$  and let  $\delta$  and  $\mathcal{H}$  be as in the hypothesis. By reducing  $\delta$ , we may also assume that

$$\delta \leq 1, \quad \delta(|\lambda - 2\varepsilon| + |f(u) + \varrho|) \leq \varepsilon.$$

Now consider  $w \in B_\delta(u)$  with  $f(w) < \lambda + \delta$  and  $t \in [0, \delta]$ . If  $f(w) \leq \lambda - 2\varepsilon$ , we have

$$\begin{aligned} f(w) + t(f(u) - f(w) + \varrho) &= (1-t)f(w) + t(f(u) + \varrho) \leq \\ &\leq (1-t)(\lambda - 2\varepsilon) + t(f(u) + \varrho) \leq \\ &\leq \lambda - 2\varepsilon + t|\lambda - 2\varepsilon| + t|f(u) + \varrho| \leq \lambda - \varepsilon, \end{aligned}$$

while, if  $f(w) > \lambda - 2\varepsilon$ , we have

$$f(w) + t(f(u) - f(w) + \varrho) \leq f(w) - (\lambda - f(u) - 2\varepsilon - \varrho)t.$$

In any case it follows

$$f(\mathcal{H}(w, t)) \leq \max \{f(w) - (\lambda - f(u) - 2\varepsilon - \varrho)t, \lambda - \varepsilon\}.$$

From Proposition 2.4 we get

$$|d\mathcal{G}_f|(u, \lambda) \geq \frac{\lambda - f(u) - 2\varepsilon - \varrho}{\sqrt{\varrho^2 + (\lambda - f(u) - 2\varepsilon - \varrho)^2}}$$

and the first assertion follows by the arbitrariness of  $\varrho$ .

The same proof works also in the symmetric case.  $\square$

### 2.3.1 Statement of the problem

Let  $n \geq 1$ ,  $N \geq 1$ ,  $\Omega$  be an open subset of  $\mathbb{R}^n$  and  $1 < p < \infty$ . In the following, we denote by  $\|\cdot\|_q$  the usual norm in  $L^q$  ( $1 \leq q \leq \infty$ ). We now define the functional setting we are interested in.

Let  $\mathcal{E} : L^p(\Omega; \mathbb{R}^N) \rightarrow \mathbb{R} \cup \{+\infty\}$  be a functional such that:

( $\mathcal{E}_1$ )  $\mathcal{E}$  is convex, lower semicontinuous and  $0 \in \mathcal{D}(\mathcal{E})$ , where

$$\mathcal{D}(\mathcal{E}) = \{u \in L^p(\Omega; \mathbb{R}^N) : \mathcal{E}(u) < +\infty\};$$

( $\mathcal{E}_2$ ) there exists  $\vartheta \in C_c(\mathbb{R}^N)$  with  $0 \leq \vartheta \leq 1$  and  $\vartheta(0) = 1$  such that

$$(\mathcal{E}_2.1) \quad \forall u \in \mathcal{D}(\mathcal{E}), \forall v \in \mathcal{D}(\mathcal{E}) \cap L^\infty(\Omega; \mathbb{R}^N), \forall c > 0 :$$

$$\lim_{h \rightarrow \infty} \left[ \sup_{\substack{\|z-u\|_p \leq c \\ \mathcal{E}(z) \leq c}} \mathcal{E} \left( \vartheta \left( \frac{z}{h} \right) v \right) \right] = \mathcal{E}(v);$$

$$(\mathcal{E}_2.2) \quad \forall u \in \mathcal{D}(\mathcal{E}) : \lim_{h \rightarrow \infty} \mathcal{E} \left( \vartheta \left( \frac{u}{h} \right) u \right) = \mathcal{E}(u).$$

Moreover, let  $G : \Omega \times \mathbb{R}^N \rightarrow \mathbb{R}$  be a function such that

( $G_1$ )  $G(\cdot, s)$  is measurable for every  $s \in \mathbb{R}^N$ ;

( $G_2$ ) for every  $t > 0$  there exists  $\alpha_t \in L^1(\Omega)$  such that

$$|G(x, s_1) - G(x, s_2)| \leq \alpha_t(x) |s_1 - s_2|$$

for a.e.  $x \in \Omega$  and every  $s_1, s_2 \in \mathbb{R}^N$  with  $|s_j| \leq t$ ; for a.e.  $x \in \Omega$  we set

$$G^\circ(x, s; \hat{s}) = \gamma^\circ(s; \hat{s}), \quad \partial_s G(x, s) = \partial \gamma(s),$$

where  $\gamma(s) = G(x, s)$ ;

( $G_3$ ) there exist  $a_0 \in L^1(\Omega)$  and  $b_0 \in \mathbb{R}$  such that

$$G(x, s) \geq -a_0(x) - b_0 |s|^p \quad \text{for a.e. } x \in \Omega \text{ and every } s \in \mathbb{R}^N;$$

( $G_4$ ) there exist  $a_1 \in L^1(\Omega)$  and  $b_1 \in \mathbb{R}$  such that

$$G^\circ(x, s; -s) \leq a_1(x) + b_1 |s|^p \quad \text{for a.e. } x \in \Omega \text{ and every } s \in \mathbb{R}^N.$$

Because of  $(\mathcal{E}_1)$  and  $(G_3)$ , we can define a lower semicontinuous functional  $f : L^p(\Omega; \mathbb{R}^N) \rightarrow \mathbb{R} \cup \{+\infty\}$  by

$$f(u) = \mathcal{E}(u) + \int_{\Omega} G(x, u(x)) dx.$$

**Remark 2.1** *According to  $(\mathcal{E}_1)$ , the functional  $\mathcal{E}$  is lower semicontinuous. Condition  $(\mathcal{E}_2)$  ensures that  $\mathcal{E}$  is continuous at least on some particular restrictions.*

**Remark 2.2** *If  $\{s \mapsto G(x, s)\}$  is of class  $C^1$  for a.e.  $x \in \Omega$ , the estimates in  $(G_2)$  and in  $(G_4)$  are respectively equivalent to*

$$|s| \leq t \implies |D_s G(x, s)| \leq \alpha_t(x),$$

$$D_s G(x, s) \cdot s \geq -a_1(x) - b_1|s|^p.$$

Because of  $(G_2)$ , for a.e.  $x \in \Omega$  and any  $t > 0$  and  $s \in \mathbb{R}^N$  with  $|s| < t$  we have

$$\forall \hat{s} \in \mathbb{R}^N : |G^\circ(x, s; \hat{s})| \leq \alpha_t(x)|\hat{s}|; \quad (2.28)$$

$$\forall s^* \in \partial_s G(x, s) : |s^*| \leq \alpha_t(x). \quad (2.29)$$

In the following, we set  $\vartheta_h(s) = \vartheta(s/h)$ , where  $\vartheta$  is a function as in  $(\mathcal{E}_2)$ , and we fix  $M > 0$  such that  $\vartheta = 0$  outside  $B_M(0)$ . Therefore

$$\forall s \in \mathbb{R}^N : |s| \geq hM \implies \vartheta_h(s) = 0. \quad (2.30)$$

We are now concerned with the relationship between the notions of generalized directional derivative and subdifferential in the functional space  $L^p(\Omega; \mathbb{R}^N)$  and the more concrete setting of hemivariational inequalities, which also involves the notion of generalized directional derivative, but in  $\mathbb{R}^N$ .

If  $u, v \in L^p(\Omega; \mathbb{R}^N)$ , we can define  $\int_{\Omega} G^\circ(x, u; v) dx$  as

$$\int_{\Omega} G^\circ(x, u; v) dx = +\infty$$

whenever

$$\int_{\Omega} [G^\circ(x, u; v)]^+ dx = \int_{\Omega} [G^\circ(x, u; v)]^- dx = +\infty.$$

With this convention,  $\{v \mapsto \int_{\Omega} G^\circ(x, u; v) dx\}$  is a convex functional from  $L^p(\Omega; \mathbb{R}^N)$  into  $\overline{\mathbb{R}}$ .

**Theorem 2.3** *Let  $u \in \mathcal{D}(f)$ . Then the following facts hold:*

- (a) for every  $v \in \mathcal{D}(\mathcal{E})$  there exists a sequence  $(v_h)$  in  $\mathcal{D}(\mathcal{E}) \cap L^\infty(\Omega; \mathbb{R}^N)$  satisfying  $[G^\circ(x, u; v_h - u)]^+ \in L^1(\Omega)$ ,  $\|v_h - v\|_p \rightarrow 0$  and  $\mathcal{E}(v_h) \rightarrow \mathcal{E}(v)$ ;
- (b) for every  $v \in \mathcal{D}(\mathcal{E})$  we have

$$f^\circ(u; v - u) \leq \mathcal{E}(v) - \mathcal{E}(u) + \int_{\Omega} G^\circ(x, u; v - u) dx; \quad (2.31)$$

- (c) if  $\partial f(u) \neq \emptyset$ , we have  $G^\circ(x, u; -u) \in L^1(\Omega)$  and

$$\mathcal{E}(v) - \mathcal{E}(u) + \int_{\Omega} G^\circ(x, u; v - u) dx \geq \int_{\Omega} u^* \cdot (v - u) dx \quad (2.32)$$

for every  $u^* \in \partial f(u)$  and  $v \in \mathcal{D}(\mathcal{E})$  (the dual space of  $L^p(\Omega; \mathbb{R}^N)$  is identified with  $L^p'(\Omega; \mathbb{R}^N)$  in the usual way);

- (d) if  $N = 1$ , we have  $[G^\circ(x, u; v - u)]^+ \in L^1(\Omega)$  for every  $v \in L^\infty(\Omega; \mathbb{R}^N)$ .

*Proof* (a) Given  $\varepsilon > 0$ , by  $(\mathcal{E}_2.2)$  we have  $\|\vartheta_h(v)v - v\|_p < \varepsilon$  and  $|\mathcal{E}(\vartheta_h(v)v) - \mathcal{E}(v)| < \varepsilon$  for  $h$  large enough. Then, by  $(\mathcal{E}_2.1)$  we get  $\|\vartheta_k(u)\vartheta_h(v)v - v\|_p < \varepsilon$  and  $|\mathcal{E}(\vartheta_k(u)\vartheta_h(v)v) - \mathcal{E}(v)| < \varepsilon$  for  $k$  large enough. Of course,  $\vartheta_k(u)\vartheta_h(v)v \in L^\infty(\Omega; \mathbb{R}^N)$  and by (2.28) we have

$$\begin{aligned} G^\circ(x, u; \vartheta_k(u)\vartheta_h(v)v - u) &\leq \vartheta_k(u)\vartheta_h(v)G^\circ(x, u; v - u) + \\ &\quad + (1 - \vartheta_k(u)\vartheta_h(v))G^\circ(x, u; -u) \leq \\ &\leq (h + k)M\alpha_{kM}(x) + [G^\circ(x, u; -u)]^+. \end{aligned}$$

From  $(G_4)$  we infer that  $[G^\circ(x, u; -u)]^+ \in L^1(\Omega)$  and assertion (a) follows.

(b) Without loss of generality, we may assume that  $[G^\circ(x, u; v - u)]^+ \in L^1(\Omega)$ . Suppose first that  $v \in \mathcal{D}(\mathcal{E}) \cap L^\infty(\Omega; \mathbb{R}^N)$  and take  $\varepsilon > 0$ .

We claim that for every  $z \in L^p(\Omega; \mathbb{R}^N)$ ,  $t \in ]0, 1/2]$  and  $h \geq 1$  with  $hM > \|v\|_\infty$ , we have

$$\frac{G(x, z + t(\vartheta_h(z)v - z)) - G(x, z)}{t} \leq 2(\|v\|_\infty \alpha_{hM} + a_1 + b_1(|z| + |v|)^p). \quad (2.33)$$

In fact, for a.e.  $x \in \Omega$ , by Lebourg's mean value theorem (see Appendix D) there exist  $\bar{t} \in ]0, t[$  and  $u^* \in \partial_s G(x, z + \bar{t}(\vartheta_h(z)v - z))$  such that

$$\begin{aligned} \frac{G(x, z + t(\vartheta_h(z)v - z)) - G(x, z)}{t} &= u^* \cdot (\vartheta_h(z)v - z) = \\ &= \frac{1}{1 - \bar{t}} [\vartheta_h(z)u^* \cdot v - u^* \cdot (z + \bar{t}(\vartheta_h(z)v - z))]. \end{aligned}$$

By relations (2.29) and (2.30), it follows that

$$\frac{|\vartheta_h(z)u^* \cdot v|}{1-\bar{t}} \leq 2 \|v\|_\infty \alpha_{hM}.$$

On the other hand, from  $(G_4)$  we deduce that for a.e.  $x \in \Omega$

$$\begin{aligned} \frac{u^* \cdot (z + \bar{t}(\vartheta_h(z)v - z))}{1-\bar{t}} &\geq -\frac{1}{1-\bar{t}} G^\circ(x, z + \bar{t}(\vartheta_h(z)v - z); -(z + \bar{t}(\vartheta_h(z)v - z))) \geq \\ &\geq -\frac{1}{1-\bar{t}} (a_1 + b_1 |z + \bar{t}(\vartheta_h(z)v - z)|^p) \geq -2(a_1 + b_1(|z| + |v|)^p). \end{aligned}$$

Then (2.33) easily follows.

For a.e.  $x \in \Omega$  we have

$$\begin{aligned} G^\circ(x, u; \vartheta_h(u)v - u) &\leq \vartheta_h(u)G^\circ(x, u; v - u) + (1 - \vartheta_h(u))G^\circ(x, u; -u) \leq \\ &\leq [G^\circ(x, u; v - u)]^+ + [G^\circ(x, u; -u)]^+. \end{aligned}$$

Furthermore, for a.e.  $x \in \Omega$  and every  $s \in \mathbb{R}^N$ ,  $(G_2)$  implies  $G^\circ(x, s; \cdot)$  to be Lipschitz continuous, so in particular

$$\lim_h G^\circ(x, u; \vartheta_h(u)v - u) = G^\circ(x, u; v - u) \quad \text{a.e. in } \Omega.$$

Then, given

$$\lambda > \int_\Omega G^\circ(x, u; v - u) dx,$$

by Fatou's lemma there exists  $\bar{h} \geq 1$  such that

$$\forall h \geq \bar{h} : \int_\Omega G^\circ(x, u; \vartheta_h(u)v - u) dx < \lambda \quad \text{and} \quad \|\vartheta_h(u)v - v\|_p < \varepsilon. \quad (2.34)$$

By the lower semicontinuity of  $\mathcal{G}$ , there exists  $\bar{\delta} \in ]0, 1/2]$  such that for every  $z \in B_{\bar{\delta}}(u)$  it is  $\mathcal{G}(z) \geq \mathcal{G}(u) - \frac{1}{2}$ . Then for every  $(z, \mu) \in B_{\bar{\delta}}(u, f(u)) \cap \text{epi}(f)$  it follows

$$\mathcal{E}(z) \leq \mu - \mathcal{G}(z) \leq \mu + \frac{1}{2} - \mathcal{G}(u) \leq f(u) + \bar{\delta} - \mathcal{G}(u) + \frac{1}{2} \leq \mathcal{E}(u) + 1.$$

Let now  $\sigma > 0$ . By assumptions  $(\mathcal{E}_1)$  and  $(\mathcal{E}_2.1)$  there exist  $h \geq \bar{h}$  and  $\delta \leq \bar{\delta}$  such that

$$\|v\|_\infty < hM,$$

$$\mathcal{E}(z) > \mathcal{E}(u) - \sigma, \quad \mathcal{E}(\vartheta_h(z)v) < \mathcal{E}(v) + \sigma, \quad \|(\vartheta_h(z)v - z) - (v - u)\|_p < \varepsilon,$$

for any  $z \in B_\delta(u)$  with  $\mathcal{E}(z) \leq \mathcal{E}(u) + 1$ .

Taking into account (D.1), (2.33) and (2.34), we deduce by Fatou's

lemma that, possibly reducing  $\delta$ , for any  $t \in ]0, \delta]$  and for any  $z \in B_\delta(u)$  we have

$$\int_{\Omega} \frac{G(x, z + t(\vartheta_h(z)v - z)) - G(x, z)}{t} dx < \lambda.$$

Now let  $\mathcal{V} : (B_\delta(u, f(u)) \cap \text{epi}(f)) \times ]0, \delta] \rightarrow B_\varepsilon(v - u)$  be defined setting

$$\mathcal{V}((z, \mu), t) = \vartheta_h(z)v - z.$$

Since  $\mathcal{V}$  is evidently continuous and

$$\begin{aligned} f(z + t\mathcal{V}((z, \mu), t)) &= f(z + t(\vartheta_h(z)v - z)) \leq \\ &\leq \mathcal{E}(z) + t(\mathcal{E}(\vartheta_h(z)v) - \mathcal{E}(z)) + \mathcal{G}(z + t(\vartheta_h(z)v - z)) \leq \\ &\leq \mathcal{E}(z) + (\mathcal{E}(v) - \mathcal{E}(u) + 2\sigma)t + \mathcal{G}(z) + \lambda t = \\ &= f(z) + (\mathcal{E}(v) - \mathcal{E}(u) + \lambda + 2\sigma)t, \end{aligned}$$

we have

$$f_\varepsilon^\circ(u; v - u) \leq \mathcal{E}(v) - \mathcal{E}(u) + \lambda + 2\sigma.$$

By the arbitrariness of  $\sigma > 0$  and  $\lambda > \int_{\Omega} G^\circ(x, u; v - u) dx$ , it follows

$$f_\varepsilon^\circ(u; v - u) \leq \mathcal{E}(v) - \mathcal{E}(u) + \int_{\Omega} G^\circ(x, u; v - u) dx.$$

Passing to the limit as  $\varepsilon \rightarrow 0^+$ , we get (2.31) when  $v \in \mathcal{D}(\mathcal{E}) \cap L^\infty(\Omega; \mathbb{R}^N)$ .

Let us now treat the general case. If we set  $v_h = \vartheta_h(v)v$ , we have  $v_h \in L^\infty(\Omega; \mathbb{R}^N)$ . Arguing as before, it is easy to see that

$$G^\circ(x, u; v_h - u) \leq [G^\circ(x, u; v - u)]^+ + [G^\circ(x, u; -u)]^+,$$

so that

$$\limsup_h \int_{\Omega} G^\circ(x, u; v_h - u) dx \leq \int_{\Omega} G^\circ(x, u; v - u) dx.$$

On the other hand, by the previous step it holds

$$f^\circ(u; v_h - u) \leq \mathcal{E}(v_h) - \mathcal{E}(u) + \int_{\Omega} G^\circ(x, u; v_h - u) dx.$$

Passing to the lower limit as  $h \rightarrow \infty$  and taking into account the lower semicontinuity of  $f^\circ(u, \cdot)$  and  $(\mathcal{E}_2.2)$ , we get (2.31).

(c) We already know that  $[G^\circ(x, u; -u)]^+ \in L^1(\Omega)$ . If we choose  $v = 0$  in (2.31), we obtain

$$f^\circ(u; -u) \leq \mathcal{E}(0) - \mathcal{E}(u) + \int_{\Omega} G^\circ(x, u; -u) dx.$$

Since  $\partial f(u) \neq \emptyset$ , it is  $f^\circ(u; -u) > -\infty$ , hence

$$\int_{\Omega} [G^\circ(x, u; -u)]^- dx < +\infty.$$

Finally, if  $u^* \in \partial f(u)$  we have by definition that

$$f^\circ(u; v - u) \geq \int_{\Omega} u^* \cdot (v - u) dx$$

and (2.32) follows from (2.31).

(d) From (2.28) it readily follows that  $G^\circ(x, u; v - u)$  is summable where  $|u(x)| \leq \|v\|_\infty$ . On the other hand, where  $|u(x)| > \|v\|_\infty$  we have

$$G^\circ(x, u; v - u) = \left(1 - \frac{v}{u}\right) G^\circ(x, u; -u)$$

and the assertion follows from  $(G_4)$ .  $\square$

Since  $f$  is only lower semicontinuous, we are interested in the verification of the condition  $(epi)_c$ . For this purpose, we consider an assumption  $(G'_3)$  on  $G$  stronger than  $(G_3)$ .

**Theorem 2.4** *Assume that*

$(G'_3)$  *there exist*  $a \in L^1(\Omega)$  *and*  $b \in \mathbb{R}$  *such that*

$$|G(x, s)| \leq a(x) + b|s|^p \quad \text{for a.e. } x \in \Omega \text{ and every } s \in \mathbb{R}^N.$$

*Then for every*  $(u, \lambda) \in \text{epi}(f)$  *with*  $\lambda > f(u)$  *it is*  $|d\mathcal{G}_f|(u, \lambda) = 1$ . *Moreover, if*  $\mathcal{E}$  *and*  $G(x, \cdot)$  *are even, for every*  $\lambda > f(0)$  *we have*  $|d_{\mathbb{Z}_2}\mathcal{G}_f|(0, \lambda) = 1$ .

*Proof* Let  $\varrho > 0$ . Since

$$\forall \tau \in [0, 1] : G^\circ(x, u; \tau u - u) = (1 - \tau)G^\circ(x, u; -u) \leq [G^\circ(x, u; -u)]^+,$$

by  $(\mathcal{E}_2.2)$  and  $(G_4)$  there exists  $\bar{h} \geq 1$  such that

$$\|\vartheta_{\bar{h}}(u)u - u\|_p < \varrho, \quad \mathcal{E}(\vartheta_{\bar{h}}(u)u) < \mathcal{E}(u) + \varrho,$$

$$\forall h \geq \bar{h} : \int_{\Omega} G^\circ(x, u; \vartheta_h(u)\vartheta_{\bar{h}}(u)u - u) dx < \varrho.$$

Set  $v = \vartheta_{\bar{h}}(u)u$ .

By  $(\mathcal{E}_2.1)$  there exist  $h \geq \bar{h}$  and  $\delta \in ]0, 1]$  such that

$$\|\vartheta_h(z)v - z\|_p < \varrho, \quad \mathcal{E}(\vartheta_h(z)v) < \mathcal{E}(u) + \varrho,$$

whenever  $\|z - u\|_p < \delta$  and  $\mathcal{E}(z) \leq \lambda + 1 - \mathcal{G}(u) + \varrho$ .

By decreasing  $\delta$ , from  $(G'_3)$ , (2.33) and (D.1) we deduce that

$$|\mathcal{G}(z) - \mathcal{G}(u)| < \varrho, \quad \int_{\Omega} \frac{G(x, z + t(\vartheta_h(z)v - z)) - G(x, z)}{t} dx < \varrho$$

whenever  $\|z - u\|_p < \delta$  and  $0 < t \leq \delta$ .

Define a continuous map

$$\mathcal{H} : \{z \in B_{\delta}(u) : f(z) < \lambda + \delta\} \times [0, \delta] \rightarrow X$$

by  $\mathcal{H}(z, t) = z + t(\vartheta_h(z)v - z)$ . It is readily seen that  $\|\mathcal{H}(z, t) - z\|_p \leq \varrho t$ .

If  $z \in B_{\delta}(u)$ ,  $f(z) < \lambda + \delta$  and  $0 \leq t \leq \delta$ , we have

$$\mathcal{E}(z) = f(z) - \mathcal{G}(z) < \lambda + \delta - \mathcal{G}(u) + \varrho \leq \lambda + 1 - \mathcal{G}(u) + \varrho,$$

hence, taking into account the convexity of  $\mathcal{E}$ ,

$$\mathcal{E}(z + t(\vartheta_h(z)v - z)) \leq \mathcal{E}(z) + t(\mathcal{E}(\vartheta_h(z)v) - \mathcal{E}(z)) \leq \mathcal{E}(z) + t(\mathcal{E}(u) - \mathcal{E}(z) + \varrho).$$

Moreover, we also have

$$\mathcal{G}(z + t(\vartheta_h(z)v - z)) \leq \mathcal{G}(z) + t\varrho \leq \mathcal{G}(z) + t(\mathcal{G}(u) - \mathcal{G}(z) + 2\varrho).$$

Therefore

$$f(z + t(\vartheta_h(z)v - z)) \leq f(z) + t(f(u) - f(z) + 3\varrho).$$

and the first assertion follows by Corollary 2.1.

Now assume that  $\mathcal{E}$  and  $G(x, \cdot)$  are even and that  $u = 0$ . Then, in the previous argument, we have  $v = 0$ , so that  $\mathcal{H}(-z, t) = -\mathcal{H}(z, t)$  and the second assertion also follows.  $\square$

Now we want to provide a criterion which helps in the verification of the Palais-Smale condition. For this purpose, we consider further assumptions on  $\mathcal{E}$ , which ensure a suitable coerciveness, and a new condition  $(G'_4)$  on  $G$ , stronger than  $(G_4)$ , which is a kind of one-sided subcritical growth condition.

**Theorem 2.5** *Let  $c \in \mathbb{R}$ . Assume that*

$(\mathcal{E}_3)$  *for every  $(u_h)$  bounded in  $L^p(\Omega; \mathbb{R}^N)$  with  $(\mathcal{E}(u_h))$  bounded, there exists a subsequence  $(u_{h_k})$  and a function  $u \in L^p(\Omega; \mathbb{R}^N)$  such that*

$$\lim_{k \rightarrow \infty} u_{h_k}(x) = u(x) \quad \text{for a.e. } x \in \Omega;$$

$(\mathcal{E}_4)$  *if  $(u_h)$  is a sequence in  $L^p(\Omega; \mathbb{R}^N)$  weakly convergent to  $u \in \mathcal{D}(\mathcal{E})$  and  $\mathcal{E}(u_h)$  converges to  $\mathcal{E}(u)$ , then  $(u_h)$  converges to  $u$  strongly in  $L^p(\Omega; \mathbb{R}^N)$ ;*

( $G'_4$ ) for every  $\varepsilon > 0$  there exists  $a_\varepsilon \in L^1(\Omega)$  such that

$$G^\circ(x, s; -s) \leq a_\varepsilon(x) + \varepsilon|s|^p \quad \text{for a.e. } x \in \Omega \text{ and every } s \in \mathbb{R}^N.$$

Then any  $(PS)_c$ -sequence  $(u_h)$  for  $f$  bounded in  $L^p(\Omega; \mathbb{R}^N)$  admits a subsequence strongly convergent in  $L^p(\Omega; \mathbb{R}^N)$ .

*Proof* From ( $G_3$ ) we deduce that  $(\mathcal{G}(u_h))$  is bounded from below. Taking into account ( $\mathcal{E}_1$ ), it follows that  $(\mathcal{E}(u_h))$  is bounded. By ( $\mathcal{E}_3$ ) there exists a subsequence, still denoted by  $(u_h)$ , converging weakly in  $L^p(\Omega; \mathbb{R}^N)$  and a.e. to some  $u \in \mathcal{D}(\mathcal{E})$ .

Given  $\varepsilon > 0$ , by ( $\mathcal{E}_2.2$ ) and ( $G_4$ ) we may find  $k_0 \geq 1$  such that

$$\mathcal{E}(\vartheta_{k_0}(u)u) < \mathcal{E}(u) + \varepsilon,$$

$$\int_{\Omega} (1 - \vartheta_{k_0}(u))G^\circ(x, u; -u) dx < \varepsilon.$$

Since  $\vartheta_{k_0}(u)u \in \mathcal{D}(\mathcal{E}) \cap L^\infty(\Omega; \mathbb{R}^N)$ , by ( $\mathcal{E}_2.1$ ) there exists  $k_1 \geq k_0$  such that

$$\forall h \in \mathbb{N} : \quad \mathcal{E}(\vartheta_{k_1}(u_h)\vartheta_{k_0}(u)u) < \mathcal{E}(u) + \varepsilon, \quad (2.35)$$

$$\int_{\Omega} (1 - \vartheta_{k_1}(u)\vartheta_{k_0}(u))G^\circ(x, u; -u) dx < \varepsilon.$$

It follows that  $\vartheta_{k_1}(u_h)\vartheta_{k_0}(u)u \in \mathcal{D}(\mathcal{E})$ . Moreover, from (2.28) and ( $G'_4$ ) we get

$$\begin{aligned} G^\circ(x, u_h; \vartheta_{k_1}(u_h)\vartheta_{k_0}(u)u - u_h) &\leq \\ &\leq \vartheta_{k_1}(u_h)G^\circ(x, u_h; \vartheta_{k_0}(u)u - u_h) + (1 - \vartheta_{k_1}(u_h))G^\circ(x, u_h; -u_h) \leq \\ &\leq \alpha_{k_1 M}(x)(k_0 M + k_1 M) + a_\varepsilon(x) + \varepsilon|u_h|^p. \end{aligned}$$

From (D.2) and Fatou's Lemma we deduce that

$$\begin{aligned} \limsup_{h \rightarrow \infty} \int_{\Omega} [G^\circ(x, u_h; \vartheta_{k_1}(u_h)\vartheta_{k_0}(u)u - u_h) - \varepsilon|u_h|^p] dx &\leq \\ &\leq \int_{\Omega} [G^\circ(x, u; \vartheta_{k_1}(u)\vartheta_{k_0}(u)u - u) - \varepsilon|u|^p] dx \leq \\ &\leq \int_{\Omega} (1 - \vartheta_{k_1}(u)\vartheta_{k_0}(u))G^\circ(x, u; -u) dx < \varepsilon, \end{aligned}$$

hence

$$\limsup_{h \rightarrow \infty} \int_{\Omega} G^\circ(x, u_h; \vartheta_{k_1}(u_h)\vartheta_{k_0}(u)u - u_h) dx < \varepsilon \sup_h \|u_h\|_p^p + \varepsilon. \quad (2.36)$$

Since  $(u_h)$  is a  $(PS)_c$ -sequence, by Theorem D.2 there exists  $u_h^* \in \partial f(u_h)$  with  $\|u_h^*\|_{p'} \leq |df|(u_h)$ , so that  $\lim_{h \rightarrow \infty} \|u_h^*\|_{p'} = 0$ . Applying (c) of Theorem 2.3, we get

$$\begin{aligned} \mathcal{E}(\vartheta_{k_1}(u_h)\vartheta_{k_0}(u)u) &\geq \mathcal{E}(u_h) - \int_{\Omega} G^\circ(x, u_h; \vartheta_{k_1}(u_h)\vartheta_{k_0}(u)u - u_h) dx + \\ &\quad + \int_{\Omega} u_h^* \cdot (\vartheta_{k_1}(u_h)\vartheta_{k_0}(u)u - u_h) dx. \end{aligned}$$

Taking into account (2.35), (2.36) and passing to the upper limit, we obtain

$$\limsup_{h \rightarrow \infty} \mathcal{E}(u_h) \leq \mathcal{E}(u) + 2\varepsilon + \varepsilon \sup_h \|u_h\|_p^p.$$

By the arbitrariness of  $\varepsilon > 0$ , we finally have

$$\limsup_{h \rightarrow \infty} \mathcal{E}(u_h) \leq \mathcal{E}(u)$$

and the strong convergence of  $(u_h)$  to  $u$  follows from  $(\mathcal{E}_4)$ .  $\square$

### 2.3.2 Area type functionals

Let  $n \geq 2$ ,  $N \geq 1$ ,  $\Omega$  be a bounded open subset of  $\mathbb{R}^n$  with Lipschitz boundary and let

$$\Psi : \mathbb{R}^{nN} \rightarrow \mathbb{R}$$

be a convex function satisfying

$$(\Psi) \quad \begin{cases} \Psi(0) = 0, \Psi(\xi) > 0 \text{ for any } \xi \neq 0 \text{ and} \\ \text{there exists } c > 0 \text{ such that } \Psi(\xi) \leq c|\xi| \text{ for any } \xi \in \mathbb{R}^{nN}. \end{cases}$$

We want to study the functional  $\mathcal{E} : L^{\frac{n}{n-1}}(\Omega; \mathbb{R}^N) \rightarrow \mathbb{R} \cup \{+\infty\}$  defined by

$$\mathcal{E}(u) = \begin{cases} \int_{\Omega} \Psi(Du^a) dx + \int_{\Omega} \Psi^\infty\left(\frac{Du^s}{|Du^s|}\right) d|Du^s|(x) + \\ \quad + \int_{\partial\Omega} \Psi^\infty(u \otimes \nu) d\mathcal{H}^{n-1}(x) & \text{if } u \in BV(\Omega; \mathbb{R}^N), \\ +\infty & \text{if } u \notin BV(\Omega; \mathbb{R}^N), \end{cases}$$

where  $Du = Du^a dx + Du^s$  is the Lebesgue decomposition of  $Du$ ,  $|Du^s|$  is the total variation of  $Du^s$ ,  $Du^s/|Du^s|$  is the Radon-Nikodym derivative

of  $Du^s$  with respect to  $|Du^s|$ ,  $\Psi^\infty$  is the recession functional associated with  $\Psi$ ,  $\nu$  is the outer normal to  $\Omega$  and the trace of  $u$  on  $\partial\Omega$  is still denoted by  $u$ .

**Theorem 2.6** *The functional  $\mathcal{E}$  satisfies conditions  $(\mathcal{E}_1)$ ,  $(\mathcal{E}_2)$ ,  $(\mathcal{E}_3)$  and  $(\mathcal{E}_4)$ .*

The section will be devoted to the proof of this result. We begin establishing some technical lemmas.

In  $BV(\Omega; \mathbb{R}^N)$  we consider the norm

$$\|u\|_{BV} = \int_{\Omega} |Du^a| dx + |Du^s|(\Omega) + \int_{\partial\Omega} |u| d\mathcal{H}^{n-1}(x),$$

which is equivalent to the standard norm of  $BV(\Omega; \mathbb{R}^N)$ .

**Lemma 2.2** *For every  $u \in BV(\Omega; \mathbb{R}^N)$  and every  $\varepsilon > 0$  there exists  $v \in C_c^\infty(\Omega; \mathbb{R}^N)$  such that  $\|v\|_\infty \leq \text{esssup}_\Omega |u|$  and*

$$\|v - u\|_{\frac{n}{n-1}} < \varepsilon, \quad \left| \int_{\Omega} |Dv| dx - \|u\|_{BV} \right| < \varepsilon, \quad |\mathcal{E}(v) - \mathcal{E}(u)| < \varepsilon.$$

*Proof* Let  $\delta > 0$ , let  $R > 0$  with  $\bar{\Omega} \subseteq B_R(0)$  and let

$$\vartheta_h(x) = 1 - \min \left\{ \max \left\{ \frac{h+1}{h} [1 - h d(x, \mathbb{R}^n \setminus \Omega)], 0 \right\}, 1 \right\}.$$

Define  $\hat{u} \in BV(B_R(0); \mathbb{R}^N)$  by

$$\hat{u}(x) = \begin{cases} u(x) & \text{if } x \in \Omega, \\ 0 & \text{if } x \in B_R(0) \setminus \Omega. \end{cases}$$

Then, if  $h$  is sufficiently large, we have  $\vartheta_h u \in BV(\Omega; \mathbb{R}^N)$ ,  $\|\vartheta_h u - u\|_{\frac{n}{n-1}} < \delta$  and

$$\begin{aligned} & \int_{\Omega} \sqrt{1 + |D(\vartheta_h u)^a|^2} d\mathcal{L}^n + |D(\vartheta_h u)^s|(\Omega) < \\ & < \int_{\Omega} \sqrt{1 + |Du^a|^2} d\mathcal{L}^n + |Du^s|(\Omega) + \int_{\partial\Omega} |u| d\mathcal{H}^{n-1} + \delta = \\ & = \int_{B_R(0)} \sqrt{1 + |D\hat{u}^a|^2} d\mathcal{L}^n + |D\hat{u}^s|(B_R(0)) + \delta. \end{aligned}$$

Moreover,  $\vartheta_h u$  has compact support in  $\Omega$  and  $\text{esssup}_\Omega |\vartheta_h u| \leq \text{esssup}_\Omega |u|$ .

If we regularize  $\vartheta_h u$  by convolution, we easily get  $v \in C_c^\infty(\Omega; \mathbb{R}^N)$  with

$$\|v\|_\infty \leq \text{esssup}_\Omega |u|, \quad \|v - u\|_{\frac{n}{n-1}} < \delta$$

and

$$\int_{\Omega} \sqrt{1 + |Dv|^2} d\mathcal{L}^n < \int_{B_R(0)} \sqrt{1 + |D\hat{u}^a|^2} d\mathcal{L}^n + |D\hat{u}^s|(B_R(0)) + \delta.$$

Since

$$\begin{aligned} \|u\|_{BV} &= \int_{B_R(0)} |D\hat{u}^a| dx + |D\hat{u}^s|(B_R(0)), \\ \mathcal{E}(u) &= \int_{B_R(0)} \Psi(D\hat{u}^a) dx + \int_{B_R(0)} \Psi^\infty \left( \frac{D\hat{u}^s}{|D\hat{u}^s|} \right) d|D\hat{u}^s|, \end{aligned}$$

our assertion follows.  $\square$

**Lemma 2.3** *The following facts hold:*

- (a)  $\Psi : \mathbb{R}^{nN} \rightarrow \mathbb{R}$  is Lipschitz continuous of some constant  $\text{Lip}(\Psi) > 0$ ;
- (b) for any  $\xi \in \mathbb{R}^{nN}$  and  $s \in [0, 1]$  we have  $\Psi(s\xi) \leq s\Psi(\xi)$ ;
- (c) for every  $\sigma > 0$  there exists  $d_\sigma > 0$  such that

$$\forall \xi \in \mathbb{R}^{nN} : \quad \Psi(\xi) \geq d_\sigma(|\xi| - \sigma);$$

- (d)  $\mathcal{E} : BV(\Omega; \mathbb{R}^N) \rightarrow \mathbb{R}$  is Lipschitz continuous of constant  $\text{Lip}(\Psi)$ ;
- (e) if  $\sigma$  and  $d_\sigma$  are as in (c), we have

$$\forall u \in BV(\Omega; \mathbb{R}^N) : \quad \mathcal{E}(u) \geq d_\sigma \left( \|u\|_{BV} - \sigma \mathcal{L}^n(\Omega) \right).$$

*Proof* Properties (a) and (b) easily follow from the convexity of  $\Psi$  and assumption  $(\Psi)$ .

To prove (c), assume by contradiction that  $\sigma > 0$  and  $(\xi_h)$  is a sequence with  $\Psi(\xi_h) < \frac{1}{h}(|\xi_h| - \sigma)$ . If  $|\xi_h| \rightarrow +\infty$ , we have eventually

$$\Psi \left( \frac{\xi_h}{|\xi_h|} \right) \leq \frac{\Psi(\xi_h)}{|\xi_h|} < \frac{1}{h} \left( 1 - \frac{\sigma}{|\xi_h|} \right).$$

Up to a subsequence,  $(\xi_h/|\xi_h|)$  is convergent to some  $\eta \neq 0$  with  $\Psi(\eta) \leq 0$ , which is impossible. Since  $|\xi_h|$  is bounded, up to a subsequence we have  $\xi_h \rightarrow \xi$  with  $|\xi| \geq \sigma$  and  $\Psi(\xi) \leq 0$ , which is again impossible.

Finally, (d) easily follows from (a) and the definition of  $\|\cdot\|_{BV}$ , while (e) follows from (c).  $\square$

Let now  $\vartheta \in C_c^1(\mathbb{R}^N)$  with  $0 \leq \vartheta \leq 1$ ,  $\|\nabla \vartheta\|_\infty \leq 2$ ,  $\vartheta(s) = 1$  for  $|s| \leq 1$  and  $\vartheta(s) = 0$  for  $|s| \geq 2$ . Define  $\vartheta_h : \mathbb{R}^N \rightarrow \mathbb{R}$  and  $T_h, R_h : \mathbb{R}^N \rightarrow \mathbb{R}^N$  by

$$\vartheta_h(s) = \vartheta \left( \frac{s}{h} \right), \quad T_h(s) = \vartheta_h(s)s, \quad R_h(s) = (1 - \vartheta_h(s))s.$$

**Lemma 2.4** *There exists a constant  $c_\Psi > 0$  such that*

$$\mathcal{E}\left(\vartheta\left(\frac{u}{h}\right)v\right) \leq \mathcal{E}(v) + \frac{c_\Psi}{h}\|v\|_\infty\|u\|_{BV},$$

$$\begin{aligned} \mathcal{E}(T_h \circ u) &\leq \mathcal{E}(u) + c_\Psi \left[ |Du|(\{x \in \Omega \setminus S_u : |\tilde{u}(x)| > h\}) + \right. \\ &\quad \left. + \int_{\{x \in S_u : |u^+(x)| > h \text{ or } |u^-(x)| > h\}} |u^+ - u^-| d\mathcal{H}^{n-1}(x) + \right. \\ &\quad \left. + \int_{\{x \in \partial\Omega : |u(x)| > h\}} |u| d\mathcal{H}^{n-1}(x) \right], \end{aligned}$$

$$\mathcal{E}(T_h \circ w) + \mathcal{E}(R_h \circ w) \leq \mathcal{E}(w) + c_\Psi \int_{\{x \in \Omega : h < |w(x)| < 2h\}} |Dw| dx$$

whenever  $h \geq 1$ ,  $u \in BV(\Omega; \mathbb{R}^N)$ ,  $v \in BV(\Omega; \mathbb{R}^N) \cap L^\infty(\Omega; \mathbb{R}^N)$  and  $w \in C_c^\infty(\Omega; \mathbb{R}^N)$ .

*Proof* Suppose first that  $u, v \in C_c^\infty(\Omega; \mathbb{R}^N)$ . Then, since

$$D\left[\vartheta\left(\frac{u}{h}\right)v\right] = \vartheta\left(\frac{u}{h}\right)Dv + \frac{1}{h}v \otimes \left[D\vartheta\left(\frac{u}{h}\right)Du\right],$$

by  $(\Psi)$  and Lemma 2.3 it follows that

$$\mathcal{E}\left(\vartheta\left(\frac{u}{h}\right)v\right) \leq \mathcal{E}(v) + \text{Lip}(\Psi) \frac{\|D\vartheta\|_\infty}{h} \|v\|_\infty \int_\Omega |Du| dx. \quad (2.37)$$

In the general case, let us consider two sequences  $(u_k), (v_k)$  in  $C_c^\infty(\Omega; \mathbb{R}^N)$  converging to  $u, v$  in  $L^1(\Omega; \mathbb{R}^N)$  with  $\int_\Omega |Du_k| dx \rightarrow \|u\|_{BV}$ ,  $\mathcal{E}(v_k) \rightarrow \mathcal{E}(v)$  and  $\|v_k\|_\infty \leq \|v\|_\infty$ . Passing to the lower limit in (2.37), we obtain the first inequality in the assertion.

To prove the second inequality, we first observe that by Lemma 2.3 we have

$$\mathcal{E}(T_h \circ u) \leq \mathcal{E}(u) + \text{Lip}(\Psi) \|R_h \circ u\|_{BV}. \quad (2.38)$$

In order to estimate the last term in (2.38), we apply the chain rule. Since  $R_h(s) = 0$  if  $|s| \leq h$  and  $\|DR_h\|_\infty \leq k_\vartheta$  for some  $k_\vartheta > 0$ , we have

$$\begin{aligned} \int_\Omega |D(R_h(u))^a| dx &\leq \int_{\Omega \setminus S_u} |DR_h(\tilde{u})| |Du^a| dx \leq \\ &k_\vartheta \int_{\{x \in \Omega \setminus S_u : |\tilde{u}(x)| > h\}} |Du^a| dx, \end{aligned}$$

$$\begin{aligned}
\left|D(R_h(u))^s\right|(\Omega) &\leq \int_{\Omega \setminus S_u} |DR_h(\tilde{u})| |dDu^s|(x) + \\
&\quad + \int_{S_u} |R_h(u^+) - R_h(u^-)| d\mathcal{H}^{n-1}(x) \leq \\
&\leq k_\vartheta \left( |Du^s|(\{x \in \Omega \setminus S_u : |\tilde{u}(x)| > h\}) + \right. \\
&\quad \left. + \int_{\{x \in S_u : |u^+(x)| > h \text{ or } |u^-(x)| > h\}} |u^+ - u^-| d\mathcal{H}^{m-1}(x) \right)
\end{aligned}$$

and

$$\int_{\partial\Omega} |R_h(u)| d\mathcal{H}^{n-1}(x) \leq k_\vartheta \int_{\{x \in \partial\Omega : |u(x)| > h\}} |u| d\mathcal{H}^{n-1}(x).$$

Combining these three estimates, we find

$$\begin{aligned}
\|R_h \circ u\|_{BV} &\leq k_\vartheta \left( \int_{\{x \in \Omega \setminus S_u : |\tilde{u}(x)| > h\}} |Du^s| dx \right. \\
&\quad + |Du^s|(\{x \in \Omega \setminus S_u : |\tilde{u}(x)| > h\}) \\
&\quad + \int_{\{x \in S_u : |u^+(x)| > h \text{ or } |u^-(x)| > h\}} |u^+ - u^-| d\mathcal{H}^{m-1}(x) \\
&\quad \left. + \int_{\{x \in \partial\Omega : |u(x)| > h\}} |u| d\mathcal{H}^{n-1}(x) \right). \tag{2.39}
\end{aligned}$$

Then the second inequality follows from (2.38) and (2.39).

Again, since  $\Psi$  is Lipschitz continuous, we have

$$\begin{aligned}
&\left| \int_{\Omega} \Psi(D(T_h \circ w)) dx - \int_{\Omega} \Psi(\vartheta_h(w)) Dw dx \right| \leq \\
&\frac{\text{Lip}(\Psi)}{h} \int_{\Omega} \left| D\vartheta\left(\frac{w}{h}\right) Dw \right| |w| dx \leq \\
&2 \text{Lip}(\Psi) \|\nabla\vartheta\|_{\infty} \int_{\{h < |w| < 2h\}} |Dw| dx.
\end{aligned}$$

In a similar way, we obtain

$$\begin{aligned}
&\left| \int_{\Omega} \Psi(D(R_h \circ w)) dx - \int_{\Omega} \Psi((1 - \vartheta_h(w))Dw) dx \right| \leq \\
&2 \text{Lip}(\Psi) \|\nabla\vartheta\|_{\infty} \int_{\{h < |w| < 2h\}} |Dw| dx.
\end{aligned}$$

Hence, combining the last two estimates and taking into account (b) of

Lemma 2.3, we get

$$\begin{aligned} \int_{\Omega} \Psi(D(T_h \circ w)) \, dx + \int_{\Omega} \Psi(D(R_h \circ w)) \, dx &\leq \\ &\leq \int_{\Omega} \Psi(Dw) \, dx + 4 \operatorname{Lip}(\Psi) \|\nabla \vartheta\|_{\infty} \int_{\{h < |w| < 2h\}} |Dw| \, dx \end{aligned}$$

and the proof is complete.  $\square$

**Lemma 2.5** *Let  $(u_h)$  be a sequence in  $C_c^{\infty}(\Omega; \mathbb{R}^N)$  and assume that  $(u_h)$  is bounded in  $BV(\Omega; \mathbb{R}^N)$ .*

*Then for every  $\varepsilon > 0$  and every  $\bar{k} \in \mathbb{N}$  there exists  $k \geq \bar{k}$  such that*

$$\liminf_{h \rightarrow \infty} \int_{\{k < |u_h| < 2k\}} |Du_h| \, dx < \varepsilon.$$

*Proof* Let  $m \geq 1$  be such that

$$\sup_h \int_{\Omega} |Du_h| \, dx \leq \frac{m\varepsilon}{2}$$

and let  $i_0 \in \mathbb{N}$  with  $2^{i_0} \geq \bar{k}$ . Then, since

$$\sum_{i=i_0}^{i_0+m-1} \int_{\{2^i < |u_h| < 2^{i+1}\}} |Du_h| \, dx \leq \int_{\Omega} |Du_h| \, dx \leq \frac{m\varepsilon}{2},$$

there exists  $i_h$  between  $i_0$  and  $i_0 + m - 1$  such that

$$\int_{\{2^{i_h} < |u_h| < 2^{i_h+1}\}} |Du_h| \, dx \leq \frac{\varepsilon}{2}.$$

Passing to a subsequence  $(i_{h_j})$ , we can suppose  $i_{h_j} \equiv i \geq i_0$ , and setting  $k = 2^i$  we get

$$\forall j \in \mathbb{N} : \int_{\{k < |u_{h_j}| < 2k\}} |Du_{h_j}| \, dx \leq \frac{\varepsilon}{2}.$$

Then the assertion follows.  $\square$

**Lemma 2.6** *Let  $(u_h)$  be a sequence in  $C_c^{\infty}(\Omega; \mathbb{R}^N)$  and let  $u \in BV(\Omega; \mathbb{R}^N)$  with  $\|u_h - u\|_1 \rightarrow 0$  and  $\mathcal{E}(u_h) \rightarrow \mathcal{E}(u)$ .*

*Then for every  $\varepsilon > 0$  and every  $\bar{k} \in \mathbb{N}$  there exists  $k \geq \bar{k}$  such that*

$$\liminf_{h \rightarrow \infty} \|R_k \circ u_h\|_{BV} < \varepsilon.$$

*Proof* Given  $\varepsilon > 0$ , let  $d > 0$  be such that

$$\forall \xi \in \mathbb{R}^{nN} : \quad \Psi(\xi) \geq d \left( |\xi| - \frac{\varepsilon}{3\mathcal{L}^n(\Omega)} \right),$$

according to Lemma 2.3. Let also  $c_\Psi > 0$  be as in Lemma 2.4. By (2.39) and Lemma 2.5, there exists  $k \geq \bar{k}$  such that

$$\|R_k \circ u\|_{BV} < \frac{d\varepsilon}{3\text{Lip}(\Psi)},$$

$$\liminf_{h \rightarrow \infty} \int_{\{k < |u_h| < 2k\}} |Du_h| dx < \frac{d\varepsilon}{3c_\Psi}.$$

From Lemma 2.4 we deduce that

$$\begin{aligned} & \mathcal{E}(T_k \circ u) + \liminf_{h \rightarrow \infty} \mathcal{E}(R_k \circ u_h) \leq \\ & \liminf_{h \rightarrow \infty} \mathcal{E}(T_k \circ u_h) + \liminf_{h \rightarrow \infty} \mathcal{E}(R_k \circ u_h) \leq \\ & \liminf_{h \rightarrow \infty} \left( \mathcal{E}(T_k \circ u_h) + \mathcal{E}(R_k \circ u_h) \right) \leq \\ & \mathcal{E}(u) + c_\Psi \liminf_{h \rightarrow \infty} \int_{\{k < |u_h| < 2k\}} |Du_h| dx < \\ & \mathcal{E}(u) + \frac{d\varepsilon}{3} \leq \mathcal{E}(T_k \circ u) + \text{Lip}(\Psi) \|R_k \circ u\|_{BV} + \frac{d\varepsilon}{3} < \\ & \mathcal{E}(T_k \circ u) + \frac{2}{3}d\varepsilon, \end{aligned}$$

whence

$$\liminf_{h \rightarrow \infty} \mathcal{E}(R_k \circ u_h) < \frac{2}{3}d\varepsilon.$$

On the other hand, by Lemma 2.3 we have

$$\mathcal{E}(R_k \circ u_h) \geq d \left( \|R_k \circ u_h\|_{BV} - \frac{\varepsilon}{3} \right)$$

and the assertion follows.  $\square$

Now we can prove the main auxiliary result we need for the proof of Theorem 2.6. It is a property of the space  $BV$  which could be interesting also in itself.

**Theorem 2.7** *Let  $(u_h)$  be a sequence in  $BV(\Omega; \mathbb{R}^N)$ . Assume  $u \in BV(\Omega; \mathbb{R}^N)$  such that  $\|u_h - u\|_1 \rightarrow 0$  and  $\mathcal{E}(u_h) \rightarrow \mathcal{E}(u)$ .*

*Then  $(u_h)$  is strongly convergent to  $u$  in  $L^{\frac{n}{n-1}}(\Omega; \mathbb{R}^N)$ .*

*Proof* By Lemma 2.2 we may find  $v_h \in C_c^\infty(\Omega; \mathbb{R}^N)$  with

$$\|v_h - u_h\|_1 < \frac{1}{h}, \quad \|v_h - u_h\|_{\frac{n}{n-1}} < \frac{1}{h}, \quad |\mathcal{E}(v_h) - \mathcal{E}(u_h)| < \frac{1}{h}.$$

Therefore it is sufficient to treat the case in which  $u_h \in C_c^\infty(\Omega; \mathbb{R}^N)$ .

By contradiction, up to a subsequence we may assume that there exists  $\varepsilon > 0$  such that  $\|u_h - u\|_{\frac{n}{n-1}} \geq \varepsilon$ . Let  $\tilde{c}$  be a constant such that  $\|w\|_{\frac{n}{n-1}} \leq \tilde{c}\|w\|_{BV}$  for any  $w \in BV(\Omega; \mathbb{R}^N)$ . According to Lemma 2.6, let  $k \in \mathbb{N}$  be such that

$$\|R_k \circ u\|_{\frac{n}{n-1}} < \frac{\varepsilon}{2}, \quad \liminf_{h \rightarrow \infty} \|R_k \circ u_h\|_{\frac{n}{n-1}} \leq \tilde{c} \liminf_{h \rightarrow \infty} \|R_k \circ u_h\|_{BV} < \frac{\varepsilon}{2}.$$

Then we have

$$\|u_h - u\|_{\frac{n}{n-1}} \leq \|R_k \circ u_h\|_{\frac{n}{n-1}} + \|T_k \circ u_h - T_k \circ u\|_{\frac{n}{n-1}} + \|R_k \circ u\|_{\frac{n}{n-1}}. \quad (2.40)$$

Since  $T_k \circ u_h \rightarrow T_k \circ u$  in  $L^{\frac{n}{n-1}}(\Omega; \mathbb{R}^N)$  as  $h \rightarrow \infty$ , passing to the lower limit in (2.40) we get

$$\liminf_{h \rightarrow \infty} \|u_h - u\|_{\frac{n}{n-1}} < \varepsilon,$$

whence a contradiction.  $\square$

*Proof of Theorem 2.6.* It is well known that  $\mathcal{E}$  satisfies condition  $(\mathcal{E}_1)$ . Conditions  $(\mathcal{E}_2)$  are an immediate consequence of Lemma 2.4. From (e) of Lemma 2.3 and Rellich's theorem, it follows that  $\mathcal{E}$  satisfies condition  $(\mathcal{E}_3)$ . To prove  $(\mathcal{E}_4)$ , let  $(u_h)$  be a sequence in  $L^{\frac{n}{n-1}}(\Omega; \mathbb{R}^N)$  weakly convergent to  $u \in BV(\Omega; \mathbb{R}^N)$  such that  $\mathcal{E}(u_h)$  converges to  $\mathcal{E}(u)$ . Again by (e) of Lemma 2.3 and Rellich's theorem we deduce that  $(u_h)$  is strongly convergent to  $u$  in  $L^1(\Omega; \mathbb{R}^N)$ . Then the assertion follows from Theorem 2.7.

### 2.3.3 A result of Clark type

Let  $n \geq 2$  and  $\Omega$  be a bounded open subset of  $\mathbb{R}^n$  with Lipschitz boundary, let  $\Psi : \mathbb{R}^{nN} \rightarrow \mathbb{R}$  be an even convex function satisfying  $(\Psi)$  and let  $G : \Omega \times \mathbb{R}^N \rightarrow \mathbb{R}$  be a function satisfying  $(G_1)$ ,  $(G_2)$ ,  $(G'_3)$ ,  $(G'_4)$  with  $p = n/(n-1)$  and the following conditions:

$$\begin{cases} \text{there exist } \tilde{a} \in L^1(\Omega) \text{ and } \tilde{b} \in L^n(\Omega) \text{ such that} \\ G(x, s) \geq -\tilde{a}(x) - \tilde{b}(x)|s| \quad \text{for a.e. } x \in \Omega \text{ and every } s \in \mathbb{R}^N; \end{cases} \quad (2.41)$$

$$\lim_{|s| \rightarrow \infty} \frac{G(x, s)}{|s|} = +\infty \quad \text{for a.e. } x \in \Omega; \quad (2.42)$$

$$\{s \mapsto G(x, s)\} \text{ is even for a.e. } x \in \Omega. \quad (2.43)$$

Finally, define  $\mathcal{E}$  as in Section 4. The main result of this section is:

**Theorem 2.8** *For every  $k \in \mathbb{N}$  there exists  $\Lambda_k$  such that for any  $\lambda \geq \Lambda_k$  the problem*

$$\begin{cases} u \in BV(\Omega; \mathbb{R}^N) \\ \mathcal{E}(v) - \mathcal{E}(u) + \int_{\Omega} G^{\circ}(x, u; v-u) dx \geq \lambda \int_{\Omega} \frac{u}{\sqrt{1+|u|^2}} \cdot (v-u) dx \quad \forall v \in BV(\Omega; \mathbb{R}^N) \end{cases}$$

*admits at least  $k$  pairs  $(u, -u)$  of distinct solutions.*

For the proof we need the following auxiliary result.

**Lemma 2.7** *Let  $(u_h)$  be a bounded sequence in  $L^{\frac{n}{n-1}}(\Omega; \mathbb{R}^N)$ , which is convergent a.e. to  $u$ , and let  $(\varrho_h)$  be a positively divergent sequence of real numbers.*

*Then we have*

$$\begin{aligned} \lim_h \int_{\Omega} \frac{G(x, \varrho_h u_h)}{\varrho_h} dx &= +\infty \quad \text{if } u \neq 0, \\ \liminf_h \int_{\Omega} \frac{G(x, \varrho_h u_h)}{\varrho_h} dx &\geq 0 \quad \text{if } u = 0. \end{aligned}$$

*Proof* If  $u = 0$ , the assertion follows directly from (2.41). If  $u \neq 0$ , we have

$$\int_{\Omega} \frac{G(x, \varrho_h u_h)}{\varrho_h} dx \geq \int_{\{u \neq 0\}} \frac{G(x, \varrho_h u_h)}{\varrho_h} dx - \frac{1}{\varrho_h} \int_{\{u=0\}} \tilde{a} dx - \int_{\{u=0\}} \tilde{b}|u_h| dx.$$

From (2.41), (2.42) and Fatou's Lemma, we deduce that

$$\lim_h \int_{\{u \neq 0\}} \frac{G(x, \varrho_h u_h)}{\varrho_h} dx = +\infty,$$

whence the assertion.  $\square$

*Proof of Theorem 2.8.* First of all, set

$$\tilde{G}(x, s) = G(x, s) - \lambda(\sqrt{1+|s|^2} - 1).$$

It is easy to see that also  $\tilde{G}$  satisfies  $(G_1)$ ,  $(G_2)$ ,  $(G'_3)$ ,  $(G'_4)$ , (2.41), (2.42), (2.43) and that

$$\tilde{G}^{\circ}(x, s; \hat{s}) = G^{\circ}(x, s; \hat{s}) - \lambda \frac{s}{\sqrt{1+|s|^2}} \cdot \hat{s}.$$

Now define a lower semicontinuous functional  $f : L^{\frac{n}{n-1}}(\Omega; \mathbb{R}^N) \rightarrow \mathbb{R} \cup \{+\infty\}$  by

$$f(u) = \mathcal{E}(u) + \int_{\Omega} \tilde{G}(x, u) dx.$$

Then  $f$  is even by (2.43) and satisfies condition  $(epi)_c$  by Theorem 2.4. We claim that

$$\lim_{\|u\|_{\frac{n}{n-1}} \rightarrow \infty} f(u) = +\infty. \quad (2.44)$$

To prove it, let  $(u_h)$  be a sequence in  $BV(\Omega; \mathbb{R}^N)$  with  $\|u_h\|_{\frac{n}{n-1}} = 1$  and let  $\varrho_h \rightarrow +\infty$ . By (e) of Lemma 2.3 there exist  $\tilde{c} > 0$  and  $\tilde{d} > 0$  such that

$$\forall u \in BV(\Omega; \mathbb{R}^N) : \quad \mathcal{E}(u) \geq \tilde{d} \left( \|u\|_{BV} - \tilde{c} \mathcal{L}^n(\Omega) \right).$$

If  $\|u_h\|_{BV} \rightarrow +\infty$ , it readily follows from (2.41) that  $f(\varrho_h u_h) \rightarrow +\infty$ . Otherwise, up to a subsequence,  $u_h$  is convergent a.e. and the assertion follows from the previous Lemma and the inequality

$$f(\varrho_h u_h) \geq \varrho_h \left[ \tilde{d} \left( \|u_h\|_{BV} - \frac{\tilde{c}}{\varrho_h} \mathcal{L}^n(\Omega) \right) + \int_{\Omega} \frac{\tilde{G}(x, \varrho_h u_h)}{\varrho_h} dx \right].$$

Since  $f$  is bounded below on bounded subsets of  $L^{\frac{n}{n-1}}(\Omega; \mathbb{R}^N)$ , it follows from (2.44) that  $f$  is bounded below on all  $L^{\frac{n}{n-1}}(\Omega; \mathbb{R}^N)$ ; furthermore, it also turns out from (2.44) that any  $(PS)_c$  sequence is bounded, hence  $f$  satisfies  $(PS)_c$  by Theorem 2.5.

Finally, let  $k \geq 1$ , let  $w_1, \dots, w_k$  be linearly independent elements of  $BV(\Omega; \mathbb{R}^N)$  and let  $\psi : S^{k-1} \rightarrow L^{\frac{n}{n-1}}(\Omega; \mathbb{R}^N)$  be the odd continuous map defined by

$$\psi(\xi) = \sum_{j=1}^k \xi_j w_j.$$

Because of  $(G'_3)$ , it is easily seen that

$$\sup \left\{ \mathcal{E}(u) + \int_{\Omega} G(x, u) dx : u \in \psi(S^{k-1}) \right\} < +\infty$$

and

$$\inf \left\{ \int_{\Omega} (\sqrt{1 + |u|^2} - 1) dx : u \in \psi(S^{k-1}) \right\} > 0.$$

Therefore there exists  $\Lambda_k > 0$  such that  $\sup_{\xi \in S^{k-1}} f(\psi(\xi)) < 0$  whenever  $\lambda \geq \Lambda_k$ .

Next, we recall the following result which is due to Clark. We refer to Theorem 2.5 in [86] for a nonsmooth version and complete proof.

**Theorem 2.9** *Let  $X$  be a Banach space and  $f : X \rightarrow \mathbb{R} \cup \{+\infty\}$  an even lower semicontinuous function. Assume that*

- (a)  $f$  is bounded from below;
- (b) for every  $c < f(0)$ , the function  $f$  satisfies  $(PS)_c$  and  $(epi)_c$ ;
- (c) there exist  $k \geq 1$  and an odd continuous map  $\psi : S^{k-1} \rightarrow X$  such that

$$\sup \{f(\psi(x)) : x \in S^{k-1}\} < f(0),$$

where  $S^{k-1}$  denotes the unit sphere in  $\mathbb{R}^k$ .

Then  $f$  admits at least  $k$  pairs  $(u_1, -u_1), \dots, (u_k, -u_k)$  of critical points with  $f(u_j) < f(0)$ .

Applying now Theorem 2.9, it follows that  $f$  admits at least  $k$  pairs  $(u_k, -u_k)$  of critical points. Therefore, by Theorem D.2, for any  $u_k$  it is possible to apply Theorem 2.3 (with  $\tilde{G}$  instead of  $G$ ), which completes the proof.

### 2.3.4 A superlinear potential

Let  $n \geq 2$  and  $\Omega$  be a bounded open subset of  $\mathbb{R}^n$  with Lipschitz boundary, let  $\Psi : \mathbb{R}^{nN} \rightarrow \mathbb{R}$  be an even convex function satisfying  $(\Psi)$  and let  $G : \Omega \times \mathbb{R}^N \rightarrow \mathbb{R}$  be a function satisfying  $(G_1)$ ,  $(G_2)$ ,  $(G'_3)$ ,  $(G'_4)$ , (2.43) with  $p = \frac{n}{n-1}$  and the following condition:

$$\left\{ \begin{array}{l} \text{there exist } q > 1 \text{ and } R > 0 \text{ such that} \\ G^\circ(x, s; s) \leq qG(x, s) < 0 \quad \text{for a.e. } x \in \Omega \text{ and every } s \in \mathbb{R}^N \text{ with } |s| \geq R. \end{array} \right. \quad (2.45)$$

Define  $\mathcal{E}$  as in section 4 and an even lower semicontinuous functional  $f : L^{\frac{n}{n-1}}(\Omega; \mathbb{R}^N) \rightarrow \mathbb{R} \cup \{+\infty\}$  by

$$f(u) = \mathcal{E}(u) + \int_{\Omega} G(x, u) dx.$$

**Theorem 2.10** *There exists a sequence  $(u_h)$  of solutions of the problem*

$$\left\{ \begin{array}{l} u \in BV(\Omega; \mathbb{R}^N) \\ \mathcal{E}(v) - \mathcal{E}(u) + \int_{\Omega} G^\circ(x, u; v - u) dx \geq 0 \quad \forall v \in BV(\Omega; \mathbb{R}^N) \end{array} \right.$$

with  $f(u_h) \rightarrow +\infty$ .

*Proof* According to (2.28), we have

$$|s| < R \implies |G^\circ(x, s; s)| \leq \alpha_R(x)|s|.$$

Combining this fact with (2.45) and  $(G'_3)$ , we deduce that there exists  $a_0 \in L^1(\Omega)$  such that

$$G^\circ(x, s; s) \leq qG(x, s) + a_0(x) \quad \text{for a.e. } x \in \Omega \text{ and every } s \in \mathbb{R}^N. \quad (2.46)$$

Moreover, from (2.45) and Lebourg's mean value theorem it follows that for every  $s \in \mathbb{R}^N$  with  $|s| = 1$  the function  $\{t \rightarrow t^{-q}G(x, ts)\}$  is nonincreasing on  $[R, +\infty[$ . Taking into account  $(G'_3)$  and possibly substituting  $a_0$  with another function in  $L^1(\Omega)$ , we deduce that

$$G(x, s) \leq a_0(x) - b_0(x)|s|^q \quad \text{for a.e. } x \in \Omega \text{ and every } s \in \mathbb{R}^N, \quad (2.47)$$

where

$$b_0(x) = \inf_{|s|=1} (-R^{-q}G(x, Rs)) > 0 \quad \text{for a.e. } x \in \Omega.$$

Finally, since  $\{\hat{s} \rightarrow G^\circ(x, s; \hat{s})\}$  is a convex function vanishing at the origin, we have  $G^\circ(x, s; s) \geq -G^\circ(x, s; -s)$ . Combining (2.46) with  $(G'_4)$ , we deduce that for every  $\varepsilon > 0$  there exists  $\tilde{a}_\varepsilon \in L^1(\Omega)$  such that

$$G(x, s) \geq -\tilde{a}_\varepsilon(x) - \varepsilon|s|^{\frac{n}{n-1}} \quad \text{for a.e. } x \in \Omega \text{ and every } s \in \mathbb{R}^N. \quad (2.48)$$

By Theorem 2.4 we have that  $f$  satisfies  $(epi)_c$  for any  $c \in \mathbb{R}$  and that  $|d_{\mathbb{Z}_2} \mathcal{G}_f|(0, \lambda) = 1$  for any  $\lambda > f(0)$ .

We also recall that, since  $\Psi$  is Lipschitz continuous, there exists  $M \in \mathbb{R}$  such that

$$(q+1)\Psi(\xi) - \Psi(2\xi) \geq \frac{q-1}{2}\Psi(\xi) - M, \quad (2.49)$$

$$(q+1)\Psi^\infty(\xi) - \Psi^\infty(2\xi) \geq \frac{q-1}{2}\Psi^\infty(\xi). \quad (2.50)$$

We claim that  $f$  satisfies the condition  $(PS)_c$  for every  $c \in \mathbb{R}$ . Let  $(u_h)$  be a  $(PS)_c$ -sequence for  $f$ . By Theorem D.2 there exists a sequence  $(u_h^*)$  in  $L^n(\Omega; \mathbb{R}^N)$  with  $u_h^* \in \partial f(u_h)$  and  $\|u_h^*\|_n \rightarrow 0$ . According to Theorem 2.3 and (2.46), we have

$$\begin{aligned} \mathcal{E}(2u_h) &\geq \mathcal{E}(u_h) - \int_{\Omega} G^\circ(x, u_h; u_h) dx + \int_{\Omega} u_h^* \cdot u_h dx \geq \\ &\geq \mathcal{E}(u_h) - q \int_{\Omega} G(x, u_h) dx + \int_{\Omega} u_h^* \cdot u_h dx - \int_{\Omega} a_0(x) dx. \end{aligned}$$

By the definition of  $f$ , it follows

$$qf(u_h) + \|u_h^*\|_n \|u_h\|_{\frac{n}{n-1}} + \int_{\Omega} a_0(x) dx \geq (q+1)\mathcal{E}(u_h) - \mathcal{E}(2u_h).$$

Finally, applying (2.49) and (2.50) we get

$$qf(u_h) + \|u_h^*\|_n \|u_h\|_{\frac{n}{n-1}} + \int_{\Omega} a_0(x) dx \geq \frac{q-1}{2}\mathcal{E}(u_h) - M\mathcal{L}^n(\Omega).$$

By (e) of Lemma 2.3 we deduce that  $(u_h)$  is bounded in  $BV(\Omega; \mathbb{R}^N)$ , hence in  $L^{\frac{n}{n-1}}(\Omega; \mathbb{R}^N)$ . Applying Theorem 2.5 we get that  $(u_h)$  admits a strongly convergent subsequence and  $(PS)_c$  follows.

Next, there exist a strictly increasing sequence  $(W_h)$  of finite-dimensional subspaces of  $BV(\Omega; \mathbb{R}^N) \cap L^\infty(\Omega; \mathbb{R}^N)$  and a strictly decreasing sequence  $(Z_h)$  of closed subspaces of  $L^{\frac{n}{n-1}}(\Omega; \mathbb{R}^N)$  such that  $L^{\frac{n}{n-1}}(\Omega; \mathbb{R}^N) = W_h \oplus Z_h$  and  $\bigcap_{h=0}^{\infty} Z_h = \{0\}$ . By (e) of Lemma 2.3 there exists  $\varrho > 0$  such that

$$\forall u \in L^{\frac{n}{n-1}}(\Omega; \mathbb{R}^N) : \quad \|u\|_{\frac{n}{n-1}} = \varrho \quad \implies \quad \mathcal{E}(u) \geq 1.$$

We claim that

$$\lim_h \left( \inf \{ f(u) : u \in Z_h, \|u\|_{\frac{n}{n-1}} = \varrho \} \right) > f(0).$$

Actually, assume by contradiction that  $(u_h)$  is a sequence with  $u_h \in Z_h$ ,  $\|u_h\|_{\frac{n}{n-1}} = \varrho$  and

$$\limsup_h f(u_h) \leq f(0).$$

Taking into account  $(G'_3)$  and Lemma 2.3, we deduce that  $(\mathcal{E}(u_h))$  is bounded, so that  $(u_h)$  is bounded in  $BV(\Omega; \mathbb{R}^N)$ . Therefore, up to a subsequence,  $(u_h)$  is convergent a.e. to 0. From (2.48) it follows that

$$\liminf_h \int_{\Omega} \left( G(x, u_h) + \varepsilon |u_h|^{\frac{n}{n-1}} \right) dx \geq \int_{\Omega} G(x, 0) dx,$$

hence

$$\liminf_h \int_{\Omega} G(x, u_h) dx \geq \int_{\Omega} G(x, 0) dx$$

by the boundedness of  $(u_h)$  in  $L^{\frac{n}{n-1}}(\Omega; \mathbb{R}^N)$  and the arbitrariness of  $\varepsilon$ . Therefore

$$\limsup_h \mathcal{E}(u_h) \leq \mathcal{E}(0) = 0$$

which contradicts the choice of  $\varrho$ .

Now, fix  $\bar{h}$  with

$$\inf\{f(u) : u \in Z_{\bar{h}}, \|u\|_{\frac{n}{n-1}} = \varrho\} > f(0)$$

and set  $Z = Z_{\bar{h}}$  and  $V_h = W_{\bar{h}+h}$ . Then  $Z$  satisfies assumption (a) of Theorem D.3 for some  $\alpha > f(0)$ .

Finally, since  $V_h$  is finite-dimensional,

$$\|u\|_G := \left( \int_{\Omega} b_0 |u|^q dx \right)^{\frac{1}{q}}$$

is a norm on  $V_h$  equivalent to the norm of  $BV(\Omega; \mathbb{R}^N)$ . Then, combining (2.47) with (d) of Lemma 2.3, we see that also assumption (b) of Theorem D.3 is satisfied.

Therefore there exists a sequence  $(u_h)$  of critical points for  $f$  with  $f(u_h) \rightarrow +\infty$  and, by Theorems D.2 and 2.3, the result follows.  $\square$

## 2.4 Historical notes and comments

We start with the following elementary proof due to H. Brézis of the Hartman–Stampacchia theorem. We first observe that since  $V$  is a finite dimensional space (say,  $\dim V = n$ ), then we can identify  $V^*$  with  $\mathbb{R}^n$  through the mapping  $\pi : V^* \rightarrow \mathbb{R}^n$  defined by

$$\langle a, x \rangle = (\pi a, x) \quad \text{for all } a \in V^*, x \in \mathbb{R}^n,$$

where  $(\cdot, \cdot)$  stands for the scalar product in  $\mathbb{R}^n$ . This enables us to observe that for proving relation (2.1), it is enough to show that there is some  $u \in K$  such that for every  $v \in K$ ,

$$(u, v - u) \geq (u - \pi A u, v - u). \quad (2.51)$$

But the mapping  $P_K(I - \pi A) : K \rightarrow K$  is continuous, where  $I : K \rightarrow K$  is the identity map and  $P_K : V \rightarrow K$  denotes the projection map onto the compact and convex set  $K$ . So, by the Brouwer fixed point theorem, there exists  $u \in K$  such that  $u = P_K(I - \pi A)u$ . Next, using the characterization of the projection on a closed convex subset of a Hilbert space (see [47, Théorème V.2]), we deduce (2.51).

The space of functions of bounded variation is very useful in the calculus of variations (geometric measure theory, fracture mechanics and image processing), as well as in the study of shock waves for nonlinear hyperbolic conservation laws. The inequality problem we have analyzed

in Section 2.3 is related to the problem of finding *minimal surfaces*, which was intensively studied in the last decades: *find the surface of least area among those bounded by a given curve*. This is also called the *Plateau problem*, after the physicist who did numerous experiments with soap films and bubbles. The standard nonparametric minimal surface equation is

$$\operatorname{div} \left( \frac{\nabla u}{\sqrt{1 + |\nabla u|^2}} \right) = 0, \quad u : \Omega \subset \mathbb{R}^2 \rightarrow \mathbb{R}.$$

This equation was first derived in 1762 by Lagrange, as an illustration of the Euler–Lagrange variational principle. We also point out that, up to a multiplicative factor, the expression  $\operatorname{div} \left( \nabla u / \sqrt{1 + |\nabla u|^2} \right)$  gives the mean curvature of the hyper-surface in  $\mathbb{R}^3$  defined by the graph of the function  $u$ .

We recall in what follows some basic definitions and properties in the theory of functions of bounded variation. We refer to the excellent monograph by Ziemer [305] for more details and complete proofs.

Let  $\Omega$  be an open set in  $\mathbb{R}^n$ ,  $n \geq 2$ . Recall first that if  $N \geq 1$  then  $u$  belongs to the space of functions of bounded variation  $BV(\Omega; \mathbb{R}^N)$  if the distributional derivative of  $u$  is representable by a finite Radon measure in  $\Omega$ . Next, the total variation of  $u$  is given by

$$\int_{\Omega} |Du| = \sup \left\{ \int_{\Omega} u \operatorname{div} \varphi \, dx; \varphi \in C_0^1(\Omega; \mathbb{R}^{nN}), |\varphi(x)| \leq 1 \text{ on } \Omega \right\}.$$

An useful result asserts that a  $L^1$  function belongs to  $BV(\Omega; \mathbb{R}^N)$  if and only if the total variation of  $u$  is finite.

We illustrate the notion of total variation of a function with the following elementary example. Assume that  $u \in C^1(\Omega; \mathbb{R})$ . Integrating by parts we obtain, for all  $\varphi \in C_0^1(\Omega; \mathbb{R}^n)$ ,

$$\int_{\Omega} u \operatorname{div} \varphi \, dx = - \int_{\Omega} \sum_{j=1}^n \frac{\partial u}{\partial x_j} \varphi_j \, dx.$$

Therefore

$$\int_{\Omega} |Du| = \int_{\Omega} |\operatorname{grad} u| \, dx, \quad \text{where } \operatorname{grad} u = \left( \frac{\partial u}{\partial x_1}, \dots, \frac{\partial u}{\partial x_n} \right). \quad (2.52)$$

More generally, if  $u \in W^{1,1}(\Omega)$ , then relation (2.52) still remains true, where now  $\operatorname{grad} u = (u_1, \dots, u_n)$  and  $u_1, \dots, u_n$  are the generalized derivatives of  $u$ .

To consider functionals defined on measures, let  $\Psi : \mathbb{R}^{nN} \rightarrow \mathbb{R}$  be convex, continuous, of linear growth and satisfying  $\Psi(0) = 0$ . Then the recession functional  $\Psi^\infty : \mathbb{R}^{nN} \rightarrow \mathbb{R}$  can be characterized as the support function of the convex function conjugate of  $\Psi$  and is defined by

$$\Psi^\infty(Z) = \limsup_{t \rightarrow +\infty} \frac{\Psi(tZ)}{t} = \lim_{t \rightarrow +\infty} \frac{\Psi(tZ)}{t} \quad \text{for all } Z \in \mathbb{R}^{nN}.$$

Then  $\Psi^\infty$  is positively homogeneous of degree one. The definition of the recession functional (also called *asymptotic functional*) shows that, in some sense,  $\Psi^\infty$  characterizes the growth speed at infinity of  $\Psi$  in a similar way as the recession cone of a convex set  $C$  characterizes the behaviour of  $C$  at infinity.

### 3

## Nonlinear Eigenvalue Problems

All truths are easy to understand once they are discovered; the point is to discover them.

---

Galileo Galilei (1564–1642)

The study of nonlinear eigenvalue problems for quasilinear operators on unbounded domains involving the  $p$ -Laplacian is motivated by various applications. We refer only to fluid mechanics, to mathematical models of the torsional creep, and to nonlinear field equations arising in quantum mechanics. For instance, in fluid mechanics, the shear stress  $\vec{\tau}$  and the velocity gradient  $\nabla_p u$  of certain fluids obey a relation of the form  $\vec{\tau}(x) = a(x)\nabla_p u(x)$ , where  $\nabla_p u := |\nabla u|^{p-2}\nabla u$  and  $p > 1$  is an arbitrary real number. The case  $p = 2$  (respectively  $p < 2$ ,  $p > 2$ ) corresponds to a Newtonian (respectively pseudo-plastic, dilatant) fluid. Then the resulting equations of motion involve  $\operatorname{div}(a\nabla_p u)$ , which reduces to  $a\Delta_p u = a \operatorname{div}(\nabla_p u)$ , provided that  $a$  is a constant.

The  $p$ -Laplace operator also appears in the study of flows through porous media ( $p = 3/2$ , see Showalter-Walkington [279]) or glacial sliding ( $p \in (1, 4/3]$ , see Pélissier-Reynaud [239]). We refer to Aronsson-Janfalk [14] for the mathematical treatment of the Hele-Shaw flow of “power-law fluids”. The concept of Hele-Shaw flow corresponds to a flow between two closely-spaced parallel plates, where the gap between the plates is small compared to the dimension of the plates. Quasilinear problems with a variable coefficient also appear in mathematical models of torsional creeps (elastic for  $p = 2$ , plastic as  $p \rightarrow \infty$ , see Bhattacharya-DiBenedetto-Manfredi [39] and Kawohl [157]). This study is based on the observation that a prismatic material rod subject to a torsional mo-

ment, at sufficiently high temperature and for an extended period of time, exhibits a permanent deformation, called *creep*. The corresponding equations are derived under the assumptions that the components of strain and stress are linked by a power law referred to as the *creep-law* see Kachanov [153, Chapters IV, VIII], [154], and Findley-Lai-Onaran [122]). A nonlinear field equation in quantum mechanics involving the  $p$ -Laplacian, for  $p = 6$ , has been proposed in Benci-Fortunato-Pisani [35].

In this chapter we are concerned with the study of the following quasi-linear eigenvalue problem

$$(P_{\lambda,\mu}) \quad \begin{cases} -\operatorname{div}(a(x)|\nabla u|^{p-2}\nabla u) = \lambda\alpha(x,u) + \beta(x,u) & \text{in } \Omega, \\ a(x)|\nabla u|^{p-2}\nabla u \cdot \mathbf{n} + b(x)|u|^{p-2}u = \mu h(x,u) & \text{on } \Gamma, \end{cases}$$

where  $\Omega \subset \mathbb{R}^N$  is an unbounded domain with (possible noncompact) smooth boundary  $\Gamma$ ,  $\mathbf{n}$  denotes the unit outward normal on  $\Gamma$ ,  $a \in L^\infty(\Omega)$ ,  $b : \Gamma \rightarrow \mathbb{R}$ ,  $\alpha, \beta : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$  and  $h : \Gamma \times \mathbb{R} \rightarrow \mathbb{R}$  are functions which satisfy some growth conditions and  $\lambda, \mu \in \mathbb{R}$  are parameters. Our aim is to study the existence and multiplicity of the solutions of problem  $(P_{\lambda,\mu})$  for different values of the parameters  $\lambda, \mu \in \mathbb{R}$  and for various functions  $\alpha, \beta$  and  $h$ . Eigenvalue problems involving the  $p$ -Laplacian have been the subject of much recent interest, see for example the papers of Allegretto-Huang [5], Anane [12], Drábek [94], Drábek-Pohozaev [96], Drábek-Simader [97], García-Montefusco-Peral [124], García-Peral [125].

In the first section we describe the weighted Sobolev spaces, where we define the energy functional associated to the problem  $(P_{\lambda,\mu})$ . In the next section our attention is focused to prove existence and multiplicity result for problem  $(P_{\lambda,\mu})$ , when  $\mu = 1$ , using the mountain pass theorem and Ljusternik-Schnirelmann type theory. In the last parts of this chapter we prove that the problem  $(P_{\lambda,\mu})$  has two nontrivial weak solutions if the parameters  $\lambda$  and  $\mu$  belongs to some interval.

### 3.1 Weighted Sobolev spaces

Let  $\Omega \subset \mathbb{R}^N$  be an unbounded domain with smooth boundary  $\Gamma$ . We assume that  $p, q$  and  $m$  are real numbers satisfying

$$1 < p < N, \quad p < q < p^* = \frac{Np}{N-p}, \quad (p^* = +\infty, \text{ if } p \geq N), \quad (3.1)$$

and

$$q \leq m < \frac{p(N-1)}{N-p} \text{ if } p < N \text{ (} q \leq m < +\infty, \text{ when } p \geq N\text{)}. \quad (3.2)$$

Let  $C_\delta^\infty(\Omega)$  be the space of  $C_0^\infty(\mathbb{R}^N)$ -functions restricted on  $\Omega$ .

We define the weighted Sobolev space  $E$  as the completion of  $C_\delta^\infty(\Omega)$  in the norm

$$\|u\|_E = \left( \int_{\Omega} \left( |\nabla u(x)|^p + \frac{1}{(1+|x|)^p} |u(x)|^p \right) dx \right)^{1/p}.$$

Denote by  $L^p(\Omega; w_1)$  and  $L^m(\Gamma; w_2)$  the weighted Lebesgue spaces with weight functions

$$w_i(x) = (1+|x|)^{\beta_i}, \quad (i = 1, 2), \quad (3.3)$$

and the norms defined by

$$\|u\|_{p, w_1}^p = \int_{\Omega} w_1 |u(x)|^p dx, \quad \|u\|_{q, w_2}^q = \int_{\Gamma} w_2 |u(x)|^q dx$$

where

$$-N < \beta_1 < -p, \text{ if } p < N \text{ (} \beta_1 < -p, \text{ when } p \geq N\text{)}, \quad (3.4)$$

$$-N < \beta_2 < q \frac{N-p}{p} - N + 1, \text{ if } p < N \text{ (} -N < \beta_2 < 0, \text{ when } p \geq N\text{)}. \quad (3.5)$$

We have the following weighted Hardy-type inequality.

**Lemma 3.1** *Let  $1 < p < N$ . Then, there exist positive constants  $C_1$  and  $C_2$ , such that for every  $u \in E$*

$$\int_{\Omega} \frac{1}{(1+|x|)^p} |u|^p dx \leq C_1 \int_{\Omega} |\nabla u|^p dx + C_2 \int_{\Gamma} \frac{|\mathbf{n} \cdot x|}{(1+|x|)^p} |u|^p d\Gamma. \quad (3.6)$$

*Proof* Using the divergence theorem we obtain for all  $u \in C_\delta^\infty(\Omega)$ ,

$$\begin{aligned} \int_{\Omega} x \cdot \nabla \left( \frac{1}{(1+|x|)^p} |u|^p \right) dx &= \int_{\Gamma} (\mathbf{n} \cdot x) \frac{1}{(1+|x|)^p} |u|^p d\Gamma \\ &\quad - N \int_{\Omega} \frac{1}{(1+|x|)^p} |u|^p dx. \end{aligned}$$

This implies

$$\begin{aligned} N \int_{\Omega} \frac{1}{(1+|x|)^p} |u|^p dx &\leq \int_{\Gamma} \frac{|\mathbf{n} \cdot x|}{(1+|x|)^p} |u|^p d\Gamma + p \int_{\Omega} \frac{1}{(1+|x|)^p} |u|^p dx \\ &\quad + p \int_{\Omega} \frac{1}{(1+|x|)^{p-1}} |u|^{p-1} |\nabla u| dx. \end{aligned}$$

Using Hölder's and Young's inequalities, the last term can be estimated by

$$\begin{aligned} &p \left( \int_{\Omega} \frac{1}{(1+|x|)^p} |u|^p dx \right)^{(p-1)/p} \left( \int_{\Omega} |\nabla u|^p dx \right)^{1/p} \\ &\leq \varepsilon(p-1) \int_{\Omega} \frac{1}{(1+|x|)^p} |u|^p dx + \varepsilon^{1-p} \int_{\Omega} |\nabla u|^p dx, \end{aligned}$$

where  $\varepsilon > 0$  is an arbitrary real number. It follows that

$$\begin{aligned} (N - \varepsilon(p-1) - p) \int_{\Omega} \frac{1}{(1+|x|)^p} |u|^p dx &\leq \varepsilon^{1-p} \int_{\Omega} |\nabla u|^p dx \\ &\quad + \int_{\Gamma} \frac{|\mathbf{n} \cdot x|}{(1+|x|)^p} |u|^p d\Gamma, \end{aligned}$$

and for  $\varepsilon$  small enough, the desired inequality follows by standard density arguments.  $\square$

In this chapter we use the following embedding result, see Pflüger [241] and [242].

**Theorem 3.1** *If*

$$p \leq r \leq \frac{pN}{N-p} \quad \text{and} \quad -N < \beta_1 \leq r \frac{N-p}{p} - N, \quad (3.7)$$

*then the embedding  $E \hookrightarrow L^r(\Omega; w_1)$  is continuous. If the upper bounds for  $r$  in (3.7) are strict, then the embedding is compact. If*

$$p \leq q \leq \frac{p(N-1)}{N-p} \quad \text{and} \quad -N < \beta_2 \leq q \frac{N-p}{p} - N + 1, \quad (3.8)$$

*then the trace operator  $E \hookrightarrow L^q(\Gamma; w_2)$  is continuous. If the upper bounds for  $q$  in (3.8) are strict, then the trace operator is compact.*

As a corollary of Lemma 3.1 and Theorem 3.1 we obtain:

**Lemma 3.2** *Let  $b$  satisfying  $\frac{c}{(1+|x|)^{p-1}} \leq b(x) \leq \frac{C}{(1+|x|)^{p-1}}$  for some constants  $0 < c \leq C$ . Then*

$$\|u\|_b^p = \int_{\Omega} a(x) |\nabla u|^p dx + \int_{\Gamma} b(x) |u|^p d\Gamma$$

defines an equivalent norm on  $E$ .

*Proof* The inequality  $\|u\|_E \leq C_1 \|u\|_b$  follows directly from Lemma 3.1, while Theorem 3.1 (for  $p = q$  and  $\beta_2 = -(p - 1)$ ) implies

$$\begin{aligned} \|u\|_b^p &\leq \|a\|_{L^\infty} \int_{\Omega} |\nabla u|^p dx + C \int_{\Gamma} |u|^p (1 + |x|)^{-(p-1)} d\Gamma \\ &\leq \|a\|_{L^\infty} \int_{\Omega} |\nabla u|^p dx + C_2 \|u\|_E^p, \end{aligned}$$

which shows the desired equivalence.  $\square$

We assume that  $h : \Gamma \times \mathbb{R} \rightarrow \mathbb{R}$  is a Carathéodory function that satisfies the following conditions:

( $\mathcal{H}$ )  $|h(x, s)| \leq h_0(x) + h_1(x)|s|^{m-1}$ ;  $p \leq m < p \cdot \frac{N-1}{N-p}$ , where  $h_i : \Gamma \rightarrow \mathbb{R}$  are nonnegative, measurable functions such that

$$0 \leq h_i(x) \leq C_h w_2 \text{ a.e.}, \quad h_0 \in L^{m/(m-1)}(\Gamma; w_2^{1/(1-m)}),$$

where  $-N < \beta_2 < m \cdot \frac{N-p}{p} - N + 1$  and  $w_2$  is defined as in (3.3).

Set  $H(x, s) = \int_0^s h(x, t) dt$ . We denote by  $N_h, N_H$  the corresponding Nemytskii operators.

**Lemma 3.3** *The operators*

$$N_h : L^m(\Gamma; w_2) \rightarrow L^{m/(m-1)}(\Gamma; w_2^{1/(1-m)}), \quad N_H : L^m(\Gamma; w_2) \rightarrow L^1(\Gamma)$$

are bounded and continuous.

*Proof* Let  $m' = m/(m-1)$  and  $u \in L^m(\Gamma; w_2)$ . Then, by ( $\mathcal{H}$ ),

$$\begin{aligned} \int_{\Gamma} |N_h(u)|^{m'} \cdot w_2^{1/(1-m)} d\Gamma &\leq \\ &\leq 2^{m'-1} \left( \int_{\Gamma} h_0^{m'} \cdot w_2^{1/(1-m)} d\Gamma + \int_{\Gamma} h_1^{m'} |u|^m \cdot w_2^{1/(1-m)} d\Gamma \right) \leq \\ &\leq 2^{m'-1} \left( C + C_h \cdot \int_{\Gamma} |u|^m \cdot w_2 d\Gamma \right), \end{aligned}$$

which shows that  $N_h$  is bounded. In a similar way we obtain

$$\begin{aligned} \int_{\Gamma} |N_H(u)| d\Gamma &\leq \int_{\Gamma} h_0 |u| d\Gamma + \int_{\Gamma} h_1 |u|^m d\Gamma \leq \\ &\leq \left( \int_{\Gamma} h_0^{m'} \cdot w_2^{1/(1-m)} d\Gamma \right)^{1/m'} \cdot \left( \int_{\Gamma} |u|^m \cdot w_2 d\Gamma \right)^{1/m} \\ &+ C_g \cdot \int_{\Gamma} |u|^m \cdot w_2 d\Gamma, \end{aligned}$$

which imply the boundedness of  $N_H$ .

From the usual properties of Nemytskii operators we deduce the continuity of  $N_h$  and  $N_H$ .  $\square$

### 3.2 Eigenvalue problems

We suppose that  $p, q$  and  $m$  are real numbers satisfying the conditions (3.1) and (3.2). In this section we consider the particular case for  $(P_{\lambda, \mu})$ , when  $\alpha(x, u) = f(x)|u|^{p-2}u$ ,  $\beta(x, u) = g(x)|u|^{q-2}u$  and  $\mu = 1$ . In this case problem  $(P_{\lambda, \mu})$  becomes

$$(P_{\lambda}) \begin{cases} -\operatorname{div}(a(x)|\nabla u|^{p-2}\nabla u) = \lambda f(x)|u|^{p-2}u + g(x)|u|^{q-2}u & \text{in } \Omega, \\ a(x)|\nabla u|^{p-2}\nabla u \cdot \mathbf{n} + b(x)|u|^{p-2}u = h(x, u) & \text{on } \Gamma, \end{cases}$$

where  $\mathbf{n}$  denotes the unit outward normal on  $\Gamma$ ,  $0 < a_0 \leq a$ , where  $a \in L^\infty(\Omega)$  and  $b : \Gamma \rightarrow \mathbb{R}$  is a continuous function satisfying

$$\frac{c}{(1+|x|)^{p-1}} \leq b(x) \leq \frac{C}{(1+|x|)^{p-1}},$$

for some constants  $0 < c \leq C$ .

We assume that  $f$  and  $g$  are nontrivial measurable functions satisfying  $0 \leq f(x) \leq C(1+|x|)^{\alpha_1}$  and  $0 \leq g(x) \leq C(1+|x|)^{\alpha_2}$ , for a.e.  $x \in \Omega$ , where the numbers  $\alpha_1, \alpha_2$  satisfy the conditions

$$-N < \alpha_1 < -p, \quad \text{if } p < N (\alpha_1 < -p, \quad \text{when } p \geq N), \quad (3.9)$$

and

$$-N < \alpha_2 < q \frac{N-p}{p} - N + 1, \quad \text{if } p < N (-N < \alpha_2 < 0, \quad \text{when } p \geq N). \quad (3.10)$$

The mapping  $h : \Gamma \times \mathbb{R} \rightarrow \mathbb{R}$  is a Carathéodory function which fulfills the assumption

$$(H1) \quad |h(x, s)| \leq h_0(x) + h_1(x)|s|^{m-1},$$

where  $h_i : \Gamma \rightarrow \mathbb{R}$  ( $i = 0, 1$ ) are measurable functions satisfying

$$h_0 \in L^{m/(m-1)}(\Gamma; w_3^{1/(1-m)}) \quad \text{and} \quad 0 \leq h_i \leq C_h w_3 \quad \text{a.e. on } \Gamma,$$

and  $w_3 = (1 + |x|)^{\alpha_3}$ , where  $\alpha_3$  satisfies the conditions

$$-N < \alpha_3 < m \frac{N-p}{p} - N + 1, \quad \text{if } p < N \quad (-N < \alpha_3 < 0 \quad \text{when } p \geq N). \quad (3.11)$$

We also assume:

$$(H2) \quad \lim_{s \rightarrow 0} \frac{h(x, s)}{b(x)|s|^{p-1}} = 0 \quad \text{uniformly in } x.$$

(H3) There exists  $\mu \in (p, q]$  such that

$$\mu H(x, t) \leq t h(x, t) \quad \text{for a.e. } x \in \Gamma \text{ and every } t \in \mathbb{R}.$$

(H4) There exists a nonempty open set  $U \subset \Gamma$  with  $H(x, t) > 0$  for  $(x, t) \in U \times (0, \infty)$ , where  $H(x, t) = \int_0^t h(x, s) ds$ .

By a weak solution of problem  $(P_\lambda)$  we mean a function  $u \in E$  such that for any  $v \in E$  it holds

$$\begin{aligned} & \int_{\Omega} a(x) |\nabla u|^{p-2} \nabla u \nabla v \, dx + \int_{\Gamma} b(x) |u|^{p-2} uv \, dS \\ &= \lambda \int_{\Omega} f(x) |u|^{p-2} uv \, dx + \int_{\Omega} g(x) |u|^{q-2} uv \, dx + \int_{\Gamma} h(x, u) v \, dS. \end{aligned}$$

The energy functional corresponding to  $(P_\lambda)$  is defined as  $\mathcal{E}_\lambda : E \rightarrow \mathbb{R}$

$$\begin{aligned} \mathcal{E}_\lambda(u) &= \frac{1}{p} \int_{\Omega} a(x) |\nabla u|^p \, dx + \frac{1}{p} \int_{\Gamma} b(x) |u|^p \, dS \\ &\quad - \frac{\lambda}{p} \int_{\Omega} f(x) |u|^p \, dx - \int_{\Gamma} H(x, u) \, dS - \frac{1}{q} \int_{\Omega} g(x) |u|^q \, dx. \end{aligned}$$

By Lemma 3.2 we have  $\|\cdot\|_b \simeq \|\cdot\|_E$ . We may write

$$\mathcal{E}_\lambda(u) = \frac{1}{p} \|u\|_b^p - \frac{\lambda}{p} \int_{\Omega} f(x) |u|^p \, dx - \int_{\Gamma} H(x, u) \, dS - \frac{1}{q} \int_{\Omega} g(x) |u|^q \, dx.$$

Since  $p < q < p^*$ ,  $-N < \alpha_1 < -p$  and  $-N < \alpha_2 < q \frac{N-p}{p} - N$ , we can apply Theorem 3.1 and obtain that the embedding  $E \hookrightarrow L^p(\Omega; w_1)$ ,

$E \hookrightarrow L^q(\Omega; w_2)$  and  $E \hookrightarrow L^q(\Gamma; w_w)$  are compact. So, the functional  $\mathcal{E}_\lambda$  is well defined.

**Lemma 3.4** *Under the assumptions (H1)-(H4), the functional  $\mathcal{E}_\lambda$  is Fréchet differentiable on  $E$  and satisfies the Palais-Smale condition.*

*Proof* Denote  $\Phi(x, u) = \frac{1}{p}f(x)|u|^p$  and  $\Psi(x, u) = \frac{1}{q}g(x)|u|^q$ . Set  $I(u) = \frac{1}{p}\|u\|_b^p$ ,  $K_H(u) = \int_\Gamma H(x, u) dS$ ,  $K_\Psi(u) = \int_\Omega \Psi(x, u) dx$  and  $K_\Phi(u) = \int_\Omega \Phi(x, u) dx$ .

Then the directional derivative of  $\mathcal{E}'_\lambda$  in the direction  $v \in E$  is

$$\langle \mathcal{E}'_\lambda(u), v \rangle = \langle I'(u), v \rangle - \lambda \langle K'_\Phi(u), v \rangle - \langle K'_\Psi(u), v \rangle - \langle K'_H(u), v \rangle,$$

where

$$\langle I'(u), v \rangle = \int_\Omega a(x)|\nabla u|^{p-2} \nabla u \nabla v dx + \int_\Gamma b(x)|u|^{p-2} uv dS,$$

$$\langle K'_H(u), v \rangle = \int_\Gamma h(x, u)v dS,$$

$$\langle K'_\Psi(u), v \rangle = \int_\Omega g(x)|u|^{q-2} uv dx,$$

$$\langle K'_\Phi(u), v \rangle = \int_\Omega f(x)|u|^{p-2} uv dx.$$

Clearly,  $I' : E \rightarrow E^*$  is continuous. The operator  $K'_H$  is a composition of the operators

$$K'_H : E \rightarrow L^m(\Gamma; w_3) \xrightarrow{N_h} L^{m/(m-1)}(\Gamma; w_3^{1/(1-m)}) \xrightarrow{l} E^*$$

where  $\langle l(u), v \rangle = \int_\Gamma uv dS$ . Since

$$\int_\Gamma |uv| dS \leq \left( \int_\Gamma |u|^{m'} w_3^{1/(1-m)} dS \right)^{1/m'} \cdot \left( \int_\Gamma |v|^m w_3 dS \right)^{1/m},$$

then by Theorem 3.1  $l$  is continuous. As a composition of continuous operators,  $K'_H$  is continuous, too. Moreover, by our assumptions on  $w_3$ , the trace operator  $E \hookrightarrow L^m(\Gamma; w_3)$  is compact and therefore,  $K'_H$  is also compact.

Set  $\varphi(u) = f(x)|u|^{p-2}u$ . Lemma 3.3 implies that the Nemytskii operator corresponding to any function which satisfies (H1) is bounded and continuous. Hence  $N_h$  and  $N_\varphi$  are bounded and continuous. Note that

$$K'_\Phi : E \subset L^p(\Omega; w_1) \xrightarrow{N_\varphi} L^{p/(p-1)}(\Omega; w_1^{1/(1-p)}) \xrightarrow{\eta} E^*,$$

where  $\langle \eta(u), v \rangle = \int_{\Omega} uv \, dx$ . Since

$$\int_{\Omega} |uv| \, dx \leq \left( \int_{\Omega} |u|^{p/(p-1)} w_1^{1/(1-p)} \, dx \right)^{(p-1)/p} \cdot \left( \int_{\Omega} |v|^p w_1 \, dx \right)^{1/p},$$

it follows that  $\eta$  is continuous. But  $K'_{\Phi}$  is the composition of three continuous operators and by the assumptions on  $w_1$ , the embedding  $E \hookrightarrow L^p(\Omega; w_1)$  is compact. This implies that  $K'_{\Phi}$  is compact. In a similar way we obtain that  $K'_{\Psi}$  is compact and this implies the continuous Fréchet differentiability of  $\mathcal{E}_{\lambda}$ .

Now, let  $u_n \in E$  be a Palais-Smale sequence, that is,

$$|\mathcal{E}_{\lambda}(u_n)| \leq C \text{ for all } n \quad (3.12)$$

and

$$\|\mathcal{E}'_{\lambda}(u_n)\|_{E^*} \rightarrow 0 \text{ as } n \rightarrow \infty. \quad (3.13)$$

We first prove that  $\{u_n\}$  is bounded in  $E$ . Remark that (3.13) implies that

$$|\langle \mathcal{E}'_{\lambda}(u_n), u_n \rangle| \leq \mu \cdot \|u_n\|_b \text{ for } n \text{ large enough.}$$

Then by (3.12) we obtain

$$C + \|u_n\|_b \geq \mathcal{E}_{\lambda}(u_n) - \frac{1}{\mu} \langle \mathcal{E}'_{\lambda}(u_n), u_n \rangle. \quad (3.14)$$

But

$$\begin{aligned} \langle \mathcal{E}'_{\lambda}(u_n), u_n \rangle &= \int_{\Omega} a(x) |\nabla u_n|^p \, dx + \int_{\Gamma} b(x) |u_n|^p \, dS \\ &\quad - \lambda \int_{\Omega} f(x) |u_n|^p \, dx - \int_{\Omega} g(x) |u_n|^q \, dx - \int_{\Gamma} h(x, u_n) u_n \, dS. \end{aligned}$$

We have

$$\begin{aligned} \mathcal{E}_{\lambda}(u_n) - \frac{1}{\mu} \langle \mathcal{E}'_{\lambda}(u_n), u_n \rangle &= \left( \frac{1}{p} - \frac{1}{\mu} \right) \left( \|u_n\|_b^p - \lambda \int_{\Omega} f(x) |u|^p \, dx \right) \\ &\quad - \left( \int_{\Gamma} H(x, u_n) \, dS - \frac{1}{\mu} \int_{\Gamma} h(x, u_n) u_n \, dS \right) - \left( \frac{1}{q} - \frac{1}{\mu} \right) \int_{\Omega} g(x) |u_n|^q \, dx. \end{aligned}$$

By (H3) we deduce that

$$\int_{\Gamma} H(x, u_n) \, dS \leq \frac{1}{\mu} \int_{\Gamma} h(x, u_n) u_n \, dS. \quad (3.15)$$

Therefore

$$\mathcal{E}_\lambda(u_n) - \frac{1}{\mu} \langle \mathcal{E}'_\lambda(u_n), u_n \rangle \geq \left( \frac{1}{p} - \frac{1}{\mu} \right) C_0 \|u_n\|_b^p. \quad (3.16)$$

Relations (3.14) and (3.16) yield

$$C + \|u_n\|_b \geq \left( \frac{1}{p} - \frac{1}{\mu} \right) C_0 \|u_n\|_b^p.$$

This shows that  $\{u_n\}$  is bounded in  $E$ .

To prove that  $\{u_n\}$  contains a Cauchy sequence we use the following well-known inequalities:

$$|\xi - \zeta|^p \leq C(|\xi|^{p-2}\xi - |\zeta|^{p-2}\zeta)(\xi - \zeta), \quad \text{for } p \geq 2, \quad (3.17)$$

$$|\xi - \zeta|^2 \leq C(|\xi|^{p-2}\xi - |\zeta|^{p-2}\zeta)(\xi - \zeta)(|\xi| + |\zeta|)^{2-p}, \quad \text{for } 1 < p < 2, \quad (3.18)$$

for every  $\xi, \zeta \in \mathbb{R}^N$ .

In the case  $p \geq 2$  we obtain:

$$\begin{aligned} \|u_n - u_k\|_b^p &= \int_{\Omega} a(x) |\nabla u_n - \nabla u_k|^p dx + \int_{\Gamma} b(x) |u_n - u_k|^p dS \\ &\leq C(\langle I'(u_n), u_n - u_k \rangle - \langle I'(u_k), u_n - u_k \rangle) \\ &= C(\langle \mathcal{E}'_\lambda(u_n), u_n - u_k \rangle - \langle \mathcal{E}'_\lambda(u_k), u_n - u_k \rangle \\ &\quad + \lambda \langle K'_\Phi(u_n), u_n - u_k \rangle - \lambda \langle K'_\Phi(u_k), u_n - u_k \rangle \\ &\quad + \langle K'_H(u_n), u_n - u_k \rangle - \langle K'_H(u_k), u_n - u_k \rangle \\ &\quad + \langle K'_\Psi(u_n), u_n - u_k \rangle - \langle K'_\Psi(u_k), u_n - u_k \rangle) \\ &\leq C(\|\mathcal{E}'_\lambda(u_n)\|_{E^*} + \|\mathcal{E}'_\lambda(u_k)\|_{E^*} + |\lambda| \|K'_\Phi(u_n) - K'_\Phi(u_k)\|_{E^*} \\ &\quad + \|K'_H(u_n) - K'_H(u_k)\|_{E^*} \\ &\quad + \|K'_\Psi(u_n) - K'_\Psi(u_k)\|_{E^*}) \|u_n - u_k\|_b. \end{aligned}$$

Since  $\mathcal{E}'_\lambda(u_n) \rightarrow 0$  and  $K'_\Phi, K'_\Psi, K'_H$  are compact, we conclude (passing eventually to a subsequence) that  $\{u_n\}$  converges in  $E$ .

If  $1 < p < 2$ , then we use the estimate

$$\|u_n - u_k\|_b^2 \leq C' |\langle I'(u_n), u_n - u_k \rangle - \langle I'(u_k), u_n - u_k \rangle| (\|u_n\|_b^{2-p} + \|u_k\|_b^{2-p}). \quad (3.19)$$

Since  $\|u_n\|_b$  is bounded, the same arguments lead to a convergent subsequence. In order to prove the estimate (3.19) we recall the following result: for all  $s \in (0, \infty)$  there is a constant  $C_s > 0$  such that

$$(x + y)^s \leq C_s(x^s + y^s) \quad \text{for any } x, y \in (0, \infty). \quad (3.20)$$

Then we obtain

$$\begin{aligned} \|u_n - u_k\|_b^2 &= \left( \int_{\Omega} a(x) |\nabla u_n - \nabla u_k|^p dx + \int_{\Gamma} b(x) |u_n - u_k|^p dS \right)^{2/p} \leq \\ C_p &\left[ \left( \int_{\Omega} a(x) |\nabla u_n - \nabla u_k|^p dx \right)^{2/p} + \left( \int_{\Gamma} b(x) |u_n - u_k|^p dS \right)^{2/p} \right]. \end{aligned} \quad (3.21)$$

Using relations (3.18), (3.20) and the Hölder inequality we find

$$\begin{aligned} &\int_{\Omega} a(x) |\nabla u_n - \nabla u_k|^p dx = \int_{\Omega} a(x) (|\nabla u_n - \nabla u_k|^2)^{p/2} dx \leq \\ &C \int_{\Omega} a(x) (|\nabla u_n|^{p-2} \nabla u_n - |\nabla u_k|^{p-2} \nabla u_k) (\nabla u_n - \nabla u_k) dx \times \\ &(|\nabla u_n| + |\nabla u_k|)^{p(2-p)/2} dx = \\ &C \int_{\Omega} (a(x) (|\nabla u_n|^{p-2} \nabla u_n - |\nabla u_k|^{p-2} \nabla u_k) (\nabla u_n - \nabla u_k))^{p/2} \times \\ &(a(x) (|\nabla u_n| + |\nabla u_k|)^p)^{(2-p)/2} dx \leq \\ &C \left( \int_{\Omega} a(x) (|\nabla u_n|^{p-2} \nabla u_n - |\nabla u_k|^{p-2} \nabla u_k) (\nabla u_n - \nabla u_k) dx \right)^{p/2} \times \\ &\left( \int_{\Omega} a(x) (|\nabla u_n| + |\nabla u_k|)^p dx \right)^{(2-p)/2} \leq \\ &\tilde{C}_p \left( \int_{\Omega} a(x) |\nabla u_n|^p dx + \int_{\Omega} a(x) |\nabla u_k|^p dx \right)^{(2-p)/2} \\ &\left( \int_{\Omega} a(x) (|\nabla u_n|^{p-2} \nabla u_n - |\nabla u_k|^{p-2} \nabla u_k) (\nabla u_n - \nabla u_k) dx \right)^{p/2} \leq \\ &\bar{C}_p \left[ \left( \int_{\Omega} a(x) |\nabla u_n|^p dx \right)^{(2-p)/2} + \left( \int_{\Omega} a(x) |\nabla u_k|^p dx \right)^{(2-p)/2} \right] \times \\ &\left( \int_{\Omega} a(x) (|\nabla u_n|^{p-2} \nabla u_n - |\nabla u_k|^{p-2} \nabla u_k) (\nabla u_n - \nabla u_k) dx \right)^{p/2} \leq \\ &\bar{C}_p \left[ \int_{\Omega} a(x) (|\nabla u_n|^{p-2} \nabla u_n - |\nabla u_k|^{p-2} \nabla u_k) (\nabla u_n - \nabla u_k) dx \right]^{p/2} \times \\ &\left( \|u_n\|_b^{(2-p)p/2} + \|u_k\|_b^{(2-p)p/2} \right). \end{aligned}$$

Using the last inequality and relation (3.20) we have the estimate

$$\begin{aligned} &\left( \int_{\Omega} a(x) |\nabla u_n - \nabla u_k|^p dx \right)^{2/p} \quad (3.22) \\ &\leq C'_p \left( \int_{\Omega} a(x) (|\nabla u_n|^{p-2} \nabla u_n - |\nabla u_k|^{p-2} \nabla u_k) (\nabla u_n - \nabla u_k) dx \right) \\ &\cdot (\|u_n\|_b^{2-p} + \|u_k\|_b^{2-p}). \end{aligned}$$

In a similar way we deduce the estimate

$$\begin{aligned} & \left( \int_{\Gamma} b(x) |u_n - u_k|^p dS \right)^{2/p} \leq \\ & C'_p \left( \int_{\Gamma} b(x) (|u_n|^{p-2} u_n - |u_k|^{p-2} u_k) (u_n - u_k) dx \right) \cdot \\ & (\|u_n\|_b^{2-p} + \|u_k\|_b^{2-p}). \end{aligned} \quad (3.23)$$

At this stage, inequalities (3.21), (3.22) and (3.23) imply the estimate (3.19). The proof of Lemma 3.4 is complete.  $\square$

Define the Rayleigh quotient

$$\tilde{\lambda} := \inf_{u \in E; u \neq 0} \left( \frac{\int_{\Omega} a(x) |\nabla u|^p dx + \int_{\Gamma} b(x) |u|^p dS}{\int_{\Omega} f(x) |u|^p dx} \right).$$

We have the following existence result.

**Theorem 3.2** *Assume that the conditions (H1)-(H4) hold. Then, for every  $\lambda < \tilde{\lambda}$ , problem  $(P_{\lambda})$  has a nontrivial weak solution.*

Our hypothesis  $\lambda < \tilde{\lambda}$  implies the existence of some  $C_0 > 0$  such that, for every  $v \in E$

$$\|v\|_b^p - \lambda \int_{\Omega} f(x) |v|^p dx \geq C_0 \|v\|_b^p. \quad (3.24)$$

*Proof of Theorem 3.2.* We have to verify the geometric assumptions of the mountain pass theorem, see Theorem 1.7. First we show that there exist positive constants  $R$  and  $c_0$  such that

$$\mathcal{E}_{\lambda}(u) \geq c_0, \quad \text{for any } u \in E \text{ with } \|u\| = R. \quad (3.25)$$

By Theorem 3.1 there exists  $A > 0$  such that

$$\|u\|_{q,w_2}^q \leq A \|u\|_b^q \quad \text{for all } u \in E.$$

This inequality together with (3.24) imply that

$$\begin{aligned} \mathcal{E}_{\lambda}(u) &= \frac{1}{p} (\|u\|_b^p - \lambda \|u\|_{p,w_1}^p) - \frac{1}{q} \int_{\Omega} g(x) |u|^q dx - \int_{\Gamma} H(x, u) dS \\ &\geq \frac{C_0}{p} \|u\|_b^p - \frac{A}{q} \|u\|_b^q - \int_{\Gamma} H(x, u) dS. \end{aligned}$$

By (H1) and (H2) we deduce that for every  $\varepsilon > 0$  there exists  $C_\varepsilon > 0$  such that

$$\frac{1}{q}|g(x)||u|^q \leq \varepsilon b(x)|u|^p + C_\varepsilon w_3(x)|u|^m.$$

Consequently,

$$\begin{aligned} \mathcal{E}_\lambda(u) &\geq \frac{C_0}{p}\|u\|_b^p - \frac{A}{q}\|u\|_b^q - \int_\Gamma (\varepsilon b(x)|u|^p + C_\varepsilon w_3(x)|u|^m) dS \\ &\geq \frac{C_0}{p}\|u\|_b^p - \frac{A}{q}\|u\|_b^q - \varepsilon c_1\|u\|_b^p - C_\varepsilon C_2\|u\|_b^m. \end{aligned}$$

For  $\varepsilon > 0$  and  $R > 0$  small enough, we deduce that for every  $u \in E$  with  $\|u\|_b = R$  we have  $\mathcal{E}_\lambda(u) \geq c_0 > 0$ , which yields (3.25).

In what follows we verify the second geometric assumption of Theorem 1.7, namely

$$\exists v \in E \text{ with } \|v\| > R \text{ such that } \mathcal{E}_\lambda(v) < c_0. \quad (3.26)$$

Choose  $\psi \in C_\delta^\infty(\Omega)$ ,  $\psi \geq 0$ , such that  $\emptyset \neq \text{supp}\psi \cap \Gamma \subset U$ . From

$$\frac{1}{q}g(x)u^q \geq c_3u^\mu - c_4 \quad \text{on } (u, s) \in U \times (0, \infty)$$

and (H1) we claim that

$$\begin{aligned} \mathcal{E}_\lambda(t\psi) &= \frac{t^p}{p}(\|\psi\|_b^p - \lambda\|\psi\|_{p,w_1}^p) - \frac{1}{q} \int_\Omega g(x)|t\psi|^q dx - \int_\Gamma H(x, t\psi) dS \\ &\leq \frac{t^p}{p}(\|\psi\|_b^p - \lambda\|\psi\|_{p,w_1}^p) - c_3t^\mu \int_U \psi^\mu dS + c_4|U| - \frac{t^q}{q} \int_\Omega w_2\psi^q dx. \end{aligned}$$

Since  $q \geq \mu > p$ , we obtain  $\mathcal{E}_\lambda(t\psi) \rightarrow -\infty$  as  $t \rightarrow \infty$ . It follows that if  $t > 0$  is large enough,  $\mathcal{E}_\lambda(t\psi) < 0$ , so  $v = t\psi$  satisfies (3.26).

By Theorem 1.7 it follows that problem  $(P_\lambda)$  has a nontrivial weak solution.  $\square$

In the special case  $h(x, s) \equiv 0$ , if we use Theorem 1.12, we are able to show that, the problem  $(P_\lambda)$  has an infinitely many solutions. The statement is the following:

**Theorem 3.3** *Assume  $h(x, s) \equiv 0$  and  $q \geq 2$ . Then, for every  $\lambda < \tilde{\lambda}$ , problem  $(P_\lambda)$  possesses infinitely many solutions.*

*Proof* Consider the even functional

$$J(v) = \frac{1}{p} \int_\Omega a(x)|\nabla v|^p dx + \frac{1}{p} \int_\Gamma b(x)|v|^p dS - \frac{\lambda}{p} \int_\Omega f(x)|v|^p dx,$$

on the closed symmetric manifold

$$M = \{v \in E : \int_{\Omega} g(x)|v|^q = 1\}.$$

Note that  $M$  is only a  $C^1$ -manifold, since we have assumed  $1 < p < q$ . By our hypotheses on  $f$ ,  $g$ ,  $b$  and  $h$  (note that (H1)-(H4) are easily satisfied), Lemma 3.4 and Theorem 1.12, we have that  $J|_M$  possesses at least  $\gamma(M)$  pairs of critical points (where  $\gamma(M)$  stands for the genus of  $M$ ).

Now we have to estimate  $\gamma(M)$ . Since  $g \not\equiv 0$  there exists an open set  $\omega \subset \Omega$  such that  $g(x) \geq \delta > 0$  on  $\omega$ . By the properties of the genus, it follows that  $\gamma(\omega) \geq \gamma(B)$ , where  $B$  is the unit ball of  $W_0^{1,p}(\omega) \subset E$ . Since the genus of the unit ball in an infinite dimensional Banach space is infinity, we have  $\gamma(M) = \infty$ . Hence, there exists a sequence  $\{v_n\} \subset E$ , such that any  $v_n$  (and also  $-v_n$ ) is a constrained critical point of  $J$  on  $M$ .

By the Lagrange multipliers rule we obtain that there exists a sequence  $\{\lambda_n\} \subset \mathbb{R}$  such that

$$\int_{\Omega} a(x)|\nabla v_n|^p dx + \int_{\Gamma} b(x)|v_n|^p dS - \lambda \int_{\Omega} f(x)|v_n|^p dx = \lambda_n \int_{\Omega} g(x)|v_n|^q dx.$$

Since  $v_n \in M$ , using our assumption  $\lambda < \tilde{\lambda}$  we find

$$\lambda_n = \|v_n\|_b^p - \lambda \int_{\Omega} f(x)|v_n|^p dx > 0,$$

so we can apply the usual scaling. Setting  $u_n = \lambda_n^{1/(q-p)}v_n$ , we have that  $u_n$  satisfies for any  $n$  the equation

$$\int_{\Omega} a(x)|\nabla u_n|^p dx + \int_{\Gamma} b(x)|u_n|^p dS = \lambda \int_{\Omega} f(x)|u_n|^p dx + \int_{\Omega} g(x)|u_n|^q dx,$$

so the claim is proved.  $\square$

### 3.3 Superlinear case

We assume throughout this section that  $p, q, r$  and  $\alpha_1$  are real numbers satisfying

$$1 < p < N, \quad p < q < r < p^* = \frac{pN}{N-p}, \quad -N < \alpha_1 < q \frac{N-p}{p} - N. \quad (3.27)$$

Also, we consider  $a \in L^\infty(\Omega)$  and  $b \in L^\infty(\Gamma)$  such that

$$a(x) \geq a_0 > 0, \quad \text{for a.e. } x \in \Omega, \quad (3.28)$$

and

$$\frac{c}{(1+|x|)^{p-1}} \leq b(x) \leq \frac{C}{(1+|x|)^{p-1}}, \quad \text{for a.e. } x \in \Gamma, \quad \text{where } c, C > 0. \quad (3.29)$$

We assume that  $h : \Gamma \times \mathbb{R} \rightarrow \mathbb{R}$  is a Carathéodory function, which satisfies the following conditions:

(h1)  $h(\cdot, 0) = 0$ ,  $h(x, s) + h(x, -s) \geq 0$ , for a.e.  $x \in \Gamma$  and for any  $s \in \mathbb{R}$ ;

(h2)  $h(x, s) \leq h_0(x) + h_1(x)|s|^{m-1}$ ,  $p \leq m < p \frac{N-1}{N-p}$ , where  $h_i : \Gamma \rightarrow \mathbb{R}$ ,  $i = 1, 2$  are nonnegative, measurable functions such that

$$0 \leq h_i(x) \leq C_h w_2, \quad \text{a.e. } g_0 \in L^{m/(m-1)}(\Gamma; w_2^{1/(m-1)}),$$

where  $-N < \alpha_2 < m \cdot \frac{N-p}{p} - N + 1$  and  $w_2 = (1+|x|)^{\alpha_2}$ .

Set  $H(x, s) = \int_0^s h(x, t) dt$ . We denote by  $H_h, N_H$  the corresponding Nemytskii operators. From Lemma 3.3 and condition (h2) it follows that the operators  $H_h, N_H$  are bounded and continuous.

In this section we consider the following double eigenvalue problem, referred as  $(P'_{\lambda, \mu})$ :

$$\begin{cases} -\operatorname{div}(a(x)|\nabla u|^{p-2}\nabla u) + h(x)|u|^{r-2}u = \lambda(1+|x|)^{\alpha_1}|u|^{q-2}u & \text{in } \Omega \\ a(x)|\nabla u|^{p-2}\nabla u \cdot \mathbf{n} + b(x) \cdot |u|^{p-2}u = \mu g(x, u) & \text{on } \Gamma \\ u \geq 0, u \not\equiv 0 & \text{in } \Omega. \end{cases}$$

Let  $\chi : \Omega \rightarrow \mathbb{R}$  be a positive and continuous function satisfying

$$\int_{\Omega} \frac{w_1^{r/(r-q)}}{\chi^{q/(r-q)}} dx < \infty. \quad (3.30)$$

Define the Banach space

$$X = \{u \in E : \int_{\Omega} \chi(x)|u|^r dx < \infty\}$$

endowed with the norm

$$\|u\|_X^p = \|u\|_b^p + \left( \int_{\Omega} \chi(x)|u(x)|^r dx \right)^{p/r}.$$

The energy functional  $\Phi : X \rightarrow \mathbb{R}$  corresponding to problem  $(P_{\lambda,\mu})$  is given by

$$\begin{aligned} \Phi(u) &= \frac{1}{p} \int_{\Omega} a(x) |\nabla u|^p dx + \frac{1}{p} \int_{\Gamma} b(x) |u|^p d\Gamma - \frac{\lambda}{q} \int_{\Omega} w_1 |u|^q dx + \\ &+ \frac{1}{r} \int_{\Omega} \chi(x) |u|^r dx - \mu \int_{\Gamma} H(x, u) d\Gamma. \end{aligned}$$

Theorem 3.1 implies that  $\Phi$  is well defined. The solutions of problem  $(P_{\lambda,\mu})$  will be found as critical points of  $\Phi$ . Therefore, a function  $u \in X$  is a solution of problem  $(P_{\lambda,\mu})$  provided that for any  $v \in X$ ,

$$\begin{aligned} \int_{\Omega} a |\nabla u|^{p-2} \nabla u \cdot \nabla v + \int_{\Gamma} b |u|^{p-2} uv &= \lambda \int_{\Omega} w_1 |u|^{p-2} uv \\ &- \int_{\Omega} \chi |u|^{p-2} uv + \mu \int_{\Gamma} hv. \end{aligned}$$

We have the following result.

**Theorem 3.4** *Assume hypotheses (3.27), (3.28), (3.29), (3.30), (h1) and (h2) hold. Then there exist real numbers  $\mu_*$ ,  $\mu^*$  and  $\lambda^* > 0$  such that problem  $(P'_{\lambda,\mu})$  has no nontrivial solution, provided that  $\mu_* < \mu < \mu^*$  and  $0 < \lambda < \lambda^*$ .*

*Proof* Suppose that  $u$  is a solution in  $X$  of  $(P'_{\lambda,\mu})$ . Then  $u$  satisfies

$$\begin{aligned} \int_{\Omega} a(x) |\nabla u|^p dx + \int_{\Gamma} b(x) |u|^p d\Gamma - \mu \int_{\Gamma} h(x, u) u d\Gamma & \quad (3.31) \\ + \int_{\Omega} \chi(x) |u|^r dx &= \lambda \int_{\Omega} w_1 |u|^q dx. \end{aligned}$$

It follows from Young's inequality that

$$\begin{aligned} \lambda \int_{\Omega} w_1 |u|^q dx &= \int_{\Omega} \frac{\lambda w_1}{\chi^{q/r}} \cdot \chi^{q/r} |u|^q dx \leq q \frac{r-q}{r} \lambda^{r/(r-q)} \int_{\Omega} \frac{w_1^{r/(r-q)}}{\chi^{q/(r-q)}} dx \\ &+ \frac{q}{r} \int_{\Omega} \chi |u|^r dx. \end{aligned}$$

This combined with (3.31) gives

$$\begin{aligned} \|u\|_b^p - \mu \int_{\Gamma} h(x, u)u \, d\Gamma &\leq \frac{r-q}{r} \lambda^{r/(r-q)} \int_{\Omega} \frac{w_1^{r/(r-q)}}{\chi^{q/(r-q)}} \, dx + \\ &+ \frac{q-r}{r} \int_{\Omega} \chi |u|^r \, dx \leq \frac{r-q}{r} \lambda^{r/(r-q)} \int_{\Omega} \frac{w_1^{r/(r-q)}}{\chi^{q/(r-q)}} \, dx. \end{aligned} \quad (3.32)$$

Set

$$\begin{aligned} A &= \{u \in X : \int_{\Gamma} h(x, u)u \, d\Gamma < 0\}, \quad B = \{u \in X : \int_{\Gamma} h(x, u)u \, d\Gamma > 0\}, \\ \mu_* &= \sup_{u \in A} \frac{\|u\|_b^p}{\int_{\Gamma} h(x, u)u \, d\Gamma}, \quad \mu^* = \inf_{u \in B} \frac{\|u\|_b^p}{\int_{\Gamma} h(x, u)u \, d\Gamma}. \end{aligned} \quad (3.33)$$

We introduce the convention that, if  $A = \emptyset$ , then  $\mu_* = -\infty$  and if  $B = \emptyset$ , then  $\mu^* = +\infty$ .

We show that, if we take  $\mu_* < \mu < \mu^*$ , then there exists  $C_0 > 0$  such that

$$C_0 \|u\|_b^p \leq \|u\|_b^p - \mu \int_{\Gamma} h(x, u)u \, d\Gamma \quad \text{for all } u \in X. \quad (3.34)$$

If  $\mu < \mu^*$ , then there exists a constant  $C_1 \in (0, 1)$  such that

$$\mu \leq (1 - C_1)\mu^* \leq (1 - C_1) \frac{\|u\|_b^p}{\int_{\Gamma} h(x, u)u \, d\Gamma} \quad \text{for all } u \in B,$$

which implies

$$\|u\|_b^p - \mu \int_{\Gamma} h(x, u)u \, d\Gamma \geq C_1 \|u\|_b^p \quad \text{for all } u \in B. \quad (3.35)$$

If  $\mu_* < \mu$ , then there exists a constant  $C_2 \in (0, 1)$  such that

$$(1 - C_2) \frac{\|u\|_b^p}{\int_{\Gamma} h(x, u)u \, d\Gamma} \leq (1 - C_2)\mu_* \leq \mu \quad \text{for all } u \in A,$$

which yields

$$\|u\|_b^p - \mu \int_{\Gamma} h(x, u)u \, d\Gamma \geq C_2 \|u\|_b^p \quad \text{for all } u \in A. \quad (3.36)$$

From (3.35) and (3.36) we conclude that

$$\|u\|_b^p - \mu \int_{\Gamma} h(x, u) u \, d\Gamma \geq \min\{C_1, C_2\} \|u\|_b^p \quad \text{for all } u \in X$$

and taking  $C_0 = \min\{C_1, C_2\}$  we obtain (3.34).

By (3.31), (3.34) and Theorem 3.1 we have

$$C_0 \bar{C} \left( \int_{\Omega} w_1 |u|^q \, dx \right)^{p/q} \leq C_0 \|u\|_b^p \leq \lambda \int_{\Omega} w_1 |u|^q \, dx, \quad (3.37)$$

for some constant  $\bar{C} > 0$ . This inequality implies

$$(\bar{C} \lambda^{-1} C_0)^{q/(q-p)} \leq \int_{\Omega} w_1 |u|^q \, dx,$$

which combined with (3.37) yields

$$C_0 \bar{C} (\bar{C} \lambda^{-1} C_0)^{p/(q-p)} \leq C_0 \|u\|_b^p.$$

Combining this with (3.32) and (3.34) we obtain

$$C_0 \bar{C} (\bar{C} \lambda^{-1} C_0)^{p/(q-p)} \leq \frac{r-q}{r} \lambda^{r/(r-q)} \int_{\Omega} \frac{w_1^{r/(r-q)}}{\chi^{q/(r-q)}} \, dx.$$

By taking

$$\lambda^* = \left( (C_0 \bar{C})^{q/(q-p)} \frac{r}{r-q} \left( \int_{\Omega} \frac{w_1^{r/(r-q)}}{\chi^{q/(r-q)}} \, dx \right)^{-1} \right)^{\frac{(r-q)(q-p)}{q(r-p)}},$$

the result follows.  $\square$

Set

$$U = \{u \in X : \int_{\Gamma} H(x, u) \, d\Gamma < 0\}, \quad V = \{u \in X : \int_{\Gamma} H(x, u) \, d\Gamma > 0\},$$

$$\mu_- = \sup_{u \in U} \frac{\|u\|_b^p}{p \int_{\Gamma} H(x, u) \, d\Gamma}, \quad \mu^+ = \inf_{u \in V} \frac{\|u\|_b^p}{p \int_{\Gamma} H(x, u) \, d\Gamma}. \quad (3.38)$$

If  $U = \emptyset$  (resp.  $V = \emptyset$ ), then we set  $\mu_- = -\infty$  (resp.  $\mu^+ = +\infty$ ).

Proceeding in the same manner as we did for proving (3.34), we can show that if we take  $\mu_- < \mu < \mu^+$ , then there exists  $c > 0$  such that

$$\frac{1}{p} \|u\|_b^p - \mu \int_{\Gamma} H(x, u) d\Gamma \geq c \|u\|_b^p \quad \text{for all } u \in X. \quad (3.39)$$

In what follows we employ the following elementary inequality

$$k|u|^\beta - \chi|u|^\gamma \leq C_{\beta, \gamma} k \left( \frac{k}{\chi} \right)^{\frac{\beta}{\gamma-\beta}}, \quad \forall u \in \mathbb{R}, \forall \chi, k \in (0, \infty), \forall 0 < \beta < \gamma. \quad (3.40)$$

**Proposition 3.1** *If  $\mu_- < \mu < \mu^+$ , then the functional  $\Phi$  is coercive.*

*Proof* By virtue of (3.40) we have

$$\begin{aligned} \int_{\Omega} \left( \frac{\lambda}{q} |u|^q w_1 - \frac{\chi}{2r} |u|^r \right) dx &\leq C_{r, q} \int_{\Omega} \lambda w_1 \left( \frac{\lambda w_1}{\chi} \right)^{q/(r-q)} dx \\ &= C_{r, q} \lambda^{r/(r-q)} \int_{\Omega} \frac{w_1^{r/(r-q)}}{h^{q/(r-q)}} dx. \end{aligned}$$

Using (3.39) it follows that

$$\begin{aligned} \Phi(u) &= \frac{1}{p} \|u\|_b^p - \mu \int_{\Gamma} H(x, u) d\Gamma - \int_{\Omega} \left( \frac{\lambda}{q} |u|^q w_1 - \frac{\chi}{2r} |u|^r \right) dx \\ &\quad + \frac{1}{2r} \int_{\Omega} \chi |u|^r dx \geq c \|u\|_b^p + \frac{1}{2r} \int_{\Omega} \chi |u|^r dx - C_1 \end{aligned}$$

and the coercivity of  $\Phi$  follows.  $\square$

**Proposition 3.2** *Suppose  $\mu_- < \mu < \mu^+$  and let  $\{u_n\}$  be a sequence in  $X$  such that  $\Phi(u_n)$  is bounded. Then there exists a subsequence of  $\{u_n\}$ , relabeled again by  $\{u_n\}$ , such that  $u_n \rightharpoonup u_0$  in  $X$  and*

$$\Phi(u_0) \leq \liminf_{n \rightarrow \infty} \Phi(u_n).$$

*Proof* Since  $\Phi$  is coercive in  $X$  we see that the boundedness of  $\{\Phi(u_n)\}$  implies that  $\|u_n\|_b$  and  $\int_{\Omega} \chi |u_n|^r dx$  are bounded. From Theorem 3.1 we know that the embedding  $E \hookrightarrow L^q(\Omega; w_1)$  is compact and using the fact that  $\{u_n\}$  is bounded in  $E$  we may assume that  $u_n \rightharpoonup u_0$  in  $E$  and  $u_n \rightarrow u_0$  in  $L^q(\Omega; w_1)$ .

Set  $F(x, u) = \frac{\lambda}{q}|u|^q w_1 - \frac{1}{r}h|u|^r$  and  $f(x, u) = F_u(x, u)$ . A simple computation yields

$$f_u(x, u) = (q-1)\lambda|u|^{q-2}w_1 - (r-1)\chi|u|^{r-2} \leq C_{r,q}\lambda w_1 \left(\frac{\lambda w_1}{\chi}\right)^{(q-2)/(r-q)}, \quad (3.41)$$

where the last inequality follows from (3.40). We obtain

$$\begin{aligned} \Phi(u_0) - \Phi(u_n) &= \frac{1}{p} \int_{\Omega} a(x)|\nabla u_0|^p dx + \frac{1}{p} \int_{\Gamma} b(x)|u_0|^p d\Gamma \\ &\quad - \frac{1}{p} \int_{\Omega} a(x)|\nabla u_n|^p dx - \frac{1}{p} \int_{\Gamma} b(x)|u_n|^p d\Gamma - \mu \int_{\Gamma} G(x, u_0) d\Gamma \\ &\quad + \mu \int_{\Gamma} H(x, u_n) d\Gamma + \int_{\Omega} (F(x, u_n) - F(x, u_0)) dx = \|u_0\|_b^p - \|u_n\|_b^p \\ &\quad + \mu \left( \int_{\Gamma} H(x, u_n) d\Gamma - \int_{\Gamma} H(x, u_0) d\Gamma \right) \\ &\quad + \int_{\Omega} \left( \int_0^1 \int_0^s f_u(x, u_0 + t(u_n - u_0)) dt ds \right) (u_n - u_0)^2 dx \\ &\leq \|u_0\|_b^p - \|u_n\|_b^p + \mu \left( \int_{\Gamma} H(x, u_n) d\Gamma - \int_{\Gamma} H(x, u_0) d\Gamma \right) \\ &\quad + C_2 \int_{\Omega} (u_n - u_0)^2 \frac{w_1^{(r-2)/(r-q)}}{\chi^{(q-2)/(r-q)}} dx, \end{aligned}$$

where  $C_2 = \frac{1}{2}C_{r,q}\lambda^{(r-2)/(r-q)}$ .

We show that the last integral tends to 0 as  $n \rightarrow \infty$ . Indeed, applying Hölder's inequality we obtain

$$\begin{aligned} \int_{\Omega} (u_n - u_0)^2 \frac{w_1^{(r-2)/(r-q)}}{\chi^{(q-2)/(r-q)}} dx &\leq \left( \int_{\Omega} \frac{w_1^{r/(r-q)}}{\chi^{q/(r-q)}} dx \right)^{(q-2)/q} \\ &\quad \cdot \left( \int_{\Omega} w_1 |u_n - u_0|^q dx \right)^{2/q}. \end{aligned}$$

Since  $u_n \rightarrow u_0$  in  $L^q(\Omega; w_1)$  we have

$$\lim_{n \rightarrow \infty} \int_{\Omega} (u_n - u_0)^2 \frac{w_1^{(r-2)/(r-q)}}{\chi^{(q-2)/(r-q)}} dx = 0. \quad (3.42)$$

The compactness of the trace operator  $E \rightarrow L^m(\Gamma; w_2)$  and the continuity of the Nemytskii operator  $N_H : L^m(\Gamma; w_2) \rightarrow L^1(\Gamma)$  imply  $N_H(u_n) \rightarrow N_H(u_0)$  in  $L^1(\Gamma)$ , that is,  $\int_{\Gamma} |N_H(u_n) - N_H(u_0)| d\Gamma \rightarrow 0$

as  $n \rightarrow \infty$ . It follows that

$$\lim_{n \rightarrow \infty} \int_{\Gamma} H(x, u_n) d\Gamma = \int_{\Gamma} H(x, u_0) d\Gamma. \quad (3.43)$$

Since the norm in  $E$  is lower semicontinuous with respect to the weak topology our conclusion follows from (3.42) and (3.43).  $\square$

**Proposition 3.3** *If  $\mu_* < \mu < \mu^*$  and  $u$  is a solution of problem  $(P'_{\lambda, \mu})$ , then*

$$C_0 \|u\|_b^p + \frac{r-q}{r} \int_{\Omega} \chi |u|^r dx \leq \frac{r-q}{r} \lambda^{r/(r-q)} \int_{\Omega} \frac{w_1^{r/(r-q)}}{\chi^{q/(r-q)}} dx$$

and

$$\|u\|_b \geq K \lambda^{-1/(q-p)},$$

where  $K > 0$  is a constant independent of  $u$ .

*Proof* If  $u$  is a solution of  $(P'_{\lambda, \mu})$ , then

$$\begin{aligned} \|u\|_b^p - \mu \int_{\Gamma} h(x, u) u d\Gamma + \int_{\Omega} \chi |u|^r dx &= \lambda \int_{\Omega} w_1 |u|^q dx \leq \\ &\leq \frac{r-q}{r} \lambda^{r/(r-q)} \int_{\Omega} \frac{w_1^{r/(r-q)}}{\chi^{q/(r-q)}} dx + \frac{q}{r} \int_{\Omega} \chi |u|^r dx. \end{aligned}$$

Using (3.34) we obtain the first part of the assertion.

From Theorem 3.1 we have that there exists  $C_q > 0$  such that

$$\|u\|_{L^q(\Omega; w_1)}^q \leq C_q \|u\|_b^q, \quad \text{for all } u \in E.$$

The above inequality and (3.34) imply

$$\|u\|_b \geq C_0^{1/(q-p)} C_q^{-1/(q-p)} \lambda^{-1/(q-p)}.$$

By taking  $K = C_0^{1/(q-p)} C_q^{-1/(q-p)}$  the second assertion follows.  $\square$

**Theorem 3.5** *Assume that hypotheses (3.27), (3.28), (3.29), (3.30), (h1) and (h2) hold. Set  $\underline{\mu} = \max\{\mu_*, \mu_-\}$ ,  $\bar{\mu} = \min\{\mu^*, \mu^+\}$  and  $J = (\underline{\mu}, \bar{\mu})$ . Then there exists  $\lambda_0 > 0$  such that the following statements hold:*

(i) *problem  $(P'_{\lambda, \mu})$  admits a nontrivial solution, for any  $\lambda \geq \lambda_0$  and every  $\mu \in J$ ;*

(ii) *problem  $(P'_{\lambda, \mu})$  does not have any nontrivial solution, provided that  $0 < \lambda < \lambda_0$  and  $\mu \in J$ .*

*Proof* According to Propositions 3.1 and 3.2,  $\Phi$  is coercive and lower semicontinuous. Therefore, there exists  $\tilde{u} \in X$  such that  $\Phi(\tilde{u}) = \inf_X \Phi(u)$ .

To ensure that  $\tilde{u} \neq 0$ , we shall prove that  $\inf_X \Phi < 0$ . Set

$$\tilde{\lambda} := \inf \left\{ \frac{q}{p} \|u\|_b^p - q\mu \int_{\Gamma} H(x, u) d\Gamma + \frac{q}{r} \int_{\Omega} \chi |u|^r dx : u \in X, \int_{\Omega} w_1 |u|^q dx = 1 \right\}.$$

First we check that  $\tilde{\lambda} > 0$ . For this aim we consider the constrained minimization problem

$$M := \inf \left\{ \int_{\Omega} a(x) |\nabla u|^p dx + \int_{\Gamma} b(x) |u|^p d\Gamma : u \in E, \int_{\Omega} w_1 |u|^q dx = 1 \right\}.$$

Clearly,  $M > 0$ . Since  $X$  is embedded in  $E$ , we have

$$\int_{\Omega} a(x) |\nabla u|^p dx + \int_{\Gamma} b(x) |u|^p d\Gamma \geq M$$

for all  $u \in X$  with  $\int_{\Omega} w_1 |u|^q dx = 1$ . Now, applying the Hölder inequality we find

$$\begin{aligned} 1 &= \int_{\Omega} w_1 |u|^q dx = \int_{\Omega} \frac{w_1}{\chi^{q/r}} \chi^{q/r} |u|^q dx \\ &\leq \left( \int_{\Omega} \frac{w_1^{r/(r-q)}}{\chi^{q/(r-q)}} dx \right)^{(r-q)/r} \cdot \left( \int_{\Omega} \chi |u|^r dx \right)^{q/r}. \end{aligned} \quad (3.44)$$

Relation (3.39) implies

$$\frac{q}{p} \|u\|_b^p - q\mu \int_{\Gamma} H(x, u) d\Gamma \geq qc \|u\|_b^p.$$

By virtue of (3.44) we have

$$\begin{aligned} \frac{q}{p} \|u\|_b^p - q\mu \int_{\Gamma} H(x, u) d\Gamma + \frac{q}{r} \int_{\Omega} \chi |u|^r dx &\geq qc \|u\|_b^p + \frac{q}{r} \int_{\Omega} \chi |u|^r dx \geq \\ &\geq qcM + \frac{q}{r} \left( \int_{\Omega} \frac{w_1^{r/(r-q)}}{\chi^{q/(r-q)}} dx \right)^{-(r-q)/q} \end{aligned}$$

for all  $u \in X$  with  $\int_{\Omega} w_1 |u|^q dx = 1$ . Then,

$$\tilde{\lambda} \geq qcM + \frac{q}{r} \left( \int_{\Omega} \frac{w_1^{r/(r-q)}}{\chi^{q/(r-q)}} dx \right)^{-(r-q)/q}$$

and our claim follows.

Let  $\lambda > \tilde{\lambda}$ . Then, there exists a function  $u \in X$  with  $\int_{\Omega} w_1 |u|^q dx = 1$  such that

$$\lambda > \frac{q}{p} \|u\|_b^p - q\mu \int_{\Gamma} H(x, u) d\Gamma + \frac{q}{r} \int_{\Omega} \chi |u|^r dx.$$

This can be rewritten as

$$\Phi(u) = \frac{1}{p} \|u\|_b^p - \mu \int_{\Gamma} H(x, u) d\Gamma + \frac{1}{r} \int_{\Omega} \chi |u|^r dx - \frac{\lambda}{q} \int_{\Omega} w_1 |u|^q dx < 0$$

and consequently  $\inf_{u \in X} \Phi(u) < 0$ . By Propositions 3.1 and 3.2 it follows that problem  $(P'_{\lambda, \mu})$  has a solution.

We set

$$\lambda_0 = \inf\{\lambda > 0 : (P'_{\lambda, \mu}) \text{ admits a solution}\}.$$

Suppose  $\lambda_0 = 0$ . Then by taking  $\lambda_1 \in (0, \lambda^*)$  (where  $\lambda^*$  is given by Theorem 3.4), we have that there exists  $\bar{\lambda}$  such that problem  $(P'_{\bar{\lambda}, \mu})$  admits a solution. But this is a contradiction, according to Theorem 3.4. Consequently,  $\lambda_0 > 0$ .

We now show that for each  $\lambda > \lambda_0$  problem  $(P'_{\lambda, \mu})$  admits a solution. Indeed, for every  $\lambda > \lambda_0$  there exists  $\rho \in (\lambda_0, \lambda)$  such that problem  $(P'_{\rho, \mu})$  has a solution  $u_{\rho}$  which is a subsolution of problem  $(P'_{\lambda, \mu})$ . We consider the variational problem

$$\inf\{\Phi(u) : u \in X \text{ and } u \geq u_{\rho}\}.$$

By Propositions 3.1 and 3.2 this problem admits a solution  $\bar{u}$ . This minimizer  $\bar{u}$  is a solution of problem  $(P'_{\lambda,\mu})$ . Since the hypothesis  $h(x, s) + h(x, -s) \geq 0$  for a.e.  $x \in \Gamma$  and for all  $s \in \mathbb{R}$  implies that  $H(x, |\bar{u}|) \geq H(x, \bar{u})$  (that is,  $\Phi(|\bar{u}|) \leq \Phi(\bar{u})$ ), we may assume that  $\bar{u} \geq 0$  on  $\Omega$ . It remains to show that problem  $(P'_{\lambda_0,\mu})$  has also a solution. Let  $\lambda_n \rightarrow \lambda_0$  and  $\lambda_n > \lambda_0$  for each  $n$ . Problem  $(P'_{\lambda_n,\mu})$  has a solution  $u_n$  for each  $n$ . By Proposition 3.3 the sequence  $\{u_n\}$  is bounded in  $X$ . Therefore we may assume that  $u_n \rightharpoonup u_0$  in  $X$  and  $u_n \rightarrow u_0$  in  $L^q(\Omega; w_1)$ . We have that  $u_0$  is a solution of  $(P'_{\lambda_0,\mu})$ . Since  $u_n$  and  $u_0$  are solutions of  $(P'_{\lambda_n,\mu})$  and  $(P'_{\lambda_0,\mu})$ , respectively, we have

$$\begin{aligned} & \int_{\Omega} a(x)(|\nabla u_n|^{p-2}\nabla u_n - |\nabla u_0|^{p-2}\nabla u_0)(\nabla u_n - \nabla u_0) dx + \\ & \int_{\Gamma} b(x)(|u_n|^{p-2}u_n - |u_0|^{p-2}u_0)(u_n - u_0) d\Gamma + \\ & \int_{\Omega} \chi(|u_n|^{r-2}u_n - |u_0|^{r-2}u_0)(u_n - u_0) dx = \\ & \lambda_n \int_{\Omega} w_1(|u_n|^{q-2}u_n - |u_0|^{q-2}u_0)(u_n - u_0) dx + \\ & (\lambda_n - \lambda_0) \int_{\Omega} w_1|u_0|^{q-2}u_0(u_n - u_0) dx + \\ & \mu \int_{\Gamma} (h(x, u_n) - h(x, u_0))(u_n - u_0) d\Gamma = J_{1,n} + J_{2,n} + J_{3,n}, \end{aligned}$$

where

$$\begin{aligned} J_{1,n} &= \lambda_n \int_{\Omega} w_1(|u_n|^{q-2}u_n - |u_0|^{q-2}u_0)(u_n - u_0) dx, \\ J_{2,n} &= (\lambda_n - \lambda_0) \int_{\Omega} w_1|u_0|^{q-2}u_0(u_n - u_0) dx, \\ J_{3,n} &= \mu \int_{\Gamma} (h(x, u_n) - h(x, u_0))(u_n - u_0) d\Gamma. \end{aligned}$$

We have

$$|J_{1,n}| \leq \sup_{n \geq 1} \lambda_n \left( \int_{\Omega} w_1|u_n|^{q-1}|u_n - u_0| dx + \int_{\Omega} w_1|u_0|^{q-1}|u_n - u_0| dx \right)$$

and it follows from Hölder's inequality that

$$|J_{1,n}| \leq \sup_{n \geq 1} \lambda_n \left[ \left( \int_{\Omega} w_1 |u_n|^q dx \right)^{(q-1)/q} \cdot \left( \int_{\Omega} w_1 |u_n - u_0|^q dx \right)^{1/q} + \left( \int_{\Omega} w_1 |u_0|^q dx \right)^{(q-1)/q} \cdot \left( \int_{\Omega} w_1 |u_n - u_0|^q dx \right)^{1/q} \right].$$

We easily observe that  $J_{1,n} \rightarrow 0$  as  $n \rightarrow \infty$ .

From the estimate

$$|J_{2,n}| \leq |\lambda_n - \lambda_0| \left( \int_{\Omega} w_1 |u_0|^q dx \right)^{(q-1)/q} \cdot \left( \int_{\Omega} w_1 |u_n - u_0|^q dx \right)^{1/q}$$

we obtain that  $J_{2,n} \rightarrow 0$  as  $n \rightarrow \infty$ .

Using the compactness of the trace operator  $E \rightarrow L^m(\Gamma; w_2)$ , the continuity of the Nemytskii operator

$$N_h : L^m(\Gamma; w_2) \rightarrow L^{m/(m-1)}(\Gamma; w_2^{1/(1-m)})$$

and the estimate

$$\begin{aligned} & \int_{\Gamma} |h(x, u_n) - h(x, u_0)| \cdot |u_n - u_0| d\Gamma \leq \\ & \left( \int_{\Gamma} |h(x, u_n) - h(x, u_0)|^{m/(m-1)} w_2^{1/(1-m)} d\Gamma \right)^{(m-1)/m} \cdot \\ & \left( \int_{\Gamma} w_2 |u_n - u_0|^m d\Gamma \right)^{1/m}, \end{aligned}$$

we see that  $J_{3,n} \rightarrow 0$  as  $n \rightarrow \infty$ . Therefore

$$\begin{aligned} & \lim_{n \rightarrow \infty} \left( \int_{\Omega} a(x) (|\nabla u_n|^{p-2} \nabla u_n - |\nabla u_0|^{p-2} \nabla u_0) (\nabla u_n - \nabla u_0) dx + \right. \\ & \left. \int_{\Gamma} b(x) (|u_n|^{p-2} u_n - |u_0|^{p-2} u_0) (u_n - u_0) d\Gamma \right) = 0. \end{aligned}$$

Applying the inequality

$$|\xi - \zeta|^p \leq C(|\xi|^{p-2} \xi - |\zeta|^{p-2} \zeta)(\xi - \zeta), \quad \text{for all } \xi, \zeta \in \mathbb{R}^N$$

we find

$$\begin{aligned} \|u_n - u_0\|_b^p &= \int_{\Omega} a(x) |\nabla u_n - \nabla u_0|^p dx + \int_{\Gamma} b(x) |u_n - u_0|^p dx \leq \\ &\leq C \left( \int_{\Omega} a(x) (|\nabla u_n|^{p-2} \nabla u_n - |\nabla u_0|^{p-2} \nabla u_0) (\nabla u_n - \nabla u_0) dx + \right. \\ &\quad \left. + \int_{\Gamma} b(x) (|u_n|^{p-2} u_n - |u_0|^{p-2} u_0) (u_n - u_0) d\Gamma \right) \rightarrow 0 \text{ as } n \rightarrow \infty, \end{aligned}$$

which shows that  $\|u_n\|_b \rightarrow \|u_0\|_b$ . By Proposition 3.3 we have  $u_0 \neq 0$ . This concludes our proof.  $\square$

### 3.4 Sublinear case

In this section we suppose that the conditions (3.27), (3.28) and (3.29) are fulfilled. For  $\lambda, \mu > 0$ , we study the following double eigenvalue problem:

$$(P''_{\lambda, \mu}) \quad \begin{cases} -\operatorname{div}(a(x) |\nabla u|^{p-2} \nabla u) = \lambda f(x, u(x)) \text{ in } \Omega, \\ a(x) |\nabla u|^{p-2} \nabla u \cdot \mathbf{n} + b(x) |u|^{p-2} u = \mu g(x, u(x)) \text{ on } \Gamma, \\ u \neq 0 \text{ in } \Omega. \end{cases}$$

We consider the following assumptions:

(F1) Let  $f : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$  be a Carathéodory function such that  $f(\cdot, 0) = 0$  and

$$|f(x, s)| \leq f_0(x) + f_1(x) |s|^{r-1},$$

where  $p < r < \frac{pN}{N-p}$ , and  $f_0, f_1$  are measurable functions which satisfy

$$0 < f_0(x) \leq C_f w_1(x), \text{ and } 0 \leq f_1(x) \leq C_f w_1(x) \text{ a.e. } x \in \Omega,$$

$$f_0 \in L^{\frac{r}{r-1}}(\Omega; w_1^{\frac{1}{1-r}});$$

(F2)

$$\lim_{s \rightarrow 0} \frac{f(x, s)}{f_0(x) |s|^{p-1}} = 0, \text{ uniformly in } x \in \Omega;$$

- (F3)  $\limsup_{s \rightarrow +\infty} \frac{1}{f_0(x)|s^p|} F(x, s) \leq 0$  uniformly for all  $x \in \Omega$ ,  $\max_{|s| \leq M} F(\cdot, s) \in L^1(\Omega)$  for all  $M > 0$ , where  $F$  denotes the primitive function of  $f$  with respect to the second variable, i.e.  $F(x, u) = \int_0^u f(x, s) ds$ ;
- (F4) there exist  $x_0 \in \Omega$ ,  $R_0 > 0$  and  $s_0 \in \mathbb{R}$  such that  $\min_{|x-x_0| < R_0} F(x, s_0) > 0$ .
- (G1) Let  $g : \Gamma \times \mathbb{R} \rightarrow \mathbb{R}$  be a Carathéodory function such that  $g(\cdot, 0) = 0$  and

$$|g(x, s)| \leq g_0(x) + g_1(x)|s|^{m-1},$$

where  $p \leq m < p \cdot \frac{N-1}{N-p}$ , and  $g_0, g_1$  are measurable functions satisfying

$$0 < g_0(x) \leq C_g w_2(x) \text{ and } 0 \leq g_1(x) \leq C_g w_2(x), \text{ a.e. } x \in \Gamma,$$

$$g_0 \in L^{\frac{a}{a-1}}(\Gamma; w_2^{\frac{1}{1-a}});$$

(G2)

$$\lim_{s \rightarrow 0} \frac{g(x, s)}{g_0(x)|s|^{p-1}} = 0, \text{ uniformly in } x \in \Gamma;$$

- (G3)  $\limsup_{s \rightarrow +\infty} \frac{1}{g_0(x)|s^p|} G(x, s) < +\infty$  uniformly for all  $x \in \Gamma$ ,  $\max_{|s| \leq M} G(\cdot, s) \in L^1(\Gamma)$  for all  $M > 0$ , where  $G$  is the primitive function of  $g$  with respect to the second variable, i.e.  $G(x, u) = \int_0^u g(x, s) ds$ .

The functions  $w_1$  and  $w_2$  are defined as in the previous section, that is,

$$w_i = (1 + |x|)^{\alpha_i}, \quad i = 1, 2.$$

The energy functional corresponding to  $(P''_{\lambda, \mu})$  is given by  $\mathcal{E}_{\lambda, \mu} : E \rightarrow \mathbb{R}$ ,

$$\mathcal{E}_{\lambda, \mu}(u) = \frac{1}{p} \int_{\Omega} a(x) |\nabla u(x)|^p dx + \frac{1}{p} \int_{\Gamma} b(x) |u(x)|^p d\Gamma - \lambda J_F(u) - \mu J_G(u),$$

where  $J_F, J_G : E \rightarrow \mathbb{R}$  are defined by

$$J_F(u) = \int_{\Omega} F(x, u(x)) dx \quad \text{and} \quad J_G(u) = \int_{\Gamma} G(x, u(x)) d\Gamma.$$

The first multiplicity result for problem  $(P''_{\lambda, \mu})$  is the following:

**Theorem 3.6** *Let  $f : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$  be a function satisfying the conditions (F1)-(F4). Then, there exists a non-degenerate compact interval  $[a, b] \subset ]0, +\infty[$  with the following properties:*

- 1° *there exists a number  $\sigma_0 > 0$  such that for every  $\lambda \in [a, b]$  and for every function  $g : \Gamma \times \mathbb{R} \rightarrow \mathbb{R}$  satisfying the conditions (G1)-(G2), there exists  $\mu_0 > 0$  such that for each  $\mu \in ]0, \mu_0[$ , the functional  $\mathcal{E}_{\lambda, \mu}$  has at least two critical points with norms less than  $\sigma_0$ ;*
- 2° *there exists a number  $\sigma_1 > 0$  such that for every  $\lambda \in [a, b]$  and for every function  $g : \Gamma \times \mathbb{R} \rightarrow \mathbb{R}$  satisfying the conditions (G1)-(G3), there exists  $\mu_1 > 0$  such that for each  $\mu \in ]0, \mu_1[$ , the functional  $\mathcal{E}_{\lambda, \mu}$  has at least three critical points with norms less than  $\sigma_1$ .*

To prove Theorem 3.6 we use Theorem 1.17. Before we prove this result, we need some auxiliary results.

**Lemma 3.5** *We assume that the conditions (F1) and (F2) are satisfied. Then, the functional  $J_F$  is sequentially weakly continuous.*

*Proof* First we observe that, from assumption (F1) and (F2), for every  $\varepsilon > 0$ , there exists  $C_\varepsilon$  such that

$$|F(x, u)| \leq \varepsilon f_0(x)|u(x)|^p + C_\varepsilon(f_0(x) + f_1(x))|u(x)|^r. \quad (3.45)$$

Now we prove our assertion, arguing by contradiction: Let  $\{u_n\}$  be a sequence in  $E$  weakly convergent to  $u \in E$ , and let  $d > 0$  be such that

$$|J_F(u_n) - J_F(u)| \geq d \quad \text{for all } n \in \mathbb{N}.$$

Without loss of generality, we can assume that there is a positive constant  $M$  such that

$$\|u\|_b \leq M, \|u_n\|_b \leq M, \|u_n - u\|_b \leq M, \quad \text{for all } n \in \mathbb{N}.$$

Because the embedding  $E \hookrightarrow L^r(\Omega; w_1)$  is compact, follows  $\|u_n - u\|_{r, w_1} \rightarrow$

0. Using (3.45), Theorem 3.1 and the Hölder inequality we have

$$\begin{aligned}
& |J_F(u_n) - J_F(u)| \leq \int_{\Omega} |F(x, u_n(x)) - F(x, u(x))| dx \\
& \leq \varepsilon \hat{c} \int_{\Omega} f_0(x) |u_n(x) - u(x)| (|u_n(x)|^{p-1} + |u(x)|^{p-1}) dx \\
& \quad + \hat{c} C_{\varepsilon} \int_{\Omega} (f_0(x) + f_1(x)) |u_n(x) - u(x)| (|u_n(x)|^{r-1} + |u(x)|^{r-1}) dx \\
& \leq \varepsilon \hat{c} C_f \int_{\Omega} w_1(x) |u_n(x) - u(x)| (|u_n(x)|^{p-1} + |u(x)|^{p-1}) dx \\
& \quad + 2\hat{c} C_{\varepsilon} C_f \int_{\Omega} w_1(x) |u_n(x) - u(x)| (|u_n(x)|^{r-1} + |u(x)|^{r-1}) dx \\
& \leq \varepsilon \hat{c} C_f \|u_n - u\|_{p, w_1} (\|u_n\|_{p, w_1}^{\frac{p}{p'}} + \|u\|_{p, w_1}^{\frac{p}{p'}}) \\
& \quad + 2\hat{c} C_{\varepsilon} C_f \|u_n - u\|_{r, w_1} (\|u_n\|_{r, w_1}^{\frac{r}{r'}} + \|u\|_{r, w_1}^{\frac{r}{r'}}),
\end{aligned}$$

where  $\hat{c} > 0$  is a constant,  $\frac{1}{p} + \frac{1}{p'} = 1$  and  $\frac{1}{r} + \frac{1}{r'} = 1$ . By the embedding results from Theorem 3.1 it follows that

$$d \leq |J_F(u_n) - J_F(u)| \leq 2\varepsilon \hat{c} C_f C_{p, w_1}^p M^p + 4\hat{c} C_{\varepsilon} C_f C_{r, w_1}^{\frac{r}{r'}} M^{\frac{r}{r'}} \|u_n - u\|_{r, w_1}.$$

Therefore, if  $\varepsilon > 0$  is sufficiently small and  $n \in \mathbb{N}$  is large enough, we have

$$d \leq |J_F(u_n) - J_F(u)| < d,$$

which is a contradiction.  $\square$

**Remark 3.1** *Note that a similar result holds for the functional  $J_G$ , i.e. if the conditions (G1), (G2) are fulfilled, then the functional  $J_G$  is sequentially weakly continuous.*

**Lemma 3.6** *There exists  $u_0 \in E$  such that  $J_F(u_0) > 0$ .*

*Proof* Let  $R_0 > 0$  and  $s_0 \in \mathbb{R}$  from (F4) and fix  $\varepsilon \in (0, \frac{R_0}{2})$ . We consider the function  $u_{\varepsilon} \in C_0^{\infty}(\Omega)$  such that

$$u_{\varepsilon}(x) = \begin{cases} 0, & \text{if } |x - x_0| \geq R_0 \\ s_0, & \text{if } |x - x_0| \leq R_0 - \varepsilon \end{cases}$$

and  $\|u_{\varepsilon}\|_{\infty} \leq |s_0|$ . Then, we have  $u_{\varepsilon} \in E$ .

We denote by  $\text{Vol}(B_r(x_0))$  the volume of the ball

$$B_r(x_0) = \{x \in \mathbb{R}^N : |x - x_0| \leq r\}.$$

We have

$$\begin{aligned}
J_F(u_\varepsilon) &= \int_{\Omega} F(x, u_\varepsilon) dx = \int_{B_{R_0-\varepsilon}(x_0) \cap \Omega} F(x, u_\varepsilon(x)) dx \\
&+ \int_{(\mathbb{R}^N \setminus B_{R_0}(x_0)) \cap \Omega} F(x, u_\varepsilon(x)) dx \\
&+ \int_{(B_{R_0}(x_0) \setminus B_{R_0-\varepsilon}(x_0)) \cap \Omega} F(x, u_\varepsilon(x)) dx \\
&\geq A_0 \text{Vol}(B_{\frac{R_0}{2}}(x_0) \cap \Omega) \\
&- B_0 [\text{Vol}(B_{R_0}(x_0) \cap \Omega) - \text{Vol}(B_{R_0-\varepsilon}(x_0) \cap \Omega)],
\end{aligned}$$

where

$$A_0 = \min_{|x-x_0| \leq R_0} F(x, s_0) > 0 \quad \text{and} \quad B_0 = \max_{\frac{R_0}{2} \leq |x-x_0| \leq R_0} \max_{|s| \leq |s_0|} F(x, s).$$

If  $\varepsilon$  goes to 0, then  $[\text{Vol}(B_{R_0}(x_0) \cap \Omega) - \text{Vol}(B_{R_0-\varepsilon}(x_0) \cap \Omega)] \rightarrow 0$  as well, so we can choose an  $\varepsilon = \varepsilon_0$ , such that  $J_F(u_{\varepsilon_0}) > 0$ . We take  $u_0 = u_{\varepsilon_0}$  and the proof is complete.  $\square$

**Lemma 3.7** *Suppose that the conditions (F1) and (F3) are satisfied. Then, for every  $\lambda \geq 0$  the functional*

$$u \rightarrow \frac{\|u\|_b^p}{p} - \lambda J_F(u)$$

*is coercive on  $E$ .*

*Proof* If  $\lambda = 0$ , then statement is trivial. Let  $\lambda > 0$  and  $a \in \left] 0, \frac{1}{\lambda p C_{p,w_1}^p} \right[$ . There exists a positive function  $k_a \in L^1(\Omega; w_1)$  such that

$$F(x, s) \leq a f_0(x) |s|^p + k_a(x) w_1(x), \quad \text{for all } (x, s) \in \Omega \times \mathbb{R}.$$

Then, it follows that

$$\begin{aligned}
\frac{\|u\|_b^p}{p} - \lambda J_F(u) &\geq \frac{\|u\|_b^p}{p} - \lambda a \int_{\Omega} w_1(x) |u(x)|^p dx - \lambda \int_{\Omega} k_a(x) w_1(x) dx \geq \\
&\geq \|u\|_b^p \left( \frac{1}{p} - \lambda a C_{p,w_1}^p \right) - \lambda \|k_a\|_{1,w_1},
\end{aligned}$$

which converges to  $\infty$  as  $\|u\|_b \rightarrow \infty$ .  $\square$

*Proof of Theorem 3.6:* We define the function  $h : ]0, +\infty[ \rightarrow \mathbb{R}$  by

$$h(t) = \sup \left\{ J_F(u) : \frac{\|u\|_b^p}{p} \leq t \right\}, \quad \text{for all } t > 0.$$

By using (3.45) it follows that

$$0 \leq h(t) \leq \varepsilon p C_f C_{p,w_1}^p t + 2p^{\frac{r}{p}} C_\varepsilon C_f C_{r,w_1}^r t^{\frac{r}{p}} \quad \text{for all } t > 0;$$

since  $p < r$ , this implies

$$\lim_{t \rightarrow 0^+} \frac{h(t)}{t} = 0.$$

By (F4) it is clear that  $u_0 \neq 0$  (since  $J_F(0) = 0$ ). Therefore, due to the convergence relation above and Lemma 3.6 it is possible to choose a real number  $t_0$  such that  $0 < t_0 < \frac{\|u_0\|_b^p}{p}$  and

$$\frac{h(t_0)}{t_0} < \left[ \frac{\|u_0\|_b^p}{p} \right]^{-1} \cdot J_F(u_0).$$

We choose  $\rho_0 > 0$  such that

$$h(t_0) < \rho_0 < \left[ \frac{\|u_0\|_b^p}{p} \right]^{-1} \cdot J_F(u_0) t_0. \quad (3.46)$$

In particular, we get  $\rho_0 < J_F(u_0)$ .

Now, we are going to apply Theorem 1.17 to the space  $E$ , the interval  $\Lambda = ]0, +\infty[$  and the function  $\varphi : E \times \Lambda \rightarrow \mathbb{R}$  defined by

$$\varphi(u, \lambda) = \frac{\|u\|_b^p}{p} + \lambda(\rho_0 - J_F(u)) \quad \text{for all } (u, \lambda) \in E \times \Lambda$$

and  $\Upsilon : E \rightarrow \mathbb{R}$  by

$$\Upsilon(u) = -J_G(u) \quad \text{for all } u \in E.$$

Let us fix  $\lambda \in ]0, +\infty[$ : from Lemma 3.7 it follows that the functional  $\varphi(\cdot, \lambda)$  is coercive; moreover,  $\varphi(\cdot, \lambda)$  is the sum of  $u \mapsto \frac{\|u\|_b^p}{p}$ , which is sequentially weakly l.s.c., and of  $u \mapsto \lambda(\rho_0 - J_F(u))$ , which is sequentially weakly continuous (see Lemma 3.5).

We prove now that  $\varphi$  satisfies the minimax inequality from Theorem 1.17. The function

$$\lambda \mapsto \inf_{u \in E} \varphi(u, \lambda)$$

is upper semicontinuous on  $\Lambda$ . Since

$$\inf_{u \in E} \varphi(u, \lambda) \leq \varphi(u_0, \lambda) = \frac{\|u_0\|_b^p}{p} + \lambda(\rho_0 - J_F(u_0))$$

and  $\rho_0 < J_F(u_0)$ , it follows that

$$\lim_{\lambda \rightarrow +\infty} \inf_{u \in E} \varphi(u, \lambda) = -\infty.$$

Thus we can find  $\bar{\lambda} \in \Lambda$  such that

$$\beta_1 = \sup_{\lambda \in \Lambda} \inf_{u \in E} \varphi(u, \lambda) = \inf_{u \in E} \varphi(u, \bar{\lambda}).$$

In order to prove that  $\beta_1 < t_0$ , we distinguish two cases:

I. If  $0 \leq \bar{\lambda} < \frac{t_0}{\rho_0}$ , we have

$$\beta_1 \leq \varphi(0, \bar{\lambda}) = \bar{\lambda} \rho_0 < t_0.$$

II. If  $\bar{\lambda} \geq \frac{t_0}{\rho_0}$ , then we use  $\rho_0 < J_F(u_0)$  and the inequality (3.46) to get

$$\beta_1 \leq \varphi(u_0, \bar{\lambda}) \leq \frac{\|u_0\|_b^p}{p} + \frac{t_0}{\rho_0}(\rho_0 - J_F(u_0)) < t_0.$$

Let us focus next on the right hand side of the minimax inequality from Theorem 1.17. Clearly,

$$\beta_2 = \inf_{u \in E} \sup_{\lambda \in \Lambda} \varphi(u, \lambda) = \inf \left\{ \frac{\|u\|_b^p}{p} : J_F(u) \geq \rho_0 \right\}.$$

On the other hand, by applying again (3.46) we easily get

$$t_0 \leq \inf \left\{ \frac{\|u\|_b^p}{p} : J_F(u) \geq \rho_0 \right\}.$$

We conclude that

$$\beta_1 < t_0 \leq \beta_2,$$

i.e., the minimax inequality from Theorem 1.17 holds .

Thus, we can apply Theorem 1.17. Fix  $\delta > \beta_1$ , and for every  $\lambda \in \Lambda$  denote

$$S_\lambda = \{u \in E : \varphi(u, \lambda) < \delta\}.$$

There exists a non-empty open set  $\Lambda_0 \subset ]0, +\infty[$  with the following property: for every  $\lambda \in \Lambda_0$  and every sequentially weakly l.s.c.  $\Upsilon : E \rightarrow \mathbb{R}$ , there exists  $\mu_0 > 0$ , such that for each  $\mu \in ]0, \mu_0[$ , the functional

$$u \rightarrow \varphi(u, \lambda) + \mu \Upsilon(u)$$

has at least two local minima lying in the set  $S_\lambda$ . Let  $[a, b] \subset \Lambda_0$  be a non-degenerate compact interval.

We prove now the two assertions of our theorem.

1° Let  $\lambda \in [a, b]$  be a real number, and let  $g : \Gamma \times \mathbb{R} \rightarrow \mathbb{R}$  satisfying the conditions (G1)-(G2), and let  $\Upsilon = -J_G$ . Then, by Remark 3.1,  $\Upsilon$  is sequentially weakly continuous. From what stated above, there exists  $\mu_0 > 0$  such that for all  $\mu \in ]0, \mu_0[$  the functional  $\mathcal{E}_{\lambda, \mu}$  admits at least two local minima  $u_{\lambda, \mu}^1, u_{\lambda, \mu}^2 \in S_\lambda$ , therefore these are critical points of  $\mathcal{E}_{\lambda, \mu}$ .

Observe that

$$S := \bigcup_{\lambda \in [a, b]} S_\lambda \subseteq S_a \cup S_b.$$

Since  $\varphi(\cdot, \lambda)$  is coercive for all  $\lambda \geq 0$ , the latter sets are bounded, hence  $S$  is bounded as well. Chosen  $\sigma_0 > \sup_{u \in S} \|u\|_b$ , we get

$$\|u_{\lambda, \mu}^1\|_b, \|u_{\lambda, \mu}^2\|_b < \sigma_0.$$

2° Let  $\lambda \in [a, b]$  be a real number, and let  $g : \Gamma \times \mathbb{R} \rightarrow \mathbb{R}$  satisfying the conditions (G1)-(G3): as above, there exists  $\mu_0 > 0$  such that for all  $\mu \in ]0, \mu_0[$  the functional  $\mathcal{E}_{\lambda, \mu}$  has at least two local minima  $u_{\lambda, \mu}^1, u_{\lambda, \mu}^2 \in E$  with norms less than  $\sigma_0$ . To prove the existence of a third critical point for  $\mathcal{E}_{\lambda, \mu}$ , we are going to apply Theorem 1.8. For this, it is enough to prove that the functional  $\mathcal{E}_{\lambda, \mu}$  satisfies the (PS) condition for  $\mu > 0$  small enough. Since (G3) holds, arguing as in Lemma 3.7, it is easy to prove that there exists  $\mu_1 \in ]0, \mu_0[$  such that  $\mathcal{E}_{\lambda, \mu}$  is coercive in  $E$  for all  $\mu \in [0, \mu_1]$ . Let  $\{u_n\}$  be a sequence such that  $\{\mathcal{E}_{\lambda, \mu}(u_n)\}$  is bounded and  $\mathcal{E}'_{\lambda, \mu}(u_n) \rightarrow 0$  holds. The coercivity of  $\mathcal{E}_{\lambda, \mu}$  implies that  $\{u_n\}$  is bounded in  $E$ . Because  $E$  is a reflexive Banach space we can find a subsequence, which we still denote by  $\{u_n\}$ , weakly convergent to a point  $u_* \in E$ . We denote  $I(u) = \frac{1}{p} \|u\|_b^p$ . Then the directional derivative of  $\mathcal{E}_{\lambda, \mu}$  in the direction  $h \in E$  is

$$\langle \mathcal{E}'_{\lambda, \mu}(u), h \rangle = \langle I'(u), h \rangle - \lambda \langle J'_F(u), h \rangle - \mu \langle J'_G(u), h \rangle,$$

where

$$\langle I'(u), h \rangle = \int_{\Omega} a(x) |\nabla u|^{p-2} \nabla u(x) \nabla h(x) dx + \int_{\Gamma} b(x) |u(x)|^{p-2} u(x) h(x) d\Gamma,$$

$$\langle J'_F(u), h \rangle = \int_{\Omega} f(x, u(x)) h(x) dx \quad \text{and} \quad \langle J'_G(u), h \rangle = \int_{\Gamma} g(x, u(x)) h(x) d\Gamma.$$

To show that  $u_n \rightarrow u_*$  strongly in  $E$  we use the inequalities (3.17) and (3.18).

In the case  $p \geq 2$  we use (3.17) and we obtain:

$$\begin{aligned}
\|u_n - u_\star\|_b^p &= \int_{\Omega} a(x) |\nabla u_n(x) - \nabla u_\star(x)|^p dx + \int_{\Gamma} b(x) |u_n(x) - u_\star(x)|^p d\Gamma \\
&\leq C \left( \langle I'(u_n), u_n - u_\star \rangle - \langle I'(u_\star), u_n - u_\star \rangle \right) \\
&= C \left( \langle \mathcal{E}'_{\lambda, \mu}(u_n), u_n - u_\star \rangle - \langle \mathcal{E}'_{\lambda, \mu}(u_\star), u_n - u_\star \rangle + \langle \lambda J'_F(u_n) \right. \\
&\quad \left. + \mu J'_G(u_n), u_n - u_\star \rangle - \langle \lambda J'_F(u_\star) + \mu J'_G(u_\star), u_n - u_\star \rangle \right) \\
&\leq C \left( \|\mathcal{E}'_{\lambda, \mu}(u_n)\|_{E'} + \lambda \|J'_F(u_n) - J'_F(u_\star)\|_{E'} \right. \\
&\quad \left. + \mu \|J'_G(u_n) - J'_G(u_\star)\|_{E'} \right) \|u_n - u_\star\|_b - C \langle \mathcal{E}'_{\lambda, \mu}(u_\star), u_n - u_\star \rangle.
\end{aligned}$$

Since  $\mathcal{E}'_{\lambda, \mu}(u_n) \rightarrow 0$  and  $J'_F, J'_G$  are compact, it follows that  $u_n \rightarrow u_\star$  converges strongly in  $E$ .

In the case  $1 < p < 2$ , we use (3.18) and Hölder's inequality to obtain the estimate

$$\|u_n - u_\star\|_b^p \leq \hat{C} \left| \langle I'(u_n), u_n - u_\star \rangle - \langle I'(u_\star), u_n - u_\star \rangle \right| \left( \|u_n\|_b^p + \|u_\star\|_b^p \right)^{(2-p)/p},$$

where  $\hat{C} > 0$  is a positive constant depending on  $p$  and  $C$ .

Thus, the condition (PS) is fulfilled for all  $\mu \in [0, \mu_1]$ . This concludes our proof.

**Corollary 3.1** *Let  $f : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$  be a function satisfying conditions (F1)-(F4). Then, there exists a non-degenerate compact interval  $[a, b] \subset ]0, +\infty[$  with the following properties:*

- I. *there exists a number  $\sigma_0 > 0$  such that for every  $\lambda \in [a, b]$  and for every function  $g : \Gamma \times \mathbb{R} \rightarrow \mathbb{R}$  satisfying conditions (G1)-(G2), there exists  $\mu_0 > 0$  such that for each  $\mu \in ]0, \mu_0[$ , problem  $(P''_{\lambda, \mu})$  has at least one non-trivial solution in  $E$  with norm less than  $\sigma_0$ ;*
- II. *for every  $\lambda \in [a, b]$  and for every function  $g : \Gamma \times \mathbb{R} \rightarrow \mathbb{R}$  satisfying conditions (G1)-(G3), there exists  $\mu_1 > 0$  such that for each  $\mu \in ]0, \mu_1[$ , problem  $(P''_{\lambda, \mu})$  has at least two non-trivial solutions in  $E$ .*

In the last part of this section we consider for  $\lambda > 0$  and  $\mu \in \mathbb{R}$  the following double eigenvalue problem:

$$(P'''_{\lambda,\mu}) \quad \begin{cases} -\operatorname{div}(a(x)|\nabla u|^{p-2}\nabla u) = \lambda f(x, u) \text{ in } \Omega, \\ a(x)|\nabla u|^{p-2}\nabla u \cdot \mathbf{n} + b(x)|u|^{p-2}u = \lambda\mu g(x, u) \text{ on } \Gamma \\ u \neq 0 \text{ in } \Omega, \end{cases}$$

where we use the same notations as in the case of problem  $(P''_{\lambda,\mu})$ .

The energy functional  $\mathcal{F}_{\lambda,\mu} : E \rightarrow \mathbb{R}$  corresponding to  $(P'''_{\lambda,\mu})$  is given by:

$$\mathcal{F}_{\lambda,\mu}(u) = \frac{1}{p} \int_{\Omega} a(x)|\nabla u(x)|^p dx + \frac{1}{p} \int_{\Gamma} b(x)|u(x)|^p d\Gamma - \lambda J_{\mu}(u),$$

where  $J_{\mu}(u) = J_F(u) + \mu J_G(u)$ .

**Lemma 3.8** *Suppose that the conditions (F1), (F3), (G1) and (G3) are satisfied. Then, for every  $\lambda \geq 0$  and  $\mu \in \mathbb{R}$  the functional  $\mathcal{F}_{\lambda,\mu}$  is coercive on  $E$  and satisfies the (PS) condition.*

*Proof* If  $\lambda = 0$ , then the statement is trivial.

Let  $\lambda > 0$ ,  $\mu \in \mathbb{R}$  and  $a, b > 0$  such that  $\lambda a C_f C_{p,w_1}^p + \lambda |\mu| b C_g C_{p,w_2}^p < \frac{1}{p}$ . There exist positive functions  $k_a \in L^1(\Omega; w_1)$ ,  $k_b \in L^1(\Gamma; w_2)$  such that

$$F(x, s) \leq a f_0(x) |s|^p + k_a(x) w_1(x), \quad \text{for all } (x, s) \in \Omega \times \mathbb{R},$$

$$G(x, s) \leq b g_0(x) |s|^p + k_b(x) w_2(x), \quad \text{for all } (x, s) \in \Gamma \times \mathbb{R}.$$

Then, it follows that

$$\begin{aligned} \mathcal{F}_{\lambda,\mu}(u) &= \frac{\|u\|_b^p}{p} - \lambda J_{\mu}(u) \\ &= \frac{\|u\|_b^p}{p} - \lambda \left( \int_{\Omega} F(x, u(x)) dx + \mu \int_{\Gamma} G(x, u(x)) d\Gamma \right) \\ &\geq \frac{\|u\|_b^p}{p} - \lambda a C_f \int_{\Omega} w_1(x) |u(x)|^p dx - \lambda \int_{\Omega} k_a(x) w_1(x) dx \\ &\quad - \lambda |\mu| b C_g \int_{\Gamma} w_2(x) |u(x)|^p d\Gamma - \lambda |\mu| \int_{\Gamma} k_b(x) w_2(x) d\Gamma \\ &\geq \|u\|_b^p \left( \frac{1}{p} - \lambda a C_f C_{p,w_1}^p - \lambda |\mu| b C_g C_{p,w_2}^p \right) \\ &\quad - \lambda \|k_a\|_{1,w_1} - \lambda |\mu| \|k_b\|_{1,w_2}, \end{aligned}$$

which goes to  $\infty$  as  $\|u\|_b \rightarrow \infty$ . We omit to prove that  $\mathcal{F}_{\lambda,\mu}$  satisfies the (PS) condition, since it is likewise to the proof given in Lemma 3.4.  $\square$

**Theorem 3.7** *We suppose that the functions  $f : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$  and  $g : \Gamma \times \mathbb{R} \rightarrow \mathbb{R}$  satisfy the conditions (F1) – (F4) and (G1) – (G3) respectively. Then, there exists  $\lambda_0 > 0$  such that to every  $\lambda \in ]\lambda_0, +\infty[$  it corresponds a nonempty open interval  $I_\lambda \subset \mathbb{R}$  such that for every  $\mu \in I_\lambda$  problem  $(P''_{\lambda,\mu})$  has at least two distinct, nontrivial weak solutions  $u_{\lambda,\mu}$  and  $v_{\lambda,\mu}$ , with the property*

$$\mathcal{F}_{\lambda,\mu}(u_\lambda, \mu) < 0 < \mathcal{F}_{\lambda,\mu}(v_\lambda, \mu).$$

**Lemma 3.9** *For  $\lambda > \lambda_0$  and  $|\mu| \in ]0, \mu_\lambda^*]$  we have*

$$\inf_{u \in E} \mathcal{F}_{\lambda,\mu}(u) < 0.$$

**Proof.** It is sufficient to prove, that for  $\lambda > \lambda_0$  and  $|\mu| \in ]0, \mu_\lambda^*]$  we have  $\mathcal{F}_{\lambda,\mu}(u_0) < 0$ . Indeed,

$$\begin{aligned} \mathcal{F}_{\lambda,\mu}(u_0) &= I(u) - \lambda J_f(u_0) - \lambda \mu J_G(u_0) \leq \\ &\leq \lambda_0 J_F(u_0) - \lambda J_F(u_0) + \lambda |\mu| m = \\ &= (\lambda_0 - \lambda) J_F(u_0) + \lambda |\mu| m = \\ &= (\lambda_0 - \lambda) \frac{\lambda(1+c)\mu_\lambda^*}{\lambda - \lambda_0} + \lambda |\mu| m = \\ &= -(1+m)\lambda \mu_\lambda^* + \lambda |\mu| m = \\ &= -\lambda \mu_\lambda^* - m\lambda(|\mu| - \mu_\lambda^*) < 0 \end{aligned}$$

for all  $\lambda > \lambda_0$  and  $|\mu| \in ]0, \mu_\lambda^*]$ . □

**Lemma 3.10** *For every  $\lambda > \lambda_0$  and  $\mu \in ]0, \mu_\lambda^*]$ , the functional  $\mathcal{F}_{\lambda,\mu}$  satisfies the mountain pass geometry.*

**Proof.** The assumptions (F1),(F2), (G1) and (G2) imply for every  $\varepsilon > 0$  the existence of  $c_1(\varepsilon), c_2(\varepsilon) > 0$  such that

$$|F(x, u(x))| \leq \varepsilon f_0(x) |u(x)|^p + c_1(\varepsilon) (f_0(x) + f_1(x)) |u(x)|^r, \quad (3.47)$$

$$|G(x, u(x))| \leq \varepsilon g_0(x) |u(x)|^p + c_2(\varepsilon) (g_0(x) + g_1(x)) |u(x)|^m, \quad (3.48)$$

where  $r \in ]p, p^*[$  and  $m \in \left[ p, p \frac{N-1}{N-p} \right]$ . Using again (F1) and (G1), we get

$$|F(x, u(x))| \leq \varepsilon w_1(x) C_f |u(x)|^p + 2c_1(\varepsilon) C_f w_1(x) |u(x)|^r, \quad (3.49)$$

$$|G(x, u(x))| \leq \varepsilon w_2(x) C_g |u(x)|^p + 2c_2(\varepsilon) C_g w_2(x) |u(x)|^m. \quad (3.50)$$

Fix  $\lambda > \lambda_0$  and  $\mu \in ]0, \mu_\lambda^*]$ , then using relations (3.49) and (3.50) we

have for every  $u \in E$

$$\begin{aligned}
\mathcal{F}_{\lambda,\mu}(u) &= \frac{1}{p} \|u\|_b^p - \lambda J_\mu(u) \\
&\geq \frac{1}{p} \|u\|_b^p - \lambda \int_\Omega |F(x, u(x))| dx - \lambda |\mu| \int_\Gamma |G(x, u(x))| d\Gamma \\
&\geq \frac{1}{p} \|u\|_b^p - \lambda \varepsilon C_f \int_\Omega w_1(x) |u(x)|^p dx \\
&\quad - 2\lambda c_1(\varepsilon) C_f \int_\Omega w_1(x) |u(x)|^r dx - \lambda |\mu| \varepsilon C_g \int_\Gamma w_2(x) |u(x)|^p d\Gamma \\
&\quad - 2\lambda |\mu| c_2(\varepsilon) C_g \int_\Gamma w_2(x) |u(x)|^m d\Gamma \\
&\geq \frac{1}{p} \|u\|_b^p - \lambda \varepsilon C_f \|u\|_{p,w_1}^p - 2\lambda c_1(\varepsilon) C_f \|u\|_{r,w_1}^r \\
&\quad - \lambda |\mu| \varepsilon C_g \|u\|_{p,w_2}^p - 2\lambda |\mu| c_2(\varepsilon) C_g \|u\|_{m,w_2}^m \\
&\quad \left( \frac{1}{p} - \lambda \varepsilon C_f C_{p,w_1}^p - \lambda |\mu| \varepsilon C_g C_{p,w_2}^p \right) \|u\|_b^p \\
&\quad 2\lambda c_1(\varepsilon) C_f C_{r,w_1}^r \|u\|_b^r - 2\lambda |\mu| c_2(\varepsilon) C_g C_{m,w_2}^m \|u\|_b^m.
\end{aligned}$$

Using the notations

$$A = \left( \frac{1}{p} - \lambda \varepsilon C_f C_{p,w_1}^p - \lambda |\mu| \varepsilon C_g C_{p,w_2}^p \right),$$

$$B = 2\lambda c_1(\varepsilon) C_f C_{r,w_1}^r, \quad C = 2\lambda |\mu| c_2(\varepsilon) C_g C_{m,w_2}^m,$$

we get

$$\mathcal{F}_{\lambda,\mu}(u) \geq (A - B \|u\|_b^{r-p} - C \|u\|_b^{m-p}) \|u\|_b^p.$$

We choose  $\varepsilon \in ]0, \frac{1}{2p} \frac{1}{\lambda(C_f C_{p,w_1}^p + |\mu| C_g C_{p,w_2}^p)} [$ , then  $A > 0$ . Let  $l : \mathbb{R}_+ \rightarrow \mathbb{R}$  be defined by  $l(t) = A - B t^{r-p} - C t^{m-p}$ . We can see, that  $l(0) = A > 0$ , so there exists  $\varepsilon^* > 0$  such that  $l(t) > 0$  for every  $t \in ]0, \varepsilon^*[$ . Then, for every  $u \in E$ , with  $\|u\| = \varepsilon^{**} < \min\{\varepsilon^*, \|u_0\|\}$ , we have  $\mathcal{F}_{\lambda,\mu}(u) \geq \eta(\lambda, \mu, \varepsilon^*) > 0$ . From Lemma 3.9 we obtain  $\mathcal{F}_{\lambda,\mu}(u_0) < 0$ .

Therefore, the functional  $\mathcal{F}_{\lambda,\mu}$  satisfies the hypotheses of Theorem 1.7.

□

**Proof of Theorem 3.7:** Fix  $\lambda > \lambda_0$  and  $\mu \in ]0, \mu_\lambda^*[ = I_\lambda$ . From Lemma 3.8 we have that the functional  $\mathcal{F}_{\lambda,\mu}$  is coercive and satisfies the (PS) condition. Then, there exists an element  $u_{\lambda,\mu} \in E$  such that  $\mathcal{E}_{\lambda,\mu}(u_{\lambda,\mu}) = \inf_{v \in E} \mathcal{F}_{\lambda,\mu}(v)$  (see [253]). By using Lemma 3.9 we obtain  $\mathcal{F}_{\lambda,\mu}(u_{\lambda,\mu}) < 0$ . On the other hand, by Lemma 3.10 and Theorem 1.8, it follows that there exists an element  $v_{\lambda,\mu} \in E$  such that  $\mathcal{F}'_{\lambda,\mu}(v_{\lambda,\mu}) = 0$  and  $\mathcal{F}_{\lambda,\mu}(v_{\lambda,\mu}) \geq \eta(\lambda, \mu, \varepsilon^*) > 0$ . □

### 3.5 Comments and further perspectives

*A. Comments.* The problems considered in this chapter extend in a quasilinear framework the *stationary Schrödinger equation*

$$-\Delta u + V(x)u = au \quad \text{in } \Omega.$$

This equation describes the motion of a particle interacting with a force field  $F(x) = -\nabla V(x)$ , where the function  $V : \Omega \rightarrow \mathbb{R}$  is called a *potential*. The associated energy functional is

$$E(u) = E_1(u) + E_2(u),$$

where

$$E_1(u) = \frac{1}{2} \int_{\Omega} |\nabla u|^2 dx \quad \text{and} \quad E_2(u) = \frac{1}{2} \int_{\Omega} [V(x)u^2 - au^2] dx.$$

Physically,  $E_1(u)$  is called the *kinetic energy* of the particle  $u$ ,  $E_2(u)$  is its *potential energy*, while  $E(u)$  is the *total energy* of  $u$ .

In some situations (see, e.g., Proposition 3.1), the kinetic energy dominates the potential energy. Any inequality in which the kinetic energy dominates some kind of integral of  $u$  (but not depending on  $\nabla u$ ) is called an *uncertainty principle*. As stated in [192], “the historical reason for this strange appellation is that such an inequality implies that one cannot make the potential energy very negative without also making the kinetic energy large”.

In this chapter we have been interested in the qualitative analysis of weak solutions to the quasilinear mixed boundary value problem

$$(P_{\lambda, \mu}) \quad \begin{cases} -\operatorname{div}(a(x)|\nabla u|^{p-2}\nabla u) = \lambda\alpha(x, u) + \beta(x, u) & \text{in } \Omega, \\ a(x)|\nabla u|^{p-2}\nabla u \cdot \mathbf{n} + b(x)|u|^{p-2}u = \mu h(x, u) & \text{on } \Gamma, \end{cases}$$

where  $\Omega \subset \mathbb{R}^N$  is an unbounded domain with (possible noncompact) smooth boundary  $\Gamma$ . In our treatment a central role is played by the compact embedding result Theorem 3.1, due to Pflüger [241], [242]. In our results, the functions  $\alpha, \beta$  and  $h$  satisfies subcritical growth condition, which implies that the energy functional associated to the problem  $(P_{\lambda, \mu})$  satisfies the (PS) condition. The (PS) condition play a key place in the mountain pass theorem and Ricceri type results used in the proof of our results. The results of this chapter are based on the papers Montefusco and Rădulescu [217], Cârstea and Rădulescu [71], Lisei, Varga, and Horváth [195], and Mezei and Varga [213].

*B. Further perspectives.* In the last years the fibering method introduced and developed by Pohozaev [245], [246], [247] can be applied with

success in the study of existence and multiplicity of solutions for bifurcation equations, homogenous Dirichlet or Neumann problems which contain  $p$ -Laplacian (see [248]). El Hamidi in the paper [104] use this method to study mixed homogeneous problems. Mitidieri and Bozhkov in [215] use an early modified version of fibering method and prove the existence and multiplicity results for quasilinear system, which contains  $p, q$ -Laplacians. In [144] the authors study positive solutions for indefinite inhomogeneous Neumann elliptic problems using this methods. Using the concentration compactness principle [194] combined with fibering method we propose the following problem: find  $u \in E$  such that

$$(P_{\lambda, \mu}^*) \quad \begin{cases} -\operatorname{div}(a(x)|\nabla u|^{p-2}\nabla u) = \lambda K(x)|u|^r + k(x)|u|^{p-2}u & \text{in } \Omega, \\ a(x)|\nabla u|^{p-2}\nabla u \cdot \mathbf{n} + b(x)|u|^{p-2}u = \mu L(x)|u|^q & \text{on } \Gamma, \end{cases},$$

where  $r = p^* = \frac{pN}{N-p}, q < \bar{p}^* = \frac{p(N-1)}{N-p}$  or  $r < p^* = \frac{pN}{N-p}, q = \bar{p}^* = \frac{p(N-1)}{N-p}$  or  $r = p^* = \frac{pN}{N-p}, q = \bar{p}^* = \frac{p(N-1)}{N-p}$  or  $p^* = \frac{pN}{N-p}, \bar{p}^* = \frac{p(N-1)}{N-p}$  represent the critical Sobolev exponent for the embedding  $E \hookrightarrow L^r(\Omega; w_1)$  and  $E \hookrightarrow L^q(\Gamma; w_2)$ .

The problem  $(P_{\lambda, \mu}^*)$  is similar with Yamabe problem on manifold with boundary, see the papers of Escobar [106], [107] and [108].

## 4

# Elliptic Systems of Gradient Type

Rigor is to the mathematician  
what morality is to men.

---

André Weil (1906–1998)

### 4.1 Introduction

This chapter examines some basic facts of the theory of boundary value problems for elliptic systems of partial differential equations, a theory which has many applications in mathematics and the physical sciences. We are mainly concerned with the existence and multiplicity results for eigenvalue problems involving elliptic systems of gradient type. Important contributions to the study of these problems are due to Esteban [109], Grossinho and Tersian [134], Fan and Zhao [112]. In this chapter we present some new results in this direction using results of Bartolo, Benci and Fortunato [25], and some recent contributions due to Ricceri [259], [261], [263], [265] and Tsar'kov [290]. A simple model treated in this chapter is given by

$$\begin{cases} -\Delta u = \lambda |\nabla v|^p & \text{in } \Omega \\ -\Delta v = \lambda |\nabla u|^q & \text{in } \Omega \\ u = v = 0 & \text{on } \partial\Omega, \end{cases} \quad (4.1)$$

where  $p$  and  $q$  are positive numbers. Problem (4.1) is often referred to as a stationary Hamilton-Jacobi system and is also related to the Kardar-Parisi-Zhang system arising in the physical theory of growth and roughening surfaces.

## 4.2 Formulation of the problems

Let  $\Omega \subseteq \mathbb{R}^N$  a stripe-like domain, that is,  $\Omega = \omega \times \mathbb{R}^l$ , where  $\omega \subset \mathbb{R}^m$  is bounded with smooth boundary and  $m \geq 1, l \geq 2, 1 < p, q < N = m+l$ . Let  $F : \Omega \times \mathbb{R}^2 \rightarrow \mathbb{R}$  be a function of class  $C^1$ . We consider the following boundary value problem:

$$(S_{p,q}^\lambda) \quad \begin{cases} -\Delta_p u = \lambda F_u(x, u, v) & \text{in } \Omega, \\ -\Delta_q v = \lambda F_v(x, u, v) & \text{in } \Omega, \\ u = v = 0 & \text{on } \partial\Omega, \end{cases}$$

We suppose that the function  $F : \Omega \times \mathbb{R}^2 \rightarrow \mathbb{R}$  satisfies the following hypotheses:

**(F1)**  $F : \Omega \times \mathbb{R}^2 \rightarrow \mathbb{R}$  is continuous,  $(s, t) \mapsto F(x, s, t)$  is of class  $C^1$  and  $F(x, 0, 0) = 0$  for every  $x \in \Omega$ .

**(F2)** There exist  $c_1 > 0$  and  $r \in (p, p^*), s \in (q, q^*)$  such that

$$|F_u(x, u, v)| \leq c_1(|u|^{p-1} + |v|^{(p-1)q/p} + |u|^{r-1}), \quad (4.2)$$

$$|F_v(x, u, v)| \leq c_1(|v|^{q-1} + |u|^{(q-1)p/q} + |v|^{s-1}) \quad (4.3)$$

for every  $x \in \Omega$  and  $(u, v) \in \mathbb{R}^2$ .

We consider the Sobolev space  $W_0^{1,\alpha}(\Omega)$  ( $\alpha \in \{p, q\}$ ) endowed with the norm

$$\|u\|_{1,\alpha} = \left( \int_{\Omega} |\nabla u|^\alpha \right)^{1/\alpha},$$

with  $\alpha \in \{p, q\}$ .

We say that  $(u, v) \in W_0^{1,p} \times W_0^{1,q}$  is a weak solution of the problem  $(S_{p,q,\Omega})$  if

$$\begin{cases} \int_{\Omega} |\nabla u|^{p-2} \nabla u \nabla w_1 dx - \lambda \int_{\Omega} F'_u(x; u(x), v(x)) w_1(x) dx = 0 \\ \int_{\Omega} |\nabla v|^{q-2} \nabla v \nabla w_2 dx - \lambda \int_{\Omega} F'_v(x; u(x), v(x)) w_2(x) dx = 0, \end{cases}$$

for every  $(w_1, w_2) \in W_0^{1,p} \times W_0^{1,q}$ .

The product space  $W_0^{1,p}(\Omega) \times W_0^{1,q}(\Omega)$  is endowed with the norm  $\|(u, v)\|_{1,p,q} = \|u\|_{1,p} + \|v\|_{1,q}$ . We define the function  $\mathcal{F} : W_0^{1,p}(\Omega) \times W_0^{1,q}(\Omega) \rightarrow \mathbb{R}$  by

$$\mathcal{F}(u, v) = \int_{\Omega} F(x; u, v) dx$$

for  $u \in W_0^{1,p}(\Omega)$ ,  $v \in W_0^{1,q}(\Omega)$ . The energy functional associated to problem  $(S_{p,q}^\lambda)$  is defined by

$$E(u, v) = \frac{1}{p} \|u\|_{1,p}^p + \frac{1}{q} \|v\|_{1,q}^q - \int_{\Omega} F(x; u, v) dx.$$

Taking into account conditions **(F1)** and **(F2)**, it follows that  $E$  is of class  $C^1$  and its critical points are weak solutions of the problem  $(S_{p,q}^\lambda)$ .

The embedding  $W_0^{1,\alpha}(\Omega) \hookrightarrow L^\beta(\Omega)$  for  $\beta \in [\alpha, \alpha^*]$  ( $\alpha \in \{p, q\}$ ) are continuous. Let  $c_{\beta,\alpha} > 0$  denote the embedding constant, that is,  $\|u\|_\beta \leq c_{\beta,\alpha} \|u\|_{1,\alpha}$  for all  $u \in W_0^{1,\alpha}(\Omega)$ . Since the embedding  $W_0^{1,\alpha}(\Omega) \hookrightarrow L^\beta(\Omega)$  for  $\beta \in [\alpha, \alpha^*]$  ( $\alpha \in \{p, q\}$ ) are not compact, we introduce the action of  $G = id^m \times O(N - m)$  on  $W_0^{1,\alpha}(\Omega)$  as

$$gu(x, y) = u(x, g_0^{-1}y)$$

for all  $(x, y) \in \omega \times \mathbb{R}^{N-m}$ ,  $g = id^m \times g_0 \in G$  and  $u \in W_0^{1,\alpha}(\Omega)$ ,  $\alpha \in \{p, q\}$ . Moreover, the action  $G$  on  $W_0^{1,\alpha}(\Omega)$  is isometric, that is  $\|gu\|_{1,\alpha} = \|u\|_{1,\alpha}$  for all  $g \in G$ ,  $u \in W_0^{1,\alpha}(\Omega)$ ,  $\alpha \in \{p, q\}$ . Let

$$W_{0,G}^{1,\alpha}(\Omega) = \{u \in W_0^{1,\alpha}(\Omega) : gu = u \text{ for all } g \in G\}, \quad \alpha \in \{p, q\},$$

denote the closed subspace of axially symmetric functions of  $W_0^{1,\alpha}(\Omega)$ . The embeddings  $W_{0,G}^{1,\alpha}(\Omega) \hookrightarrow L^\beta(\Omega)$ ,  $\alpha < \beta < \alpha^*$ ,  $\alpha \in \{p, q\}$ , are compact, by Theorem B.7. The fixed points set of the action  $G$  on  $W_0^{1,p}(\Omega) \times W_0^{1,q}(\Omega)$  is  $W_{0,G}^{1,p,q}(\Omega) = W_{0,G}^{1,p}(\Omega) \times W_{0,G}^{1,q}(\Omega)$ . In order to apply the principle of symmetric criticality, we introduce an invariance condition for the function  $F$ , hence we suppose that:

**(FI)**  $F(gx, u, v) = F(x, u, v)$  for every  $x \in \Omega$ ,  $g \in G$  and  $(u, v) \in \mathbb{R}^2$ .

Therefore to study the weak solutions of the problem  $(S_{p,q}^\lambda)$  it is sufficient to study the existence of critical points of the function  $E_G = E|_{W_{0,G}^{1,p,q}(\Omega)}$ . To manipulate easier the problem  $(S_{p,q}^\lambda)$  we introduce the following notations:

$$\langle A_p u, w_1 \rangle_p = \int_{\Omega} |\nabla u|^{p-2} \nabla u \nabla w_1 dx$$

and

$$\langle A_q v, w_2 \rangle_q = \int_{\Omega} |\nabla v|^{q-2} \nabla v \nabla w_2 dx,$$

where  $\langle \cdot, \cdot \rangle_p$  ( $\langle \cdot, \cdot \rangle_q$ ) means the duality between  $W_0^{1,p}(\Omega)$  and  $(W_0^{1,p}(\Omega))^*$  ( $W_0^{1,q}(\Omega)$  and  $(W_0^{1,q}(\Omega))^*$ ) respectively. Also we denote in the same way the restriction of the norms  $\|\cdot\|_{1,p}$ ,  $\|\cdot\|_{1,q}$  and  $\langle \cdot, \cdot \rangle_p$ ,  $\langle \cdot, \cdot \rangle_q$  restricted to  $W_{0,G}^{1,p,q}(\Omega)$ .

With the above notations we have  $A_p : W_0^{1,p}(\Omega) \rightarrow (W_0^{1,p}(\Omega))^*$  and  $A_q : W_0^{1,q}(\Omega) \rightarrow (W_0^{1,q}(\Omega))^*$  are the duality mappings induced by the functions  $t \in [0, +\infty[ \mapsto t^{p-1} \in [0, +\infty[$ , and  $t \in [0, +\infty[ \mapsto t^{q-1} \in [0, +\infty[$ , respectively.

This chapter is divided in three sections. In the first section we study the case when the function  $F$  is superlinear at infinity, the second section contain the case when  $F$  is sublinear, while the section three contains a special class of system.

### 4.3 Systems with superlinear potential

In this section we prove an existence result for problem  $((S_{p,q}^\lambda))$  in the case when  $\lambda = 1$  and the function  $F$  is independent of  $x$ , that is,  $F : \mathbb{R}^2 \rightarrow \mathbb{R}$  and satisfies the conditions **(F1)**, **(F2)**. In this case we have the following problem:

$$(S_{p,q}) \quad \begin{cases} -\Delta_p u = F_u(u, v) & \text{in } \Omega, \\ -\Delta_q v = F_v(u, v) & \text{in } \Omega, \\ u = v = 0 & \text{on } \partial\Omega, \end{cases}$$

This problem consists in finding  $(u, v) \in W_0^{1,p}(\Omega) \times W_0^{1,q}(\Omega)$  such that

$$\begin{cases} \langle A_p(u), w_1 \rangle - \int_{\Omega} F'_u(u(x), v(x)) w_1(x) dx = 0 \\ \langle A_q(v), w_2 \rangle - \int_{\Omega} F'_v(u(x), v(x)) w_2(x) dx = 0, \end{cases}$$

for every  $(w_1, w_2) \in W_0^{1,p}(\Omega) \times W_0^{1,q}(\Omega)$ .

In this case the energy functional associated to the problem  $(S_{p,q})$  is given by

$$E(u, v) = \frac{1}{p} \|u\|_{1,p}^p + \frac{1}{q} \|v\|_{1,q}^q - \mathcal{F}(u, v), \quad (4.4)$$

where  $\mathcal{F}(u, v) = \int_{\Omega} F(u(x), v(x)) dx$ . Its derivative in  $(u, v) \in W_0^{1,p}(\Omega) \times W_0^{1,q}(\Omega)$  is given by

$$E'(u, v)(w_1, w_2) = \langle A_p(u), w_1 \rangle_p + \langle A_q(v), w_2 \rangle_q - \mathcal{F}'(u, v)(w_1, w_2), \quad (4.5)$$

for every  $(w_1, w_2) \in W_0^{1,p}(\Omega) \times W_0^{1,q}(\Omega)$ , where

$$\mathcal{F}'(u, v)(w_1, w_2) = \int_{\Omega} [F_u(u, v)w_1 + F_v(u, v)w_2] dx. \quad (4.6)$$

As in introduction we denote by  $E_G = E|_{W_{0,G}^{1,p,q}(\Omega)}$  and  $\mathcal{F}_G = \mathcal{F}|_{W_{0,G}^{1,p,q}(\Omega)}$ . In the following we study the only the existence of critical points of the functional  $E_G$ .

In the following we suppose that the function  $F : \mathbb{R}^2 \rightarrow \mathbb{R}$  beside of conditions **(F1)**, **(F2)** satisfies the following assumptions:

**(F3)** there exist  $c_2 > 0$  and  $\mu, \nu \geq 1$  such that

$$-c_2(|u|^\mu + |v|^\nu) \geq F(u, v) - \frac{1}{p}F'_u(u, v)u - \frac{1}{q}F'_v(u, v)v, \quad (4.7)$$

for all  $(u, v) \in \mathbb{R}^2$ .

$$\text{(F4)} \quad \lim_{u,v \rightarrow 0} \frac{F'_u(u, v)}{|u|^{p-1}} = \lim_{u,v \rightarrow 0} \frac{F'_v(u, v)}{|v|^{q-1}} = 0,$$

The main result of this section is the following.

**Theorem 4.1** *We suppose that the function  $F : \Omega \times \mathbb{R}^2 \rightarrow \mathbb{R}$  satisfies the conditions **(F1)**-**(F4)** with  $ps = qr$  and*

$$\mu > \max\{p, N(r-p)/p\} \text{ and } \nu > \max\{q, N(s-q)/q\}. \quad (4.8)$$

*Then, (S) admits at least one nonzero solution.*

We will use the following auxiliary result.

**Proposition 4.1** *For all  $t > 1$  and  $(u, v) \in \mathbb{R}^2$  we have*

$$\begin{aligned} F(t^{\frac{1}{p}}u, t^{\frac{1}{q}}v) &\geq tF(u, v) \\ &+ c_2 \left[ \frac{p}{\mu-p}(t^{\frac{\mu}{p}} - t)|u|^\mu + \frac{q}{\nu-q}(t^{\frac{\nu}{q}} - t)|v|^\nu \right]. \end{aligned} \quad (4.9)$$

*Proof* Fix  $(u, v) \in \mathbb{R}^2$ . We define the function  $g : ]0, \infty[ \rightarrow \mathbb{R}$  by

$$g(t) = t^{-1}F(t^{\frac{1}{p}}u, t^{\frac{1}{q}}v) - c_2 \frac{p}{\mu-p} t^{\frac{\mu}{p}-1} |u|^\mu - c_2 \frac{q}{\nu-q} t^{\frac{\nu}{q}-1} |v|^\nu.$$

Since  $g$  is of class  $C^1$ , due to the mean value theorem, for a fixed  $t > 1$  there exists  $\tau = \tau(t, u, v) \in ]1, t[$  such that

$$g(t) - g(1) = g'(\tau)(t - 1).$$

We have

$$\partial_t F(t^{\frac{1}{p}}u, t^{\frac{1}{q}}v) = \frac{1}{p}F'_u(t^{\frac{1}{p}}u, t^{\frac{1}{q}}v)t^{\frac{1}{p}-1}u + \frac{1}{q}F'_v(t^{\frac{1}{p}}u, t^{\frac{1}{q}}v)t^{\frac{1}{q}-1}v.$$

Hence,

$$g'(t) = -t^{-2}F(t^{\frac{1}{p}}u, t^{\frac{1}{q}}v) + t^{-1} \left[ \frac{1}{p}F'_u(t^{\frac{1}{p}}u, t^{\frac{1}{q}}v)t^{\frac{1}{p}-1}u \right. \\ \left. + \frac{1}{q}F'_v(t^{\frac{1}{p}}u, t^{\frac{1}{q}}v)t^{\frac{1}{q}-1}v \right] - c_2 \left[ t^{\frac{\mu}{p}-2}|u|^\mu + t^{\frac{\nu}{q}-2}|v|^\nu \right].$$

Therefore,

$$g(t) - g(1) = -\tau^{-2}[F(\tau^{\frac{1}{p}}u, \tau^{\frac{1}{q}}v) - \frac{1}{p}F'_u(\tau^{\frac{1}{p}}u, \tau^{\frac{1}{q}}v)\tau^{\frac{1}{p}}u \\ - \frac{1}{q}F'_v(\tau^{\frac{1}{p}}u, \tau^{\frac{1}{q}}v)\tau^{\frac{1}{q}}v + c_2(|\tau^{\frac{1}{p}}u|^\mu + |\tau^{\frac{1}{q}}v|^\nu)](t-1).$$

Due to (4.7), we have  $g(t) \geq g(1)$ . Thus,

$$F(t^{\frac{1}{p}}u, t^{\frac{1}{q}}v) \geq tF(u, v) + c_2 \left[ \frac{p}{\mu-p}(t^{\frac{\mu}{p}} - t)|u|^\mu + \frac{q}{\nu-q}(t^{\frac{\nu}{q}} - t)|v|^\nu \right]$$

for all  $t > 1$  and  $(u, v) \in \mathbb{R}^2$ . □

**Proposition 4.2** *If the assumptions of Theorem 4.1 are satisfied, then the functional  $E_G$  satisfies the Cerami  $(C)_c$  condition for all  $c > 0$  (see Remark 1.3),*

*Proof* The norm of the Banach space  $W_{0,G}^{1,p,q}(\Omega)$  is  $\|(u, v)\|_{1,p,q} = \|u\|_{1,p} + \|v\|_{1,q}$ . We denote the norm on the dual space  $(W_{0,G}^{1,p,q}(\Omega))^*$  by  $\|\cdot\|_*$ . Now, let  $(u_n, v_n) \in W_{0,G}^{1,p,q}(\Omega)$  be such that

$$E_G(u_n, v_n) \rightarrow c > 0, \quad (4.10)$$

and

$$(1 + \|(u_n, v_n)\|_{1,p,q})\|E'_G(u_n, v_n)\|_* \rightarrow 0, \quad (4.11)$$

as  $n \rightarrow +\infty$ . Moreover, we have

$$\left\langle E'_G(u_n, v_n) \left( \frac{1}{p}u_n, \frac{1}{q}v_n \right) \right\rangle \geq -(1 + \|(u_n, v_n)\|_{1,p,q})\|E'_G(u_n, v_n)\|_*. \quad (4.12)$$

Using (4.10), (4.11) and (4.7) one has for  $n$  large enough that

$$\begin{aligned} c+1 &\geq E_G(u_n, v_n) - E'_G(u_n, v_n)\left(\frac{1}{p}u_n, \frac{1}{q}v_n\right) \\ &= -\mathcal{F}_G(u_n, v_n) + \mathcal{F}'_G(u_n, v_n)\left(\frac{1}{p}u_n, \frac{1}{q}v_n\right) \\ &= -\int_{\Omega} [F(u_n, v_n) - \frac{1}{p}F'_u(u_n, v_n)u_n \\ &\quad - \frac{1}{q}F'_v(u_n, v_n)v_n] dx \geq c_2 \int_{\Omega} [|u_n|^\mu + |v_n|^\nu] dx. \end{aligned}$$

From the previous inequality we obtain that

$$\{(u_n, v_n)\} \text{ is bounded in } L^\mu(\Omega) \times L^\nu(\Omega). \quad (4.13)$$

By **(F4)** we have that for all  $\varepsilon > 0$  there exists  $\delta(\varepsilon) > 0$  such that, if  $|u|^{p-1} + |v|^{(p-1)q/p} < \delta(\varepsilon)$ , then

$$|F'_u(u, v)| < \varepsilon \left( |u|^{p-1} + |v|^{(p-1)q/p} \right).$$

If  $|u|^{p-1} + |v|^{(p-1)q/p} \geq \delta(\varepsilon)$ , by using (4.2), we have

$$\begin{aligned} |F'_u(u, v)| &\leq c_1[ (|u|^{p-1} + |v|^{(p-1)q/p})^{\frac{r-1}{p-1}} (\delta(\varepsilon))^{\frac{p-r}{p-1}} + |u|^{r-1} ] \\ &\leq c(\varepsilon)(|u|^{r-1} + |v|^{(r-1)q/p}). \end{aligned}$$

Combining the above relations, we have that for all  $\varepsilon > 0$  there exists  $c_1(\varepsilon) > 0$  such that

$$|F'_u(u, v)| < \varepsilon \left( |u|^{p-1} + |v|^{(p-1)q/p} \right) + c_1(\varepsilon) \left( |u|^{r-1} + |v|^{(r-1)q/p} \right) \quad (4.14)$$

for all  $(u, v) \in \mathbb{R}^2$ . A similar calculation shows that for all  $\varepsilon > 0$  there exists  $c_2(\varepsilon) > 0$  such that

$$|F'_v(u, v)| < \varepsilon \left( |v|^{q-1} + |u|^{(q-1)p/q} \right) + c_2(\varepsilon) \left( |v|^{s-1} + |u|^{(s-1)p/q} \right) \quad (4.15)$$

for all  $(u, v) \in \mathbb{R}^2$ .

Using (4.14), (4.15), and keeping in mind that  $F(0, 0) = 0$ , for all  $\varepsilon > 0$  there exists  $c(\varepsilon) = c(c_1(\varepsilon), c_2(\varepsilon)) > 0$  such that

$$\begin{aligned} F(u, v) &\leq \varepsilon(|u|^p + |v|^{(p-1)q/p}|u| + |v|^q + |u|^{(q-1)p/q}|v|) \\ &\quad + c(\varepsilon)(|u|^r + |v|^{(r-1)q/p}|u| + |v|^s + |u|^{(s-1)p/q}|v|) \end{aligned} \quad (4.16)$$

for all  $(u, v) \in \mathbb{R}^2$ .

After integration we use the relation  $ps = qr$ , as well as Young's inequality and Hölder's inequality to obtain

$$\begin{aligned} \mathcal{F}_G(u_n, v_n) &\leq \varepsilon \left[ \left(2 + \frac{1}{p} - \frac{1}{q}\right) \|u_n\|_p^p + \left(2 + \frac{1}{q} - \frac{1}{p}\right) \|v_n\|_q^q \right] \\ &\quad + c(\varepsilon) \left[ \left(2 + \frac{1}{r} - \frac{1}{s}\right) \|u_n\|_r^r + \left(2 + \frac{1}{s} - \frac{1}{r}\right) \|v_n\|_s^s \right]. \end{aligned}$$

Therefore, one has

$$\begin{aligned} &\left[ \frac{1}{p} - \varepsilon \left(2 + \frac{1}{p} - \frac{1}{q}\right) c_{p,p}^p \right] \|u_n\|_{1,p}^p + \left[ \frac{1}{q} - \varepsilon \left(2 + \frac{1}{q} - \frac{1}{p}\right) c_{q,q}^q \right] \|v_n\|_{1,q}^q \leq \\ &\leq E_G(u_n, v_n) + c(\varepsilon) \left[ \left(2 + \frac{1}{r} - \frac{1}{s}\right) \|u_n\|_r^r + \left(2 + \frac{1}{s} - \frac{1}{r}\right) \|v_n\|_s^s \right]. \end{aligned}$$

Choosing

$$0 < \varepsilon < \frac{1}{3} \min \left\{ \frac{1}{pc_{p,p}^p}, \frac{1}{qc_{q,q}^q} \right\}, \quad (4.17)$$

we find  $c_3(\varepsilon), c_4(\varepsilon) > 0$  such that

$$c_3(\varepsilon) (\|u_n\|_{1,p}^p + \|v_n\|_{1,q}^q) \leq c + 1 + c_4(\varepsilon) (\|u_n\|_r^r + \|v_n\|_s^s) \quad (4.18)$$

for  $n$  large enough. Now, we will examine the behaviour of the sequences  $\{\|u_n\|_r^r\}$  and  $\{\|v_n\|_s^s\}$ , respectively.

To this end, observe that  $\mu \leq r$  and  $\nu \leq s$ . From relations (4.2), (4.3) and from the fact that for all  $\beta \in ]0, \infty[$  there exists a constant  $c(\beta) > 0$  such that

$$(x + y)^\beta \leq c(\beta)(x^\beta + y^\beta), \quad \text{for all } x, y \in [0, \infty[,$$

we have

$$\begin{aligned} &|F(u, v) - F(w, y)| \leq \\ &c_3 [|u - w| (|u|^{p-1} + |w|^{p-1} + |v|^{(p-1)q/p} + |y|^{(p-1)q/p} + |u|^{r-1} + |w|^{r-1}) + \\ &|v - y| (|v|^{q-1} + |y|^{q-1} + |u|^{(q-1)p/q} + |w|^{(q-1)p/q} + |v|^{s-1} + |y|^{s-1})], \end{aligned} \quad (4.19)$$

where  $c_3 = c_3(c_1, p, q, r, s) > 0$ .

Keeping in mind the relation  $ps = qr$ , taking  $w = y = 0$  and  $u = ut^{\frac{1}{p}}$ ,  $v = vt^{\frac{1}{q}}$  ( $t > 1$ ) in (4.19), from (4.9) follow the required relations.

We distinguish two cases:

- I)  $\mu = r$ : From (4.13) we have that  $\{\|u_n\|_r^r\}$  is bounded.  
 II)  $\mu \in ]p, N(r-p)/p\}$ ,  $r[$ : we use the interpolation inequality

$$\|u\|_r \leq \|u\|_\mu^{1-\delta} \|u\|_{p^*}^\delta \quad \text{for all } u \in L^\mu(\Omega) \cap L^{p^*}(\Omega)$$

with  $\delta = \frac{p^*}{r} \cdot \frac{r-\mu}{p^*-\mu}$ . From (4.13) and the fact that the embedding  $W_{0,G}^{1,p}(\Omega) \hookrightarrow L^{p^*}(\Omega)$ , is continuous, we have that there exists  $c_5 > 0$  such that  $\|u_n\|_r^r \leq c_5 \|u_n\|_{1,p}^{\delta r}$ , with  $\delta r < p$ .

Taking into consideration the analogous relations for the sequence  $\{\|v_n\|_s^s\}$ , we conclude from (4.18) that the sequences  $\{\|u_n\|_{1,p}\}$  and  $\{\|v_n\|_{1,q}\}$  are bounded. Because the embeddings  $W_{0,G}^{1,p}(\Omega) \hookrightarrow L^r(\Omega)$ ,  $W_{0,G}^{1,q}(\Omega) \hookrightarrow L^s(\Omega)$  are compact, therefore up to a subsequence, we have

$$(u_n, v_n) \rightarrow (u, v) \text{ weakly in } W_{0,G}^{1,p,q}(\Omega), \quad (4.20)$$

$$u_n \rightarrow u \text{ strongly in } L^r(\Omega), \quad (4.21)$$

$$v_n \rightarrow v \text{ strongly in } L^s(\Omega). \quad (4.22)$$

Moreover, we have

$$\begin{aligned} E'_G(u_n, v_n)(u - u_n, v - v_n) &= \langle A_p(u_n), u - u_n \rangle_p + \langle A_q(v_n), v - v_n \rangle_q \\ &\quad - \mathcal{F}'_G(u_n, v_n)(u - u_n, v - v_n) \end{aligned}$$

and

$$\begin{aligned} E'_G(u, v)(u_n - u, v_n - v) &= \langle A_p(u), u_n - u \rangle_p + \langle A_q(v), v_n - v \rangle_q \\ &\quad - \mathcal{F}'_G(u, v)(u_n - u, v_n - v). \end{aligned}$$

By adding the above two relations, we obtain

$$\begin{aligned} J_n &:= \langle A_p(u) - A_p(u_n), u - u_n \rangle_p + \langle A_q(v) - A_q(v_n), v - v_n \rangle_q \\ &= J_n^1 - J_n^2 - J_n^3, \end{aligned} \quad (4.23)$$

where

$$J_n^1 = \mathcal{F}'_G(u_n, v_n)(u_n - u, v_n - v) + \mathcal{F}'_G(u, v)(u - u_n, v - v_n),$$

$$J_n^2 = E'_G(u_n, v_n)(u - u_n, v - v_n) \text{ and } J_n^3 = E'_G(u, v)(u_n - u, v_n - v).$$

In the sequel, we estimate  $J_n^i$  for  $i \in \{1, 2, 3\}$ . Using relations (4.14),

(4.15) and  $ps = qr$ , one has

$$\begin{aligned}
J_n^1 &= \int_{\Omega} [F'(u_n(x), v_n(x))(u_n(x) - u(x), v_n(x) - v(x)) \\
&\quad + F'(u(x), v(x))(u(x) - u_n(x), v(x) - v_n(x))] dx \\
&\leq \int_{\Omega} [|F'_u(u_n(x), v_n(x))(u_n(x) - u(x))| \\
&\quad + |F'_v(u_n(x), v_n(x))(v_n(x) - v(x))| + |F'_u(u(x), v(x))(u(x) - u_n(x))| \\
&\quad + |F'_v(u(x), v(x))(v(x) - v_n(x))|] dx \\
&\leq \varepsilon \left[ (\|u_n\|_p^{p-1} + \|u\|_p^{p-1} + \|v_n\|_q^{(p-1)q/p} + \|v\|_q^{(p-1)q/p}) \|u - u_n\|_p \right. \\
&\quad \left. + (\|v_n\|_q^{q-1} + \|v\|_q^{q-1} + \|u_n\|_p^{(q-1)p/q} + \|u\|_p^{(q-1)p/q}) \|v - v_n\|_q \right] \\
&\quad + c_1(\varepsilon) \left( \|u_n\|_r^{r-1} + \|u\|_r^{r-1} + \|v_n\|_s^{(r-1)s/r} + \|v\|_s^{(r-1)s/r} \right) \|u - u_n\|_r \\
&\quad + c_2(\varepsilon) \left( \|v_n\|_s^{s-1} + \|v\|_s^{s-1} + \|u_n\|_r^{(s-1)r/s} + \|u\|_r^{(s-1)r/s} \right) \|v - v_n\|_s.
\end{aligned}$$

The sequences  $\{u_n\}$  and  $\{v_n\}$  are bounded in  $W_{0,G}^{1,p}(\Omega) (\hookrightarrow L^p(\Omega) \cap L^r(\Omega))$  and  $W_{0,G}^{1,q}(\Omega) (\hookrightarrow L^q(\Omega) \cap L^s(\Omega))$ , respectively, then by using the relations (4.21) and (4.22), one has

$$\limsup_{n \rightarrow \infty} J_n^1 \leq 0, \quad (4.24)$$

since  $\varepsilon > 0$  was arbitrary chosen.

The convergence in (4.11) implies  $\|E'_G(u_n, v_n)\|_{\star} \rightarrow 0$ , therefore  $J_n^2 \rightarrow 0$ . Taking into account that  $(u_n, v_n) \rightharpoonup W_{0,G}^{1,p,q}(\Omega)$  weakly, it follows that  $J_n^3 \rightarrow 0$ . Therefore, if  $n \rightarrow \infty$ , then

$$\langle A_p(u) - A_p(u_n), u - u_n \rangle_p + \langle A_q(v) - A_q(v_n), v - v_n \rangle_q \rightarrow 0.$$

On the other hand, from the inequality

$$|t-s|^\alpha \leq \begin{cases} (|t|^{\alpha-2}t - |s|^{\alpha-2}s)(t-s), & \text{if } \alpha \geq 2, \\ ((|t|^{\alpha-2}t - |s|^{\alpha-2}s)(t-s))^{\frac{\alpha}{2}} (|t|^\alpha + |s|^\alpha)^{\frac{2-\alpha}{2}}, & \text{if } 1 < \alpha < 2, \end{cases}$$

for all  $t, s \in \mathbb{R}^N$ , we obtain that

$$\lim_{n \rightarrow \infty} \int_{\Omega} (|\nabla u_n - \nabla u|^p + |\nabla v_n - \nabla v|^q) = 0,$$

that is, the sequences  $\{u_n\}$  and  $\{v_n\}$  are strongly convergent in  $W_{0,G}^{1,p}(\Omega)$  and  $W_{0,G}^{1,q}(\Omega)$ , respectively.  $\square$

To prove Theorem 4.1 we use Theorem 1.7, 1.29 combined with Remark 1.3.

**Proof of Theorem 4.1** From Proposition 4.2 it follows that the functional  $E_G$  satisfies the Cerami  $(C)_c$  condition for every  $c > 0$ . We need to verify that  $E_G$  satisfies the mountain pass geometry. Using (4.16), we obtain

$$\begin{aligned} E_G(u, v) &= \frac{1}{p} \|u\|_{1,p}^p + \frac{1}{q} \|v\|_{1,q}^q - \int_{\Omega} F(u, v) dx \geq \\ &\left[ \frac{1}{p} - \varepsilon \left( 2 + \frac{1}{p} - \frac{1}{q} \right) c_{p,p}^p \right] \|u\|_{1,p}^p + \left[ \frac{1}{q} - \varepsilon \left( 2 + \frac{1}{q} - \frac{1}{p} \right) c_{q,q}^q \right] \|v\|_{1,q}^q \\ &- c(\varepsilon) \left[ \left( 2 + \frac{1}{r} - \frac{1}{s} \right) \|u\|_r^r + \left( 2 + \frac{1}{s} - \frac{1}{r} \right) \|v\|_s^s \right]. \end{aligned}$$

Choosing  $\varepsilon$  as in (4.17), we can fix  $c_5(\varepsilon), c_6(\varepsilon) > 0$  such that

$$E_G(u, v) \geq c_5(\varepsilon) (\|u\|_{1,p}^p + \|v\|_{1,q}^q) - c_6(\varepsilon) (\|u\|_{1,p}^r + \|v\|_{1,q}^s).$$

Since the function  $t \mapsto (x^t + y^t)^{\frac{1}{t}}$ ,  $t > 0$  is non-increasing ( $x, y \geq 0$ ), using again  $ps = qr$ , we have

$$\|u\|_{1,p}^r + \|v\|_{1,q}^s \leq [\|u\|_{1,p}^p + \|v\|_{1,q}^q]^{\frac{r}{p} (= \frac{s}{q})}.$$

Therefore,

$$E_G(u, v) \geq \left[ c_5(\varepsilon) - c_6(\varepsilon) (\|u\|_{1,p}^p + \|v\|_{1,q}^q)^{\frac{r}{p} - 1} \right] (\|u\|_{1,p}^p + \|v\|_{1,q}^q).$$

Let  $0 < \rho < 1$  and denote

$$B_{\rho} = \{(u, v) \in W_{0,G}^{1,p,q}(\Omega) : \|(u, v)\|_{1,p,q} = \rho\}.$$

Then, we have  $(\rho/2)^{\max\{p,q\}} \leq \|u\|_{1,p}^p + \|v\|_{1,q}^q \leq \rho$  for all  $(u, v) \in B_{\rho}$ . Choosing  $\rho$  small enough, there exists  $\eta > 0$  such that  $E_G(u, v) \geq \eta$  for all  $(u, v) \in B_{\rho}$ , due to the fact that  $r > p$ . This proves the first condition from Theorem 1.7.

To prove the second condition from Theorem 1.7, fix  $u^0 \in W_{0,G}^{1,p}(\Omega)$  and  $v^0 \in W_{0,G}^{1,q}(\Omega)$  such that  $\|u^0\|_{1,p} = \|v^0\|_{1,q} = 1$ . Then, for every  $t > 1$  we have

$$\begin{aligned} E_G(t^{\frac{1}{p}} u^0, t^{\frac{1}{q}} v^0) &= \left( \frac{1}{p} + \frac{1}{q} \right) t - \int_{\Omega} F(t^{\frac{1}{p}} u^0, t^{\frac{1}{q}} v^0) dx \\ &\leq \left( \frac{1}{p} + \frac{1}{q} - \int_{\Omega} F(u^0, v^0) dx \right) t \\ &\quad - c_2 \left[ \frac{p}{\mu - p} (t^{\frac{\mu}{p}} - t) \|u^0\|_{\mu}^{\mu} + \frac{q}{\nu - q} (t^{\frac{\nu}{q}} - t) \|v^0\|_{\nu}^{\nu} \right]. \end{aligned}$$

Therefore,  $E_G(t^{\frac{1}{p}}u^0, t^{\frac{1}{q}}v^0) \rightarrow -\infty$  as  $t \rightarrow \infty$  (recall that  $\mu > p$  and  $\nu > q$ ). Choosing  $t = t_0$  large enough and denoting by  $e_p = t_0^{\frac{1}{p}}u_p^0$  and  $e_q = t_0^{\frac{1}{q}}v_q^0$ , we are led to (ii). This completes the proof.  $\square$

**Example 4.1** Let  $p = 3/2$ ,  $q = 9/4$ ,  $\Omega = ]a, b[ \times \mathbb{R}^2$  ( $a < b$ ) and

$$F(u, v) = u^2 + |v|^{7/2} + 1/4\{|u|^{5/2} + |v|^{5/2}\}.$$

The conditions **(F1)** and **(F4)** can be verified easily. Choosing  $r = 5/2$ ,  $s = 15/4$  and  $\mu = 5/2$ ,  $\nu = 7/2$ , the assumptions **(F2)** and **(F3)** hold too. Therefore we can apply Theorem 4.1 and obtain at least one nonzero solution for problem  $(S_{3/2, 9/4})$ .

#### 4.4 Systems with sublinear potential

In the previous section we studied problem  $(S_{p,q})$ , assuming that the function  $F$  satisfies the condition **(F3)**, which is an Ambrosetti-Rabinowitz type condition and which asserts that the energy functional  $E$  satisfies the Palais-Smale or Cerami compactness condition. This condition implies, in particular, some sort of *superlinearity* of  $F$ . In this section we will treat the case when  $F$  is *sub-(p, q)-linear*. We suppose that the conditions **(F1)**-**(F2)** are fulfilled as introduction. Also, we impose that the function  $F : \Omega \times \mathbb{R}^2 \rightarrow \mathbb{R}$  satisfies the following conditions:

**(F'3)**  $\lim_{u,v \rightarrow 0} \frac{F_u(x, u, v)}{|u|^{p-1}} = \lim_{u,v \rightarrow 0} \frac{F_v(x, u, v)}{|v|^{q-1}} = 0$  uniformly for every  $x \in \Omega$ .

**(F'4)** There exist  $p_1 \in (0, p)$ ,  $q_1 \in (0, q)$ ,  $\mu \in [p, p^*]$ ,  $\nu \in [q, q^*]$  and  $a \in L^{\mu/(\mu-p_1)}(\Omega)$ ,  $b \in L^{\nu/(\nu-q_1)}(\Omega)$ ,  $c \in L^1(\Omega)$  such that

$$F(x, u, v) \leq a(x)|u|^{p_1} + b(x)|v|^{q_1} + c(x)$$

for every  $x \in \Omega$  and  $(u, v) \in \mathbb{R}^2$ .

**(F'5)** There exists  $(u_0, v_0) \in W_G^{p,q}$  such that

$$\int_{\Omega} F(x, u_0(x), v_0(x)) dx > 0.$$

In this section we suppose that the function  $F$  satisfies the conditions **(F1)**-**(F2)** and **(F'3)**-**(F'5)**. This section is dedicated to study the problem  $(S_{p,q}^{\lambda})$  which is equivalent with, find  $(u, v) \in W_0^{1,p}(\Omega) \times W_0^{1,q}(\Omega)$  such that

$$\begin{cases} \langle A_p(u), w_1 \rangle_p - \lambda \int_{\Omega} F'_u(x; u(x), v(x)) w_1(x) dx = 0 \\ \langle A_q(v), w_2 \rangle_q - \lambda \int_{\Omega} F'_v(x; u(x), v(x)) w_2(x) dx = 0, \end{cases}$$

for every  $(w_1, w_2) \in W_0^{1,p}(\Omega) \times W_0^{1,q}(\Omega)$ , where  $\lambda \in \mathbb{R}$  is a parameter.

Using the Theorem 1.29 it is enough to study the critical points of the functional  $E_{\lambda,G} = E_{\lambda}|_{W_0^{1,p,q}(\Omega)}$ . The functional  $E_{\lambda,G} : W_0^{1,p,q}(\Omega) \rightarrow \mathbb{R}$  is given by

$$E_{\lambda,G}(u, v) = \frac{1}{p} \|u\|_{1,p}^p + \frac{1}{q} \|v\|_{1,q}^q - \lambda \mathcal{F}_G(u, v), \quad (4.25)$$

where  $\mathcal{F}_G(u, v) = \int_{\Omega} F(x; u(x), v(x)) dx$  and its critical points will be the weak solutions of the problem  $(S_{p,q}^{\lambda})$ .

The main result of this section is the following.

**Theorem 4.2** *Let  $F : \Omega \times \mathbb{R}^2 \rightarrow \mathbb{R}$  be a function which satisfies the conditions **(F1)**-**(F2)** and **(F'3)**-**(F'5)** and  $ps = qr$ . Then, there exist an open interval  $\Lambda \subset (0, \infty)$  and  $\sigma > 0$  such that for all  $\lambda \in \Lambda$  the system  $(S_{\lambda})$  has at least two distinct, nontrivial weak solutions, denoted by  $(u_{\lambda}^i, v_{\lambda}^i)$ ,  $i \in \{1, 2\}$  such that  $\|u_{\lambda}^i\|_{1,p} < \sigma$ ,  $\|v_{\lambda}^i\|_{1,q} < \sigma$ ,  $i \in \{1, 2\}$ .*

Further in this section, we suppose that all assumptions of Theorem 4.2 are fulfilled. Before we prove Theorem 4.2, the main result of this section, we need some auxiliary results.

**Lemma 4.1** *For every  $\varepsilon > 0$  there exists  $c(\varepsilon) > 0$  such that*

- i)  $|F_u(x, u, v)| \leq \varepsilon(|u|^{p-1} + |v|^{(p-1)q/p}) + c(\varepsilon)(|u|^{r-1} + |v|^{(r-1)q/p})$ ;*
- ii)  $|F_v(x, u, v)| \leq \varepsilon(|v|^{q-1} + |u|^{(q-1)p/q}) + c(\varepsilon)(|v|^{s-1} + |u|^{(s-1)p/q})$ ;*
- iii)  $|F(x, u, v)| \leq \varepsilon(|u|^p + |v|^{(p-1)q/p}|u| + |v|^q + |u|^{(q-1)p/q}|v|) + c(\varepsilon)(|u|^r + |v|^{(r-1)q/p}|u| + |v|^s + |u|^{(s-1)p/q}|v|)$*

*for every  $x \in \Omega$  and  $(u, v) \in \mathbb{R}^2$ .*

*Proof* *i)* Let  $\varepsilon > 0$  be arbitrary. Let us prove the first inequality, the second one being similar. From the first limit of **(F'3)** we have in particular that

$$\lim_{u,v \rightarrow 0} \frac{F_u(x, u, v)}{|u|^{p-1} + |v|^{(p-1)q/p}} = 0.$$

Therefore, there exists  $\delta(\varepsilon) > 0$  such that, if  $|u|^{p-1} + |v|^{(p-1)q/p} < \delta(\varepsilon)$ ,

then  $|F_u(x, u, v)| \leq \varepsilon(|u|^{p-1} + |v|^{(p-1)q/p})$ . If  $|u|^{p-1} + |v|^{(p-1)q/p} \geq \delta(\varepsilon)$ , then (4.2) implies that

$$|F_u(x, u, v)| \leq c_1[(|u|^{p-1} + |v|^{(p-1)q/p})^{(r-1)/(p-1)}\delta(\varepsilon)^{(p-r)/(p-1)} + |u|^{r-1}] + c(\varepsilon)(|u|^{r-1} + |v|^{(r-1)q/p}).$$

Combining the above inequalities, we obtain the desired relation. The inequality *iii*) follows from the mean value theorem, *i*), *ii*) and  $F(x, 0, 0) = 0$ .  $\square$

**Lemma 4.2**  $\mathcal{F}_G$  is a sequentially weakly continuous function on  $W_{0,G}^{1,p,q}(\Omega)$ .

*Proof* Suppose the contrary, that is, let  $\{(u_n, v_n)\} \subset W_{0,G}^{1,p,q}(\Omega)$  be a sequence which converges weakly to  $(u, v) \in W_{0,G}^{1,p,q}(\Omega)$  and  $\mathcal{F}_G(u_n, v_n) \not\rightarrow \mathcal{F}_G(u, v)$ . Therefore, there exists  $\varepsilon_0 > 0$  and a subsequence of  $\{(u_n, v_n)\}$  (denoted again by  $\{(u_n, v_n)\}$ ) such that

$$0 < \varepsilon_0 \leq |\mathcal{F}_G(u_n, v_n) - \mathcal{F}_G(u, v)| \quad \text{for every } n \in \mathbb{N}.$$

For some  $0 < \theta_n < 1$  we have

$$0 < \varepsilon_0 \leq |\mathcal{F}'(u_n + \theta_n(u - u_n), v_n + \theta_n(v - v_n))(u_n - u, v_n - v)| \quad (4.26)$$

for every  $n \in \mathbb{N}$ . Let us denote by  $w_n = u_n + \theta_n(u - u_n)$  and  $y_n = v_n + \theta_n(v - v_n)$ . Since the embeddings  $W_{0,G}^{1,p} \hookrightarrow L^r(\Omega)$  and  $W_{0,G}^{1,q} \hookrightarrow L^s(\Omega)$  are compact, up to a subsequence,  $\{(u_n, v_n)\}$  converges strongly to  $(u, v)$  in  $L^r(\Omega) \times L^s(\Omega)$ . By (4.6), Lemma 4.1, Hölder's inequality and  $ps = qr$  one has

$$\begin{aligned} & |\mathcal{F}'(w_n, y_n)(u_n - u, v_n - v)| \\ & \leq \int_{\Omega} [|F_u(x, w_n, y_n)||u_n - u| + |F_v(x, w_n, y_n)||v_n - v|] dx \leq \\ & \varepsilon \int_{\Omega} [(|w_n|^{p-1} + |y_n|^{(p-1)q/p})|u_n - u| + (|y_n|^{q-1} + |w_n|^{(q-1)p/q})|v_n - v|] dx \\ & + c(\varepsilon) \int_{\Omega} [(|w_n|^{r-1} + |y_n|^{(r-1)q/p})|u_n - u| + (|y_n|^{s-1} \\ & + |w_n|^{(s-1)p/q})|v_n - v|] dx \\ & \leq \varepsilon[(\|w_n\|_p^{p-1} + \|y_n\|_q^{(p-1)q/p})\|u_n - u\|_p + (\|y_n\|_q^{q-1} + \\ & \|w_n\|_p^{(q-1)p/q})\|v_n - v\|_q] + c(\varepsilon)[(\|w_n\|_r^{r-1} + \|y_n\|_s^{(r-1)q/p})\|u_n - u\|_r \\ & + (\|y_n\|_s^{s-1} + \|w_n\|_r^{(s-1)p/q})\|v_n - v\|_s]. \end{aligned}$$

Since  $\{w_n\}$  and  $\{y_n\}$  are bounded in  $W_{0,G}^{1,p} \hookrightarrow L^p(\Omega) \cap L^r(\Omega)$  and  $W_{0,G}^{1,q} \hookrightarrow L^q(\Omega) \cap L^s(\Omega)$ , respectively, while  $u_n \rightarrow u$  and  $v_n \rightarrow v$

strongly in  $L^r(\Omega)$  and  $L^s(\Omega)$ , respectively, choosing  $\varepsilon > 0$  arbitrarily small, we obtain that  $\mathcal{F}'(w_n, y_n)(u_n - u, v_n - v) \rightarrow 0$ , as  $n \rightarrow \infty$ . But this contradicts (4.26).  $\square$

**Lemma 4.3** *Let  $\lambda \geq 0$  be fixed and let  $\{(u_n, v_n)\}$  be a bounded sequence in  $W_{0,G}^{1,p,q}(\Omega)$  such that*

$$\|E'_{\lambda,G}(u_n, v_n)\|_{\star} \rightarrow 0$$

*as  $n \rightarrow \infty$ . Then  $\{(u_n, v_n)\}$  contains a strongly convergent subsequence in  $W_{0,G}^{1,p,q}(\Omega)$ .*

*Proof* Because  $W_{0,G}^{1,p,q}(\Omega)$  is a reflexive Banach space and  $\{(u_n, v_n)\}$  is a bounded sequence, we can assume that

$$(u_n, v_n) \rightarrow (u, v) \text{ weakly in } W_{0,G}^{1,p,q}; \quad (4.27)$$

$$(u_n, v_n) \rightarrow (u, v) \text{ strongly in } L^r(\Omega) \times L^s(\Omega). \quad (4.28)$$

On the other hand, we have

$$\begin{aligned} E'_{\lambda,G}(u_n, v_n)(u - u_n, v - v_n) &= \int_{\Omega} |\nabla u_n|^{p-2} \nabla u_n (\nabla u - \nabla u_n) \\ &+ \int_{\Omega} |\nabla v_n|^{q-2} \nabla v_n (\nabla v - \nabla v_n) - \lambda \mathcal{F}'_G(u_n, v_n)(u - u_n, v - v_n) \end{aligned}$$

and

$$\begin{aligned} E'_{\lambda,G}(u, v)(u_n - u, v_n - v) &= \int_{\Omega} |\nabla u|^{p-2} \nabla u (\nabla u_n - \nabla u) \\ &+ \int_{\Omega} |\nabla v|^{q-2} \nabla v (\nabla v_n - \nabla v) - \lambda \mathcal{F}'_G(u, v)(u_n - u, v_n - v). \end{aligned}$$

Adding these two relations, one has

$$\begin{aligned} a_n &\stackrel{\text{not.}}{=} \int_{\Omega} (|\nabla u_n|^{p-2} \nabla u_n - |\nabla u|^{p-2} \nabla u) (\nabla u_n - \nabla u) \\ &+ \int_{\Omega} (|\nabla v_n|^{q-2} \nabla v_n - |\nabla v|^{q-2} \nabla v) (\nabla v_n - \nabla v) \\ &= -E'_{\lambda,G}(u_n, v_n)(u - u_n, v - v_n) - E'_{\lambda,G}(u, v)(u_n - u, v_n - v) \\ &\quad - \lambda \mathcal{F}'_G(u_n, v_n)(u - u_n, v - v_n) - \lambda \mathcal{F}'_G(u, v)(u_n - u, v_n - v). \end{aligned}$$

Using (4.27) and (4.28), similar estimations as in Lemma 4.2 show that the last two terms tends to 0 as  $n \rightarrow \infty$ . Due to (4.27), the second terms tends to 0, while the inequality

$$|E'_{\lambda,G}(u_n, v_n)(u - u_n, v - v_n)| \leq \|E'_{\lambda,G}(u_n, v_n)\|_{\star} \|(u - u_n, v - v_n)\|_{1,p,q}$$

and the assumption implies that the first term tends to 0 too. Thus,

$$\lim_{n \rightarrow \infty} a_n = 0. \quad (4.29)$$

From the well-known inequality

$$|t-s|^\alpha \leq \begin{cases} (|t|^{\alpha-2}t - |s|^{\alpha-2}s)(t-s), & \text{if } \alpha \geq 2, \\ ((|t|^{\alpha-2}t - |s|^{\alpha-2}s)(t-s))^{\alpha/2}(|t|^\alpha + |s|^\alpha)^{(2-\alpha)/2}, & \text{if } 1 < \alpha < 2, \end{cases}$$

for all  $t, s \in \mathbb{R}^N$ , and (4.29), we conclude that

$$\lim_{n \rightarrow \infty} \int_{\Omega} (|\nabla u_n - \nabla u|^p + |\nabla v_n - \nabla v|^q) = 0,$$

hence, the sequence  $\{(u_n, v_n)\}$  converges strongly to  $(u, v)$  in  $W_G^{p,q}$ .  $\square$

**Proof of Theorem 4.2.** We will show that the assumptions of Theorem 1.15 are fulfilled for:  $X = W_{0,G}^{1,p,q}(\Omega)$ ,  $I = [0, \infty[$  and  $f = E_{\lambda,G}$ .

We fix  $\lambda \geq 0$ . It is clear that  $W_{0,G}^{1,p,q}(\Omega) \ni (u, v) \mapsto \frac{1}{p}\|u\|_{1,p}^p + \frac{1}{q}\|v\|_{1,q}^q$  is sequentially weakly lower semicontinuous. Thus, from Lemma 4.2 it follows that  $E_{\lambda,G}(\cdot, \cdot)$  is also sequentially weakly lower semicontinuous.

First we prove that

$$\lim_{\|(u,v)\| \rightarrow \infty} E_{\lambda,G}(u, v) = +\infty. \quad (4.30)$$

Indeed, from **(F'4)** and Hölder's inequality, one has

$$\begin{aligned} E_{\lambda,G}(u, v) &\geq \frac{1}{p}\|u\|_{1,p}^p + \frac{1}{q}\|v\|_{1,q}^q - \lambda \int_{\Omega} [a(x)|u|^{p_1} + b(x)|v|^{q_1} + c(x)] dx \\ &\geq \frac{1}{p}\|u\|_{1,p}^p + \frac{1}{q}\|v\|_{1,q}^q - \lambda[\|a\|_{\mu/(\mu-p_1)}\|u\|_{\mu}^{p_1} \\ &\quad + \|b\|_{\nu/(\nu-q_1)}\|v\|_{\nu}^{q_1} + \|c\|_1]. \end{aligned}$$

Since  $W_{0,G}^{1,p}(\Omega) \hookrightarrow L^\mu(\Omega)$  and  $W_{0,G}^{1,q}(\Omega) \hookrightarrow L^\nu(\Omega)$  are continuous,  $p_1 < p$  and  $q_1 < q$ , then relation (4.30) yields immediately.

We prove that  $E_{\lambda,G}(\cdot, \cdot)$  satisfies the Palais-Smale condition: Let  $\{(u_n, v_n)\}$  be a sequence in  $W_{0,G}^{1,p,q}(\Omega)$  such that  $\sup_{n \in \mathbb{N}} |E_{\lambda,G}(u_n, v_n)| < +\infty$  and

$\lim_{n \rightarrow \infty} \|E'_{\lambda,G}(u_n, v_n)\|_* = 0$ . According to (4.30),  $\{(u_n, v_n)\}$  must be bounded in  $W_{0,G}^{1,p,q}(\Omega)$ . The conclusion follows now by Lemma 4.3.

Now, we deal with  $(\gamma_1)$  from Theorem 1.15: Let us define the function  $g : (0, \infty) \rightarrow \mathbb{R}$  by

$$g(t) = \sup\{\mathcal{F}_G(u, v) : \frac{1}{p}\|u\|_{1,p}^p + \frac{1}{q}\|v\|_{1,q}^q \leq t\}.$$

After integration in the inequality from Lemma 4.1 *iii*), then by using the Young inequality, the fact that the embedding  $W_{0,G}^{1,p}(\Omega) \hookrightarrow L^\mu(\Omega)$  and  $W_{0,G}^{1,q}(\Omega) \hookrightarrow L^\nu(\Omega)$  are compact for  $\mu \in ]p, p^*[$  and  $\nu \in ]q, q^*[$  and the relation  $ps = qr$ , we have that for an arbitrary  $\varepsilon > 0$  there exists  $c(\varepsilon) > 0$  such that

$$\mathcal{F}_G(u, v) \leq \varepsilon(\|u\|_{1,p}^p + \|v\|_{1,q}^q) + c(\varepsilon)(\|u\|_{1,p}^r + \|v\|_{1,q}^s) \quad (4.31)$$

for every  $(u, v) \in W_{0,G}^{1,p,q}(\Omega)$ . Since the function  $x \mapsto (a^x + b^x)^{1/x}$ ,  $x > 0$  is non-increasing ( $a, b \geq 0$ ), and by using again  $ps = qr$ , one has that

$$\|u\|_{1,p}^r + \|v\|_{1,q}^s \leq [\|u\|_{1,p}^p + \|v\|_{1,q}^q]^{r/p}. \quad (4.32)$$

Therefore,

$$g(t) \leq \varepsilon \max\{p, q\}t + c(\varepsilon)(\max\{p, q\}t)^{\frac{r}{p}}, \quad t > 0.$$

On the other hand, it is clear that  $g(t) \geq 0$ ,  $t > 0$ . Taking into account the arbitrariness of  $\varepsilon > 0$  and the fact that  $r > p$ , we conclude that

$$\lim_{t \rightarrow 0^+} \frac{g(t)}{t} = 0. \quad (4.33)$$

By **(F'5)** it is clear that  $(u_0, v_0) \neq (0, 0)$  (note that  $\mathcal{F}_G(0, 0) = 0$ ). Therefore, it is possible to choose a number  $\eta$  such that

$$0 < \eta < \mathcal{F}_G(u_0, v_0) \left[ \frac{1}{p} \|u_0\|_{1,p}^p + \frac{1}{q} \|v_0\|_{1,q}^q \right]^{-1}.$$

Due to (4.33), there exists  $t_0 \in \left(0, \frac{1}{p} \|u_0\|_{1,p}^p + \frac{1}{q} \|v_0\|_{1,q}^q\right)$  such that  $g(t_0) < \eta t_0$ . Thus,  $g(t_0) < \mathcal{F}_G(u_0, v_0) t_0 \left[ \frac{1}{p} \|u_0\|_{1,p}^p + \frac{1}{q} \|v_0\|_{1,q}^q \right]^{-1}$ . Let  $\rho_0 > 0$  be such that

$$g(t_0) < \rho_0 < \mathcal{F}_G(u_0, v_0) t_0 \left[ \frac{1}{p} \|u_0\|_{1,p}^p + \frac{1}{q} \|v_0\|_{1,q}^q \right]^{-1}. \quad (4.34)$$

Define  $h : I = [0, \infty) \rightarrow \mathbb{R}$  by  $h(\lambda) = \rho_0 \lambda$ . We prove that  $h$  fulfils the inequality *iii*) from Theorem 1.15.

Due to the choice of  $t_0$  and (4.34), one has

$$\rho_0 < \mathcal{F}_G(u_0, v_0). \quad (4.35)$$

The function

$$I \ni \lambda \mapsto \inf_{(u,v) \in W_{0,G}^{1,p,q}(\Omega)} \left[ \frac{1}{p} \|u\|_{1,p}^p + \frac{1}{q} \|v\|_{1,q}^q + \lambda(\rho_0 - \mathcal{F}_G(u, v)) \right]$$

is clearly upper semicontinuous on  $I$ . Thanks to (4.35), we have

$$\begin{aligned} \lim_{\lambda \rightarrow \infty} \inf_{(u,v) \in W_{0,G}^{1,p,q}(\Omega)} (E_{\lambda,G}(u,v) + \rho_0 \lambda) &\leq \lim_{\lambda \rightarrow \infty} \left[ \frac{1}{p} \|u_0\|_{1,p}^p + \frac{1}{q} \|v_0\|_{1,q}^q \right. \\ &\quad \left. + \lambda(\rho_0 - \mathcal{F}_G(u_0, v_0)) \right] = -\infty. \end{aligned}$$

Thus we find an element  $\bar{\lambda} \in I$  such that

$$\begin{aligned} &\sup_{\lambda \in I} \inf_{(u,v) \in W_{0,G}^{1,p,q}(\Omega)} (E_{\lambda,G}(u,v) + \rho_0 \lambda) \quad (4.36) \\ &= \inf_{(u,v) \in W_{0,G}^{1,p,q}(\Omega)} \left[ \frac{1}{p} \|u\|_{1,p}^p + \frac{1}{q} \|v\|_{1,q}^q + \bar{\lambda}(\rho_0 - \mathcal{F}_G(u,v)) \right]. \end{aligned}$$

Since  $g(t_0) < \rho_0$ , for all  $(u,v) \in W_{0,G}^{1,p,q}(\Omega)$  such that  $\frac{1}{p} \|u\|_{1,p}^p + \frac{1}{q} \|v\|_{1,q}^q \leq t_0$ , we have  $\mathcal{F}(u,v) < \rho_0$ . Thus,

$$t_0 \leq \inf \left\{ \frac{1}{p} \|u\|_{1,p}^p + \frac{1}{q} \|v\|_{1,q}^q : \mathcal{F}_G(u,v) \geq \rho_0 \right\}. \quad (4.37)$$

On the other hand,

$$\begin{aligned} \inf_{(u,v) \in W_{0,G}^{1,p,q}(\Omega)} \sup_{\lambda \in I} (E_{\lambda,G}(u,v) + \rho_0 \lambda) &= \inf_{(u,v) \in W_{0,G}^{1,p,q}(\Omega)} \left[ \frac{1}{p} \|u\|_{1,p}^p + \frac{1}{q} \|v\|_{1,q}^q \right. \\ &\quad \left. + \sup_{\lambda \in I} (\lambda(\rho_0 - \mathcal{F}_G(u,v))) \right] = \inf \left\{ \frac{1}{p} \|u\|_{1,p}^p + \frac{1}{q} \|v\|_{1,q}^q : \mathcal{F}_G(u,v) \geq \rho_0 \right\}. \end{aligned}$$

Thus, (4.37) is equivalent to

$$t_0 \leq \inf_{(u,v) \in W_{0,G}^{1,p,q}(\Omega)} \sup_{\lambda \in I} (E_{\lambda,G}(u,v) + \rho_0 \lambda). \quad (4.38)$$

There are two distinct cases:

I) If  $0 \leq \bar{\lambda} < t_0/\rho_0$ , we have

$$\inf_{(u,v) \in W_{0,G}^{1,p,q}(\Omega)} \left[ \frac{1}{p} \|u\|_{1,p}^p + \frac{1}{q} \|v\|_{1,q}^q + \bar{\lambda}(\rho_0 - \mathcal{F}_G(u,v)) \right]$$

$$\leq E_{\bar{\lambda},G}(0,0) + \rho_0 \bar{\lambda} = \bar{\lambda} \rho_0 < t_0.$$

Combining the above inequality with (4.36) and (4.38), the desired relation from Theorem 1.15 ( $\gamma_1$ ) yields immediately.

II) If  $t_0/\rho_0 \leq \bar{\lambda}$ , from (4.35) and (4.34) we obtain

$$\begin{aligned} & \inf_{(u,v) \in W_{0,G}^{1,p,q}(\Omega)} \left[ \frac{1}{p} \|u\|_{1,p}^p + \frac{1}{q} \|v\|_{1,q}^q + \bar{\lambda}(\rho_0 - \mathcal{F}_G(u,v)) \right] \\ & \leq \frac{1}{p} \|u_0\|_{1,p}^p + \frac{1}{q} \|v_0\|_{1,q}^q + \bar{\lambda}(\rho_0 - \mathcal{F}_G(u_0, v_0)) \\ & \leq \frac{1}{p} \|u_0\|_{1,p}^p + \frac{1}{q} \|v_0\|_{1,q}^q + \frac{t_0}{\rho_0}(\rho_0 - \mathcal{F}_G(u_0, v_0)) < t_0. \end{aligned}$$

The conclusion holds similarly as in the first case.

Thus, the hypotheses of Theorem 1.15 are fulfilled. This implies the existence of an open interval  $\Lambda \subset [0, \infty)$  and  $\sigma > 0$  such that for all  $\lambda \in \Lambda$  the function  $E_{\lambda,G}(\cdot, \cdot)$  has at least three distinct critical points in  $W_{0,G}^{1,p,q}(\Omega)$  (denote them by  $(u_\lambda^i, v_\lambda^i)$ ,  $i \in \{1, 2, 3\}$ ) and  $\|(u_\lambda^i, v_\lambda^i)\| < \sigma$ . Therefore  $\|u_\lambda^i\|_{1,p} < \sigma$ ,  $\|v_\lambda^i\|_{1,q} < \sigma$ ,  $i \in \{1, 2, 3\}$ .

**Example 4.2** Let  $\Omega = \omega \times \mathbb{R}^2$ , where  $\omega$  is a bounded open interval in  $\mathbb{R}$ . Let  $\gamma : \Omega \rightarrow \mathbb{R}$  be a continuous, non-negative, not identically zero, axially symmetric function with compact support in  $\Omega$ . Then, there exist an open interval  $\Lambda \subset (0, \infty)$  and a number  $\sigma > 0$  such that for every  $\lambda \in \Lambda$ , the system

$$\begin{cases} -\Delta_{3/2} u = 5/2 \lambda \gamma(x) |u|^{1/2} u \cos(|u|^{5/2} + |v|^3) & \text{in } \Omega \\ -\Delta_{9/4} v = 3 \lambda \gamma(x) |v| v \cos(|u|^{5/2} + |v|^3) & \text{in } \Omega \\ u = v = 0 & \text{on } \partial\Omega, \end{cases}$$

has at least two distinct, nontrivial weak solutions with the properties from Theorem 4.2.

Indeed, let us choose  $F(x, u, v) = \gamma(x) \sin(|u|^{5/2} + |v|^3)$ ,  $r = 11/4$ ,  $s = 33/8$ . **(F1)** and **(F3)** hold immediately. For **(F4)** we choose  $a = b = 0$ ,  $c = \gamma$ . Since  $\gamma$  is an axially symmetric function,  $\text{supp} \gamma$  will be an  $id \times \mathbf{O}(2)$ -invariant set, that is, if  $(x, y) \in \text{supp} \gamma$ , then  $(x, gy) \in \text{supp} \gamma$  for every  $g \in \mathbf{O}(2)$ . Therefore, it is possible to fix an element  $u_0 \in W_{0, id \times \mathbf{O}(2)}^{1,3/2}(\Omega)$  such that  $u_0(x) = (\pi/2)^{2/5}$  for every  $x \in \text{supp} \gamma$ . Choosing  $v_0 = 0$ , one has that

$$\int_{\Omega} F(x, u_0(x), v_0(x)) dx = \int_{\text{supp} \gamma} \gamma(x) \sin |u_0(x)|^{5/2} dx = \int_{\text{supp} \gamma} \gamma(x) dx > 0.$$

The conclusion follows from Theorem 4.2.  $\square$

### 4.5 Shift solutions for gradient systems

In the above two section we was supposed that the function  $F : \Omega \times \mathbb{R}^2 \rightarrow \mathbb{R}$  is *superlinear* in  $(0, 0)$ . If  $F$  is not superlinear in  $(0, 0)$  then the problem  $(S_{p,q}^\lambda)$  cannot be handled with minimax type results. Using recent ideas of Ricceri [261] it is possible to prove a multiplicity result obtaining shift solutions for problem  $(S_{p,q}^\lambda)$ . Roughly speaking, under certain assumptions on the nonlinear term  $F$  we are able to show that the existence of a real parameter  $\lambda > 0$ , and of a pair  $(u_0, v_0) \in W_0^{1,p}(\Omega) \times W_0^{1,q}(\Omega)$  such that the problem

$$((S_{(u_0, v_0)}^\lambda)) : \begin{cases} -\Delta_p(\|u\|_{1,p}^{p'(q-1)}u) = \lambda F_u(x, u + u_0, v + v_0) \\ -\Delta_q(\|u\|_{1,q}^{q'(p-1)}u) = \lambda F_v(x, u + u_0, v + v_0), \end{cases}$$

In this section we suppose that the function  $F$  has the following form  $F(x, u, v) = b(x)G(u, v)$ , where  $G : \mathbb{R}^2 \rightarrow \mathbb{R}$  is of class  $C^1$  and  $b : \Omega \rightarrow \mathbb{R}$  is such that they fulfil the following conditions:

(G)  $G(0, 0) = 0$  and there exist real numbers  $k > 0$ ,  $p_1 \in ]0, p - 1[$ , and  $q_1 \in ]0, q - 1[$  with the following properties:

- (i) For every  $u, v \in \mathbb{R}$  we have  $|G'_u(u, v)| \leq k|u|^{p_1}$ ;
- (ii) For every  $u, v \in \mathbb{R}$  we have  $|G'_v(u, v)| \leq k|v|^{q_1}$ .

(b)  $b : \Omega \rightarrow [0, +\infty[$  belongs to  $L^1(\Omega) \cap L^\infty(\Omega)$  and is not identically zero and  $G = id^m \times O(N - m)$  invariant.

The problem  $(S_{(u_0, v_0)}^\lambda)$  can be reformulated in the following way: For  $(u_0, v_0) \in W_0^{1,p}(\Omega) \times W_0^{1,q}(\Omega)$  and  $\lambda > 0$  we denote by  $(S_{(u_0, v_0)}^\lambda)$  the problem of finding  $(u, v) \in W_0^{1,p}(\Omega) \times W_0^{1,q}(\Omega)$  such that for every  $(w_1, w_2) \in W_0^{1,p}(\Omega) \times W_0^{1,q}(\Omega)$  we have:

$$\begin{cases} \|u - u_0\|_{1,p}^{pq-p} \langle A_p(u - u_0), w_1 \rangle_p - \lambda \int_{\Omega} b(x) G'_u(u(x), v(x)) w_1(x) dx = 0 \\ \|v - v_0\|_{1,q}^{pq-q} \langle A_q(v - v_0), w_2 \rangle_q - \lambda \int_{\Omega} b(x) G'_v(u(x), v(x)) w_2(x) dx = 0. \end{cases}$$

In this case, the energy functional  $E_{(u_0, v_0), \lambda} : W_0^{1,p}(\Omega) \times W_0^{1,q}(\Omega) \rightarrow \mathbb{R}$  associated to the problem  $(S_{(u_0, v_0)}^\lambda)$  is given by:

$$E_{(u_0, v_0), \lambda}(u, v) = \frac{\|u - u_0\|_{1,p}^{pq} + \|v - v_0\|_{1,q}^{pq}}{p q} - \lambda J(u, v), \quad (4.39)$$

where  $J: W_{0,G}^{1,p,q}(\Omega) \rightarrow \mathbb{R}$  is defined by

$$J(u, v) = \int_{\Omega} b(x)G(u(x), v(x))dx.$$

In standard way from the conditions **(G)** and **(b)** it follows that the function  $E_{(u_0, v_0), \lambda}$  is of class  $C^1$  and its differential is given by:

$$\begin{aligned} E'_{(u_0, v_0), \lambda}(u, v)(w_1, w_2) &= \|u - u_0\|_{1,p}^{pq-p} \langle A_p(u - u_0), w_1 \rangle_p + (4.40) \\ &+ \|v - v_0\|_{1,q}^{pq-q} \langle A_q(v - v_0), w_2 \rangle_q - \lambda J'(u, v)(w_1, w_2), \end{aligned}$$

where

$$J'(u, v)(w_1, w_2) = \int_{\Omega} [b(x)G'_u(u, v)w_1(x) + b(x)G'_v(u, v)w_2(x)]dx, \quad (4.41)$$

for all  $(w_1, w_2) \in W_0^{1,p}(\Omega) \times W_0^{1,p}(\Omega)$ . The critical points of the functional  $E_{(u_0, v_0), \lambda}$  are the solutions of problem  $((S_{(u_0, v_0)}^\lambda))$ .

In the sequel we will use the following notations:

- (N1)**  $\|\cdot\|: W_0^{1,p}(\Omega) \times W_0^{1,q}(\Omega) \rightarrow \mathbb{R}$  denotes the homogenized Minkowski type norm defined by  $\|(u, v)\| = (\|u\|_{1,p}^{pq} + \|v\|_{1,q}^{pq})^{\frac{1}{pq}}$ , for every  $(u, v) \in W_0^{1,p}(\Omega) \times W_0^{1,q}(\Omega)$ . Throughout in this section the space  $W_0^{1,p}(\Omega) \times W_0^{1,q}(\Omega)$  is considered to be endowed with this norm. We have that the space  $W_0^{1,p}(\Omega) \times W_0^{1,q}(\Omega)$  is a separable, uniformly convex, smooth real Banach space. Note that the following inequality holds for every  $(u, v) \in W_0^{1,p}(\Omega) \times W_0^{1,q}(\Omega)$

$$\|u\|_{1,p} + \|v\|_{1,q} \leq 2\|(u, v)\|. \quad (4.42)$$

- (N2)** According to Theorem B.7, for every  $r \in ]p, p^*[$  and  $s \in ]q, q^*[$  the embedding  $W_0^{1,p}(\Omega) \hookrightarrow L^r(\Omega)$  and  $W_0^{1,q}(\Omega) \hookrightarrow L^s(\Omega)$  are compact. Therefore for the fixed numbers  $r \in [p, p^*]$  and  $s \in [q, q^*]$ , there exists a positive real number  $c$  such that the following inequalities hold for every  $(u, v) \in W_0^{1,p}(\Omega) \times W_0^{1,q}(\Omega)$

$$\|u\|_r \leq c\|u\|_{1,p}, \quad \|v\|_s \leq c\|v\|_{1,q}. \quad (4.43)$$

- (N3)**  $\nu_1 := \frac{r}{r-(p_1+1)}$ ,  $\nu_2 := \frac{s}{s-(q_1+1)}$ .

- (N4)** From condition **(G)** it follows the inequality:

$$|F(x, y) - F(\bar{x}, \bar{y})| \leq k(|x| + |\bar{x}|)^{p_1}|x - \bar{x}| + k(|y| + |\bar{y}|)^{q_1}|y - \bar{y}|, \quad (4.44)$$

for every  $(x, y), (\bar{x}, \bar{y}) \in \mathbb{R} \times \mathbb{R}$ .

In the next as in above two section we denote with  $W_{0,G}^{1,p,q}(\Omega)$  the fixed points set of the action  $G = id^m \times O(N - m)$  on  $W_0^{1,p}(\Omega) \times W_0^{1,q}(\Omega)$ . The norms  $\|\cdot\|_{1,p}$ ,  $\|\cdot\|_{1,q}$  and  $\|(\cdot, \cdot)\|$  restricted to  $W_{0,G}^{1,p,q}(\Omega)$  we will denote in the same way and we introduce the following notations  $J_G = J|_{W_{0,G}^{1,p,q}(\Omega)}$ ,  $E_{(u_0, v_0), G} = E_{(u_0, v_0), \lambda}|_{W_{0,G}^{1,p,q}(\Omega)}$ .

**Proposition 4.3** *The function  $J_G: W_{0,G}^{1,p,q}(\Omega) \rightarrow \mathbb{R}$  has the following properties:*

(a) *There exists a positive real number  $C$  such that the inequality*

$$|J_G(u, v)| \leq C(\|u\|_{1,p}^{p_1+1} + \|v\|_{1,q}^{q_1+1})$$

*holds for every pair  $(u, v) \in W_{0,G}^{1,p,q}(\Omega)$ .*

(b)  *$J_G$  is sequentially weakly continuous.*

(c) *For every  $(u, v), (w_1, w_2) \in W_{0,G}^{1,p,q}(\Omega)$  we have*

$$|J'_G(u, v)(w_1, w_2)| \leq k(\|b\|_{\nu_1} \|u\|_r^{p_1} \|w_1\|_r + \|b\|_{\nu_2} \|v\|_s^{q_1} \|w_2\|_s). \quad (4.45)$$

*Proof* (a) From (4.44) it follows that

$$|F(u(x), v(x))| \leq k|u(x)|^{p_1+1} + k|v(x)|^{q_1+1}$$

for every  $x \in \Omega$ . Thus

$$|b(x)F(u(x), v(x))| \leq k|b(x)||u(x)|^{p_1+1} + k|b(x)||v(x)|^{q_1+1}. \quad (4.46)$$

Using (4.46) and Hölder's inequality, we get

$$|J_G(u, v)| \leq k\|b\|_{\nu_1} \|u\|_r^{p_1+1} + k\|b\|_{\nu_2} \|v\|_s^{q_1+1}.$$

Hence, in view of (4.43), we obtain

$$|J_G(u, v)| \leq k\|b\|_{\nu_1} c^{p_1+1} \|u\|_{1,p}^{p_1+1} + k\|b\|_{\nu_2} c^{q_1+1} \|v\|_{1,q}^{q_1+1}.$$

By taking  $C = \max\{k\|b\|_{\nu_1} c^{p_1+1}, k\|b\|_{\nu_2} c^{q_1+1}\}$ , we obtain the asserted inequality.

(b) In view of (4.44), for every  $(u, v), (\bar{u}, \bar{v}) \in W_{0,G}^{1,p,q}$  the following inequality holds

$$\begin{aligned} |J_G(u, v) - J_G(\bar{u}, \bar{v})| &\leq k \int_{\Omega} b(x)(|u(x)| + |\bar{u}(x)|)^{p_1} |u(x) - \bar{u}(x)| dx \\ &\quad + k \int_{\Omega} b(x)(|v(x)| + |\bar{v}(x)|)^{q_1} |v(x) - \bar{v}(x)| dx. \end{aligned}$$

Using Hölder's inequality (note that  $\frac{1}{\nu_1} + \frac{p_1}{r} + \frac{1}{r} = 1$  and  $\frac{1}{\nu_2} + \frac{q_1}{s} + \frac{1}{s} = 1$ ), we get

$$\begin{aligned} |J_G(u, v) - J_G(\bar{u}, \bar{v})| &\leq k \|b\|_{\nu_1} \|(|u| + |\bar{u}|)\|_r^{p_1} \|u - \bar{u}\|_r \\ &\quad + k \|b\|_{\nu_2} \|(|v| + |\bar{v}|)\|_s^{q_1} \|v - \bar{v}\|_s. \end{aligned} \quad (4.47)$$

Now, let  $(u_n, v_n)_{i \in \mathbb{N}} \subset W_{0,G}^{1,p,q}(\Omega)$  be a sequence which converges weakly to  $(u, v) \in W_{0,G}^{1,p,q}(\Omega)$ .

From (4.47) we obtain that the following inequality holds for every  $n \in \mathbb{N}$

$$\begin{aligned} |J_G(u_n, v_n) - J_G(u, v)| &\leq k 2^{p_1} c^{p_1} M^{p_1} \|b\|_{\nu_1} \|u_n - u\|_r \\ &\quad + k 2^{q_1} c^{q_1} M^{q_1} \|b\|_{\nu_2} \|v_n - v\|_s. \end{aligned} \quad (4.48)$$

Because the embedding  $W_{0,G}^{1,p}(\Omega) \hookrightarrow L^r(\Omega)$  and  $W_{0,G}^{1,q}(\Omega) \hookrightarrow L^s(\Omega)$  are compact, follow that  $u_n \rightarrow u$  and  $v_n \rightarrow v$  strongly in  $L^r(\Omega)$  and  $L^s(\Omega)$  respectively. From (4.48) follows that  $\lim_{n \rightarrow \infty} J(u_n, v_n) = J(u, v)$ , which prove the assertion.

(c) Pick  $(u, v), (w_1, w_2) \in W_{0,G}^{1,p,q}(\Omega)$ . From the Mean Value Theorem and **(G)** we have:

$$|F'((u(x), v(x))(w_1(x), w_2(x)))| \leq k |u(x)|^{p_1} |w_1(x)| + k |v(x)|^{q_1} |w_2(x)|. \quad (4.49)$$

From Hölder's inequality it follows that

$$|J'_G(u, v)(w_1, w_2)| \leq k (\|b\|_{\nu_1} \|u\|_r^{p_1} \|w_1\|_r + \|b\|_{\nu_2} \|v\|_s^{q_1} \|w_2\|_s).$$

This concludes our proof.  $\square$

**Proposition 4.4** *Let  $(u_0, v_0) \in W_{0,G}^{1,p,q}(\Omega)$  and  $\lambda > 0$ . The energy functional  $E_{(u_0, v_0), G}$  satisfies the following conditions:*

- (a)  $E_{(u_0, v_0), G}$  is coercive.
- (b)  $E_{(u_0, v_0), G}$  verifies the Palais-Smale condition.
- (c)  $E_{(u_0, v_0), G}$  is weakly lower semicontinuous.

*Proof* (a) According to assertion (a) of Proposition 4.3, the following inequality holds for every  $(u, v) \in W_{0,G}^{1,p,q}(\Omega)$

$$\frac{\|u - u_0\|_{1,p}^{pq}}{pq} - \lambda C \|u\|_{1,p}^{p_1+1} + \frac{\|v - v_0\|_{1,q}^{pq}}{pq} - \lambda C \|v\|_{1,q}^{q_1+1} \leq E_{(u_0, v_0), G}(u, v). \quad (4.50)$$

Define  $g: W_{0,G}^{1,p}(\Omega) \rightarrow \mathbb{R}$  and  $h: W_{0,G}^{1,q}(\Omega) \rightarrow \mathbb{R}$  by

$$g(u) = \frac{\|u - u_0\|_{1,p}^{pq}}{pq} - \lambda C \|u\|_{1,p}^{p_1+1} \quad \text{and} \quad h(v) = \frac{\|v - v_0\|_{1,q}^{pq}}{pq} - \lambda C \|v\|_{1,q}^{q_1+1}.$$

Using these notations, (4.50) can be rewritten as

$$g(u) + h(v) \leq E_{(u_0, v_0), G}(u, v), \text{ for all } (u, v) \in W_{0, G}^{1, p, q}(\Omega). \quad (4.51)$$

For  $u \neq 0$  we have

$$\begin{aligned} g(u) &= \|u\|_{1, p}^p \left( \frac{\|u - u_0\|_{1, p}^{pq}}{pq \|u\|_{1, p}^p} - \lambda C \|u\|_{1, p}^{p_1 + 1 - p} \right) \\ &= \|u\|_{1, p}^p \left( \frac{\|u - u_0\|_{1, p}^p}{pq \|u\|_{1, p}^p} \|u - u_0\|_{1, p}^{pq - p} - \lambda C \|u\|_{1, p}^{p_1 + 1 - p} \right). \end{aligned}$$

Since  $\lim_{\|u\|_{1, p} \rightarrow \infty} \frac{\|u - u_0\|_{1, p}^p}{\|u\|_{1, p}^p} = 1$ ,  $\lim_{\|u\|_{1, p} \rightarrow \infty} \|u - u_0\|_{1, p}^{pq - p} = +\infty$  (because  $pq - p > 0$ ), and  $\lim_{\|u\|_{1, p} \rightarrow \infty} \|u\|_{1, p}^{p_1 + 1 - p} = 0$  (recall that  $p_1 < p - 1$ ), we obtain that  $\lim_{\|u\|_{1, p} \rightarrow \infty} g(u) = +\infty$ . Hence,  $g$  is coercive. A similar argument yields that  $h$  is coercive. Since  $g$  and  $h$  are continuous, it follows from (4.51) that  $E_{(u_0, v_0), G}$  is coercive.

(b) Let  $\{(u_n, v_n)\}$  be a sequence (PS) sequence in  $W_{0, G}^{1, p, q}(\Omega)$  for the function  $E_{(u_0, v_0), G}$ , that is

(PS<sub>1</sub>)  $E_{(u_0, v_0), G}(u_n, v_n)$  is bounded,

(PS<sub>2</sub>)  $E'_{(u_0, v_0), G}(u_n, v_n) \rightarrow 0$ .

From the coercivity of the function  $E_{(u_0, v_0), G}$  follows that the sequence  $\{(u_n, v_n)\}$  is bounded in  $W_{0, G}^{1, p, q}(\Omega)$ . As in Lemma 4.3 with some mirror modifications we obtain that the function  $E_{(u_0, v_0), G}$  satisfies the (PS) condition.

(c) The map  $(u, v) \in W_{0, G}^{1, p, q}(\Omega) \mapsto \frac{\|(u, v) - (u_0, v_0)\|^{pq}}{pq} \in \mathbb{R}$  is weakly lower semicontinuous, since all its lower level sets are weakly closed (recall that  $W_{0, G}^{1, p, q}(\Omega)$  is reflexive, since it is uniformly convex). The map  $(u, v) \in W_{0, G}^{1, p, q}(\Omega) \mapsto -\lambda J_G(u, v) \in \mathbb{R}$  is sequentially weakly continuous (according to assertion (b) of Proposition 4.3), hence the map  $E_{(u_0, v_0), G}$  is sequentially weakly lower semicontinuous. Since  $E_{(u_0, v_0), G}$  is coercive, the Eberlein-Smulyan theorem implies that  $E_{(u_0, v_0), G}$  is weakly lower semicontinuous.  $\square$

The main result of this section is the following.

**Theorem 4.3** *Assume that the hypotheses **(G)** and **(b)** hold, and that the function  $J_G$  is not constant. Then, for every  $\sigma \in ] \inf_{W_{0, G}^{1, p, q}(\Omega)} J_G, \sup_{W_{0, G}^{1, p, q}(\Omega)} J_G[$*

and every  $(u_0, v_0) \in J^{-1}(]-\infty, \sigma])$ , one of the following alternatives is true:

- (A<sub>1</sub>) There exists  $\lambda > 0$  such that the function  $E_{(u_0, v_0), G}$  has at least three critical points in  $W_{0, G}^{1, p, q}(\Omega)$ .
- (A<sub>2</sub>) There exists  $(u^*, v^*) \in J_G^{-1}(\sigma)$  such that, for all  $(u, v) \in J_G^{-1}(] \sigma, +\infty[) \setminus \{(u^*, v^*)\}$ , the inequality

$$\|(u, v) - (u_0, v_0)\| > \|(u^*, v^*) - (u_0, v_0)\|$$

holds.

*Proof* Fix  $\sigma \in ]-\infty, \sigma[$ ,  $\inf_{W_{0, G}^{1, p, q}(\Omega)} J_G$ ,  $\sup_{W_{0, G}^{1, p, q}(\Omega)} J_G$ ,  $(u_0, v_0) \in J^{-1}(]-\infty, \sigma])$ , and

assume that (A<sub>1</sub>) does not hold. This implies that for every  $\lambda > 0$  the function  $E_{(u_0, v_0), G}$  has at most one local minimum in  $W_{0, G}^{1, p, q}(\Omega)$ . Indeed, the existence of two local minima of  $E_{(u_0, v_0), G}$  would imply, by the mountain pass theorem of zero altitude 1.8 (recall that, according to assertion (b) of Proposition 4.4, the function  $E_{(u_0, v_0), G}$  satisfies the Palais-Smale condition) that this function has a third critical point. This would contradict our assumption that (A<sub>1</sub>) does not hold.

We are going to prove that in this case (A<sub>2</sub>) is satisfied. For this purpose we apply Theorem 1.19.

Set  $\Lambda := [0, +\infty[$  and define the map  $f: W_{0, G}^{1, p, q}(\Omega) \times \Lambda \rightarrow \mathbb{R}$  by

$$f(u, v, \lambda) = E_{(u_0, v_0), G}(u, v) + \lambda \sigma = \frac{\|(u, v) - (u_0, v_0)\|^{pq}}{pq} + \lambda(\sigma - J_G(u, v)).$$

We consider the space  $W_{0, G}^{1, p, q}(\Omega)$  be equipped with the weak topology. Our aim is to show that  $f$  satisfies the conditions (i), (ii), and (iii) of Theorem 1.19. Observe that for every  $(u, v) \in W_{0, G}^{1, p, q}(\Omega)$  the map  $f(u, v, \cdot): \Lambda \rightarrow \mathbb{R}$  is affine, thus it is quasi-concave and continuous, that is, condition (i) is fulfilled. Next we prove that condition (ii) is also satisfied. For this fix a real number  $\lambda \geq 0$ . By assertion (c) of Proposition 4.4 we know that  $f(\cdot, \lambda)$  is (weakly) lower semicontinuous. Assume now that  $f(\cdot, \lambda)$  has a local minimum which is not global. Since  $f(\cdot, \lambda)$  is coercive (by assertion (a) of Proposition 4.4) and weakly lower semicontinuous, it has a global minimum. Thus,  $f(\cdot, \lambda)$  has at least two local minima. It follows that  $E_{(u_0, v_0), G}$  has at least two local minima, too, which is impossible by the assumption and the remark made at the beginning of the proof. We conclude that condition (ii) holds. To prove that condition (iii) is also satisfied we show first that  $\sup_{\Lambda} \inf_{W_{0, G}^{1, p, q}(\Omega)} f(u, v, \lambda) < +\infty$ . For

this choose  $(u_1, v_1) \in W_{0,G}^{1,p,q}(\Omega)$  such that  $\sigma < J(u_1, v_1)$ . For every  $\lambda \in \Lambda$  we have

$$\inf_{W_{0,G}^{1,p,q}(\Omega)} f(u, v, \lambda) \leq f(u_1, v_1, \lambda) \leq \frac{\|(u_1, v_1) - (u_0, v_0)\|^{pq}}{pq}, \quad (4.52)$$

hence  $\sup_{\Lambda} \inf_{W_{0,G}^{1,p,q}(\Omega)} f(u, v, \lambda) < +\infty$ . For every  $\rho_0 > \sup_{\Lambda} \inf_{W_{0,G}^{1,p,q}(\Omega)} f(u, v, \lambda)$

the set  $\{(u, v) \in W_{0,G}^{1,p,q}(\Omega) \mid f(u, v, 0) \leq \rho_0\}$  is weakly compact, thus condition (iii) is satisfied. Applying Theorem 1.19, we obtain

$$\alpha := \sup_{\Lambda} \inf_{W_{0,G}^{1,p,q}(\Omega)} f = \inf_{W_{0,G}^{1,p,q}(\Omega)} \sup_{\Lambda} f. \quad (4.53)$$

Note that the function  $\lambda \in \Lambda \mapsto \inf_{W_{0,G}^{1,p,q}(\Omega)} f(u, v, \lambda)$  is upper semicontinuous,

since for every  $(u, v) \in W_{0,G}^{1,p,q}(\Omega)$  the map  $f(u, v, \cdot)$  is continuous. Also, the first inequality of (4.52) yields that

$$\lim_{\lambda \rightarrow +\infty} \inf_{W_{0,G}^{1,p,q}(\Omega)} f(u, v, \lambda) = -\infty.$$

Thus, the map  $\lambda \in \Lambda \mapsto \inf_{W_{0,G}^{1,p,q}(\Omega)} f(u, v, \lambda)$  is upper bounded and attains its maximum at a point  $\lambda^* \in \Lambda$ . Hence,

$$\alpha = \sup_{\Lambda} \inf_{W_{0,G}^{1,p,q}(\Omega)} f = \inf_{W_{0,G}^{1,p,q}(\Omega)} \left( \frac{\|(u, v) - (u_0, v_0)\|^{pq}}{pq} + \lambda^*(\sigma - J_G(u, v)) \right). \quad (4.54)$$

On the other hand, we have for every  $(u, v) \in W_{0,G}^{1,p,q}(\Omega)$  that

$$\sup_{\Lambda} f(u, v, \lambda) = \begin{cases} +\infty, & \text{if } \sigma > J_G(u, v) \\ \frac{\|(u, v) - (u_0, v_0)\|^{pq}}{pq}, & \text{if } \sigma \leq J_G(u, v). \end{cases}$$

In view of (4.53) it follows that

$$\alpha = \inf_{J_G^{-1}([\sigma, +\infty[)} \left( \frac{\|(u, v) - (u_0, v_0)\|^{pq}}{pq} \right).$$

Since the map  $(u, v) \in W_{0,G}^{1,p,q}(\Omega) \mapsto \frac{\|(u, v) - (u_0, v_0)\|^{pq}}{pq} \in \mathbb{R}$  is weakly lower semicontinuous and coercive, and since the set  $J_G^{-1}([\sigma, +\infty[)$  is weakly closed and nonempty, there exists a pair  $(u^*, v^*) \in J_G^{-1}([\sigma, +\infty[)$  such that

$$\alpha = \frac{\|(u^*, v^*) - (u_0, v_0)\|^{pq}}{pq}. \quad (4.55)$$

Observe that  $\lambda^* > 0$ . Otherwise,  $\lambda^* = 0$ , which implies  $\alpha = 0$ , hence  $(u_0, v_0) = (u^*, v^*)$ , which is impossible since  $J_G(u_0, v_0) < \sigma$ .

The relations (4.54) and (4.55) yield that

$$\frac{\|(u^*, v^*) - (u_0, v_0)\|^{pq}}{pq} \leq \frac{\|(u^*, v^*) - (u_0, v_0)\|^{pq}}{pq} + \lambda^*(\sigma - J_G(u^*, v^*)),$$

thus  $J_G(u^*, v^*) = \sigma$ . This implies (by (4.54)) that  $(u^*, v^*)$  is a global minimum of  $E_{(u_0, v_0), G}$ . By the assumption and the remark at the beginning of the proof this pair is the only global minimum of  $E_{(u_0, v_0), G}$ . Thus, for every pair  $(u, v) \in J_G^{-1}([\sigma, +\infty]) \setminus \{(u^*, v^*)\}$ , the inequality  $\|(u, v) - (u_0, v_0)\| > \|(u^*, v^*) - (u_0, v_0)\|$  holds. We conclude that  $(A_2)$  is satisfied.  $\square$

**Corollary 4.1** *Assume that the hypotheses of Theorem 4.3 are fulfilled. If  $S$  is a convex dense subset of  $W_{0,G}^{1,p,q}(\Omega)$ , and if there exists*

$$\sigma \in ] \inf_{W_{0,G}^{1,p,q}(\Omega)} J_G, \sup_{W_{0,G}^{1,p,q}(\Omega)} J_G[$$

*such that the level set  $J_G^{-1}([\sigma, +\infty])$  is not convex, then there exist  $(u_0, v_0) \in J_G^{-1}(]-\infty, \sigma]) \cap S$  and  $\lambda > 0$  such that the function  $E_{(u_0, v_0), G}$  has at least three critical points in  $W_{0,G}^{1,p,q}(\Omega)$ , that is, there exist  $(u_0, v_0) \in J_G^{-1}(]-\infty, \sigma]) \cap S$  and  $\lambda > 0$  such that problem  $((S_{(u_0, v_0)}^\lambda))$  has at least three solutions in  $W_{0,G}^{1,p,q}(\Omega)$ .*

*Proof* Since  $J_G$  is sequentially weakly continuous (by assertion (c) of Proposition 4.3), the level set  $M := J^{-1}([\sigma, +\infty])$  is sequentially weakly closed. Theorem A.1 yields the existence of pairwise distinct pairs  $(u_0, v_0) \in S$ ,  $(u_1, v_1), (u_2, v_2) \in M$  such that

$$\|(u_1, v_1) - (u_0, v_0)\| = \|(u_2, v_2) - (u_0, v_0)\| = \inf_{(u,v) \in M} \|(u, v) - (u_0, v_0)\|.$$

It follows that  $(u_0, v_0) \notin M$ , that is,  $J_G(u_0, v_0) < \sigma$ . Also, the above relations show that alternative  $(A_2)$  of Theorem 4.3 does not hold. Hence  $(A_1)$  must be satisfied, that is, there exists  $\lambda > 0$  such that the function  $E_{(u_0, v_0), G}$  has at least three critical points in  $W_{0,G}^{1,p,q}(\Omega)$ . Using Theorem 1.29, we conclude that problem  $((S_{(u_0, v_0)}^\lambda))$  has at least three solutions in  $W_0^{1,p}(\Omega) \times W_0^{1,p}(\Omega)$ .  $\square$

**Remark.** The assumption of Corollary 4.1 stating that there exists  $\sigma \in ] \inf_{W_{0,G}^{1,p,q}(\Omega)} J, \sup_{W_{0,G}^{1,p,q}(\Omega)} J[$  such that the level set  $J^{-1}([\sigma, +\infty])$  is not convex is equivalent to the fact that  $J$  is not quasi-concave.

A direct application of Corollary 4.1 is the following.

**Theorem 4.4** *Let  $G: \mathbb{R}^2 \rightarrow [0, +\infty[$  be a function which is not quasi-concave and which satisfies condition **(G)**. Let  $b: \Omega \rightarrow [0, +\infty[$  be a  $G$ -invariant function with  $b(0) > 0$  and for which condition **(b)** holds, and assume that  $S$  is a convex dense subset of  $W_{0,G}^{1,p,q}(\Omega)$ . Then, there exist  $(u_0, v_0) \in S$  and  $\lambda > 0$  with the property that problem  $((S_{(u_0, v_0)}^\lambda))$  has at least three solutions in  $W_0^{1,p}(\Omega) \times W_0^{1,p}(\Omega)$ .*

*Proof* We are going to apply Corollary 4.1 to the spaces  $W_{0,G}^{1,p}(\Omega)$  and  $W_{0,G}^{1,q}(\Omega)$  and to the map  $J|_{W_{0,G}^{1,p,q}(\Omega)}$ . First we prove that  $J|_{W_{0,G}^{1,p,q}(\Omega)}$  is not constant. For this we assume, by contradiction, that  $J|_{W_{0,G}^{1,p,q}(\Omega)}(u, v) = J|_{W_{0,G}^{1,p,q}(\Omega)}(0, 0) = 0$  for every  $(u, v) \in W_{0,G}^{1,p,q}(\Omega)$ .

Since  $b \geq 0$  and  $F \geq 0$ , it follows that  $b(x)F(u(x), v(x)) = 0$  for every  $(u, v) \in W_{0,G}^{1,p,q}(\Omega)$  and a.e.  $x \in \Omega$ . Since  $b(0) > 0$ , there exists a real number  $R > 0$  such that the closed ball  $B$  centered in 0 with radius  $R$  is contained in  $\Omega$  and  $b(x) > 0$  for a.e.  $x \in B$ . Consequently,  $G(u(x), v(x)) = 0$  for every  $(u, v) \in W_{0,G}^{1,p,q}(\Omega)$  and a.e.  $x \in B$ . Since  $F$  is not constant, there exist  $s_0, t_0 \in \mathbb{R}$  such that  $G(s_0, t_0) > 0$ . Choosing  $u \in W_{0,G}^{1,p}(\Omega)$  and  $v \in W_{0,G}^{1,q}(\Omega)$  such that  $u(x) = s_0$  and  $v(x) = t_0$  for every  $x \in B$ , we get a contradiction.

We show that  $J|_{W_{0,G}^{1,p,q}(\Omega)}$  is not quasi-concave. It follows from above that

$$t := \int_B b(x) dx > 0.$$

The fact that  $G$  is not quasi-concave implies the existence of a real number  $\rho_0 \in ]0, \sup_{\mathbb{R} \times \mathbb{R}} G[$  such that  $G^{-1}([\rho_0, +\infty[)$  is not convex. Thus, we find  $\rho \in \mathbb{R}$ ,  $\alpha \in ]0, 1[$ , and  $(s_i, t_i) \in \mathbb{R}^2$ ,  $i \in \{1, 2, 3\}$ , with the following properties

$$(s_2, t_2) = \alpha(s_1, t_1) + (1-\alpha)(s_3, t_3), \quad G(s_1, t_1) > \rho, \quad G(s_3, t_3) > \rho, \quad G(s_2, t_2) < \rho.$$

Let

$$M := \max\{F(x, y) \mid |x| \leq \max\{|s_1|, |s_3|\}, |y| \leq \max\{|t_1|, |t_3|\}\}.$$

Choose  $R_1 > R$  and  $\varepsilon > 0$  such that

$$\|b\|_\infty M \text{meas}(A) < \varepsilon < t|F(s_i, t_i) - \rho|, \quad \text{for } i \in \{1, 2, 3\},$$

where  $A := \{x \in \Omega : R < |x| < R_1\}$  and  $\text{meas}(A)$  stays for the Lebesgue

measure of  $A$ . For  $i \in \{1, 3\}$  let  $u_i, v_i \in C_c^\infty(\Omega)$  be  $G$ -invariant functions such that  $\|u_i\|_\infty = |s_i|$ ,  $\|v_i\|_\infty = |t_i|$ ,

$$u_i(x) = \begin{cases} s_i, & |x| \leq R \\ 0, & |x| \geq R_1, \end{cases} \quad \text{and} \quad v_i(x) = \begin{cases} t_i, & |x| \leq R \\ 0, & |x| \geq R_1. \end{cases}$$

Put  $(u_2, v_2) := \alpha(u_1, v_1) + (1-\alpha)(u_3, v_3)$ . It follows that  $u_2, v_2 \in C_c^\infty(\Omega)$ ,  $\|u_2\|_\infty \leq \alpha|s_1| + (1-\alpha)|s_3|$ ,  $\|v_2\|_\infty \leq \alpha|t_1| + (1-\alpha)|t_3|$ ,

$$u_2(x) = \begin{cases} s_2, & |x| \leq R \\ 0, & |x| \geq R_1, \end{cases} \quad \text{and} \quad v_2(x) = \begin{cases} t_2, & |x| \leq R \\ 0, & |x| \geq R_1. \end{cases}$$

Let  $i \in \{1, 2, 3\}$ . Note that for every  $x \in \Omega$  the following inequalities hold

$$|u_i(x)| \leq \|u_i\|_\infty \leq \max\{|s_1|, |s_3|\} \quad \text{and} \quad |v_i(x)| \leq \|v_i\|_\infty \leq \max\{|t_1|, |t_3|\},$$

thus

$$0 \leq b(x)G(u_i(x), v_i(x)) \leq \|b\|_\infty M.$$

Since

$$\begin{aligned} J|_{W_{0,G}^{1,p,q}(\Omega)}(u_i, v_i) &= \int_B b(x)G(u_i(x), v_i(x))dx + \int_A b(x)G(u_i(x), v_i(x))dx \\ &= G(s_i, t_i)t + \int_A b(x)G(u_i(x), v_i(x))dx, \end{aligned}$$

we conclude that, for  $i \in \{1, 3\}$ ,

$$J|_{W_{0,G}^{1,p,q}(\Omega)}(u_i, v_i) \geq G(s_i, t_i)t - \|b\|_\infty M \text{meas}(A) > G(s_i, t_i)t - \epsilon > t\rho.$$

and

$$J|_{W_{0,G}^{1,p,q}(\Omega)}(u_2, v_2) \leq G(s_2, t_2)t + \|b\|_\infty M \text{meas}(A) < G(s_2, t_2)t + \epsilon < t\rho.$$

Thus  $J|_{W_{0,G}^{1,p,q}(\Omega)}$  is not quasi-concave. Therefore, Corollary 4.1 yields the existence of a pair  $(u_0, v_0) \in S$  and of a real number  $\lambda > 0$  such that the function  $E_{(u_0, v_0), G}|_{W_{0,G}^{1,p,q}(\Omega)}$  has at least three critical points in  $W_{0,G}^{1,p,q}(\Omega)$ . In order to ensure that these points are also critical points of  $E_{(u_0, v_0), G}$ , we have to show that  $E_{(u_0, v_0), G}$  is  $G$ -invariant. For this pick arbitrary  $(u, v) \in W_{0,G}^{1,p,q}(\Omega)$  and  $g \in G$ . Then,

$$\begin{aligned} E_{(u_0, v_0), G}(g \cdot u, g \cdot v) &= \frac{\|(g \cdot u, g \cdot v) - (u_0, v_0)\|^{pq}}{pq} - \lambda J(g \cdot u, g \cdot v) \\ &= \frac{\|(g \cdot u, g \cdot v) - (g \cdot u_0, g \cdot v_0)\|^{pq}}{pq} - \lambda \int_\Omega b(x)G((g \cdot u)(x), (g \cdot v)(x))dx. \end{aligned}$$

Using the formula for the change of variable, and taking into account that  $b$  is  $G$ -invariant and that the elements of  $G$  are orthogonal maps (hence the absolute value of the determinant of their matrices is 1), we conclude that

$$\begin{aligned} E_{(u_0, v_0), G}(g \cdot u, g \cdot v) &= \frac{\|(u, v) - (u_0, v_0)\|^{pq}}{pq} - \lambda \int_{\Omega} b(x)G(u(x), v(x))dx \\ &= E_{(u_0, v_0), G}(u, v). \end{aligned}$$

So, by Theorem 1.29 every critical point of  $E_{(u_0, v_0), G}$  is also a critical point of  $E_{(u_0, v_0), \lambda}$ . The conclusion follows now from Corollary 4.1.  $\square$

**Example 4.3** We give an example of a non-constant function  $F: \mathbb{R}^2 \rightarrow [0, +\infty)$  which satisfies the conditions required in the hypotheses of Theorem 4.4. Let  $\gamma, \delta \in \mathbb{R}$  be such that  $1 < \gamma < p$  and  $1 < \delta < q$ . Define  $F: \mathbb{R}^2 \rightarrow \mathbb{R}$  by

$$F(u, v) = |u|^\gamma + |v|^\delta.$$

Then,  $F$  is convex and of class  $C^1$  and satisfies the condition **(G)**. The relations

$$F(0, 0) = F\left(\frac{1}{2}(-1, 0) + \frac{1}{2}(1, 0)\right) = 0 < 1 = F(-1, 0) = F(1, 0)$$

show that  $F$  is not quasi-concave.

## 4.6 Comments and historical notes

In the last few years many paper is dedicated to study the existence and multiplicity of solutions for gradient, Hamiltonian and non-variational systems on bounded or unbounded domain. For this see the papers of de Figueiredo and his collaborators.

The motivation to investigate elliptic eigenvalue problems on such domains arises in mathematical physics, see for instance Amick [9], Amick and Toland [10]; the mathematical development was initiated by Esteban [109], Grossinho and Tersian [134], Fao and Zhao [112]. Many papers study the existence and multiplicity of solutions for elliptic systems defined on unbounded domains, see Bartsch and de Figueiredo [28], Bartsch and Wang [32], Costa [76], Dinu [92], Grossinho [133]. The main tools used in the aforementioned papers are based on mountain pass theorems and on the symmetric version of the mountain pass theorem, when the energy functional associated to problem  $(S_\lambda)$  satisfies the (PS) condition.

This chapter is based on the papers Kristály [165], [166] and Breckner and Varga [46] and we present some recent results which complete the results obtained in the aforementioned papers.

## 5

### Systems with arbitrary growth nonlinearities

The universe is not required to  
be in perfect harmony with  
human ambition.

---

Carl Sagan (1934–1996)

In this chapter we are concerned with the elliptic system

$$\begin{cases} -\Delta u = v^p & \text{in } \Omega \\ -\Delta v = f(u) & \text{in } \Omega \\ u = v = 0 & \text{on } \partial\Omega, \end{cases}$$

where  $\Omega$  is a bounded domain in  $\mathbb{R}^N$ ,  $0 < p < \frac{2}{N-2}$ , and the continuous nonlinear term  $f$  has an arbitrary growth near the origin or at infinity. Here and in the sequel, we use the notation  $s^\alpha = \text{sgn}(s) |s|^\alpha$ ,  $\alpha > 0$ .

#### 5.1 Introduction

We consider the elliptic system

$$\begin{cases} -\Delta u = g(v) & \text{in } \Omega \\ -\Delta v = f(u) & \text{in } \Omega \\ u = v = 0 & \text{on } \partial\Omega, \end{cases} \quad (S)$$

where  $\Omega \subset \mathbb{R}^N$  ( $N \geq 2$ ) is an open bounded domain with smooth boundary, and  $f, g : \mathbb{R} \rightarrow \mathbb{R}$  are continuous functions.

In the particular case when  $g(s) = s^p$ ,  $f(s) = s^q$  ( $p, q > 1$ ) and  $N \geq 3$ , system (S) has been widely studied replacing the usual criticality notion (namely,  $p, q \leq \frac{N+2}{N-2}$ ) by the so-called “critical hyperbola” which involves

both parameters  $p$  and  $q$ , that is, those pairs of points  $(p, q) \in \mathbb{R}_+^2$  which verify

$$\frac{1}{p+1} + \frac{1}{q+1} = 1 - \frac{2}{N}. \quad (CH)$$

Points  $(p, q)$  on this curve meet the typical non-compactness phenomenon of Sobolev embeddings and non-existence of solutions for  $(S)$  has been pointed out by Mitidieri [214] and van der Vorst [294] via Pohozaev-type arguments. On the other hand, when

$$1 > \frac{1}{p+1} + \frac{1}{q+1} > 1 - \frac{2}{N}, \quad (5.1)$$

the existence of nontrivial solutions for  $(S)$  has been proven by de Figueiredo and Felmer [114], Hulshof, Mitidieri and van der Vorst [143]. Note that the latter results work also for nonlinearities  $g(s) \sim s^p$  and  $f(s) \sim s^q$  as  $|s| \rightarrow \infty$  with  $(p, q)$  fulfilling (5.1). The points verifying (5.1) form a proper region in the first quadrant of the  $(p, q)$ -plane situated below the critical hyperbola  $(CH)$ . Note that (5.1) is verified for any  $p, q > 1$  whenever  $N = 2$ .

In spite of the aforementioned results, the whole region below  $(CH)$  is far to be understood from the point of view of existence/multiplicity of solutions for  $(S)$ . Via a mountain pass argument, de Figueiredo and Ruf [120] proved the existence of at least one nontrivial solution to the problem

$$\begin{cases} -\Delta u = v^p & \text{in } \Omega; \\ -\Delta v = f(u) & \text{in } \Omega; \\ u = v = 0 & \text{on } \partial\Omega, \end{cases} \quad (\tilde{S})$$

when

$$\begin{cases} 0 < p, & \text{if } N = 2; \\ 0 < p < \frac{2}{N-2}, & \text{if } N \geq 3, \end{cases} \quad (5.1)$$

and  $f : \mathbb{R} \rightarrow \mathbb{R}$  has a suitable superlinear growth at infinity, formulated in the term of the Ambrosetti-Rabinowitz condition. Later, Salvatore [270] guaranteed via the Pohozaev's fibering method the existence of a solution for  $(\tilde{S})$  in a similar context as [120]. Note that in both papers [120] and [270], *no further growth restriction* is required on the nonlinear term  $f$  other than the Ambrosetti-Rabinowitz condition. This latter fact is not surprising taking into account that (5.1) is actually equivalent to

$$1 > \frac{1}{p+1} > 1 - \frac{2}{N},$$

which is nothing but a "degenerate" case of (5.1) putting formally  $q = \infty$ , that is, the growth of  $f$  may be arbitrary large.

This chapter is divided into two parts. In Section 5.2 we are dealing with the case when  $f$  fulfills an Ambrosetti-Rabinowitz type condition. Two different approaches are presented; first, by considering the special case  $f(s) = s^q$ , the one-parametric fibering method is used in order to prove the existence of a nonzero solution to  $(\tilde{S})$ . Then, the mountain pass argument is exploited whenever  $f$  fulfills a generic Ambrosetti-Rabinowitz type condition. In Section 5.3 the nonlinear term  $f$  has an oscillatory behaviour. Moreover, the nonlinear term  $f$  may enjoy an arbitrary growth at infinity (resp., at zero) whenever it oscillates near the origin (resp., at infinity) in a suitable way. In addition, the size of our solutions reflects the oscillatory behaviour of the nonlinear term, see relations (5.11) and (5.17) below; namely, the solutions are small (resp., large) in  $L^\infty$ -norm and in a suitably chosen Sobolev space whenever the nonlinearity  $f$  oscillates near the origin (resp., at infinity). We emphasize that no symmetry condition is required on  $f$ . These results are proved by means of Ricceri's variational principle, see Theorem 1.20.

Note that system  $(\tilde{S})$  is equivalent to the Poisson equation

$$\begin{cases} -\Delta(-\Delta u)^{\frac{1}{p}} = f(u) & \text{in } \Omega; \\ u = \Delta u = 0 & \text{on } \partial\Omega. \end{cases} \quad (P)$$

The suitable functional space where solutions of  $(P)$  is going to be sought is

$$X = W^{2, \frac{p+1}{p}}(\Omega) \cap W_0^{1, \frac{p+1}{p}}(\Omega)$$

endowed with the norm

$$\|u\|_X = \left( \int_{\Omega} |\Delta u|^{\frac{p+1}{p}} dx \right)^{\frac{p}{p+1}}.$$

Due to (5.1) one has  $\frac{p+1}{p} > 1 + \frac{N-2}{2} = \frac{N}{2}$ , therefore  $W^{2, \frac{p+1}{p}}(\Omega) \subset\subset C(\Omega)$ , so

$$X \subset\subset C(\Omega). \quad (5.2)$$

For further use, we denote by  $\kappa_0 > 0$  the best embedding constant of  $X \subset\subset C(\Omega)$ . The energy functional associated to the Poisson problem  $(P)$  is  $E : X \rightarrow \mathbb{R}$  defined by

$$E(u) = \frac{p}{p+1} \|u\|_X^{\frac{p+1}{p}} - \mathcal{F}(u) \quad \text{where } \mathcal{F}(u) = \int_{\Omega} F(u(x)) dx.$$

Due to (5.2), the functional  $E$  is well-defined, is of class  $C^1$  on  $X$  and

$$\langle E'(u), w \rangle = \int_{\Omega} (-\Delta u)^{\frac{1}{p}} (-\Delta w) dx - \int_{\Omega} f(u) w dx, \quad u, w \in X.$$

Here,  $\langle \cdot, \cdot \rangle$  denotes the duality pairing between  $X$  and  $X^*$ . Note that if  $u \in X$  is a critical point of  $E$  then it is a weak solution of problem (P); in such a case, the pair  $(u, (-\Delta u)^{\frac{1}{p}}) \in X \times X$  is a weak solution of system  $(\tilde{S})$ . See also [120, Subsection 3.1] and [270, Proposition 2.1]. Moreover, standard regularity arguments show that the pair  $(u, (-\Delta u)^{\frac{1}{p}}) \in X \times X$  is actually a strong solution of system  $(\tilde{S})$ , see [120].

## 5.2 Elliptic systems with mountain pass geometry

### 5.2.1 Pure power nonlinearities: application of the fibering method

For the sake of simplicity, we consider a special case of  $(\tilde{S})$ , that is,

$$\begin{cases} -\Delta u = v^p & \text{in } \Omega; \\ -\Delta v = u^q & \text{in } \Omega; \\ u = v = 0 & \text{on } \partial\Omega, \end{cases} \quad (\bar{S})$$

where  $\Omega$  is a smooth bounded domain in  $\mathbb{R}^N$ . The main result of this section reads as follows.

**Theorem 5.1** *Assume that condition (5.1) holds and  $pq > 1$ . Then the system  $(\bar{S})$  possesses a nontrivial (strong) solution.*

In order to prove Theorem 5.1, we are going to apply the fibering method. To do this, we introduce the set

$$S = \{u \in X : \|u\|_X = 1\}.$$

For any  $v \in S$ , we consider in  $\lambda \in \mathbb{R}$  the algebraic equation

$$|\lambda|^{\frac{p+1}{p}} - |\lambda|^{q+1} \int_{\Omega} |v|^{q+1} dx = 0. \quad (5.3)$$

A direct calculation implies that (5.3) has exactly two nonzero solutions

$$\lambda_{\pm}(v) = \pm \left( \int_{\Omega} |v|^{q+1} \right)^{t_0}, \quad (5.4)$$

where  $t_0 = \left( \frac{1}{p} - q \right)^{-1}$ . Moreover,  $\lambda_{\pm}(v) \in C^1(S)$ .

We associate with  $E$  a functional  $\tilde{E}$  defined on  $\mathbb{R} \times X$  by

$$\tilde{E}(\lambda, v) = E(\lambda v) = \frac{p}{p+1} |\lambda|^{\frac{p+1}{p}} \int_{\Omega} |\Delta v|^{\frac{p+1}{p}} dx - \frac{1}{q+1} |\lambda|^{q+1} \int_{\Omega} |v|^{q+1} dx.$$

Clearly, the restriction of  $\tilde{E}$  to  $\mathbb{R} \times S$ , still denoted by  $\tilde{E}$ , becomes

$$\tilde{E}(\lambda, v) = \frac{p}{p+1} |\lambda|^{\frac{p+1}{p}} - \frac{1}{q+1} |\lambda|^{q+1} \int_{\Omega} |v|^{q+1} dx.$$

On account of Theorem 1.32, one can prove that if  $(\lambda, v) \in (\mathbb{R} \setminus \{0\}) \times S$  is a conditionally critical point of the functional  $\tilde{E}$  on  $\mathbb{R} \times S$  then the vector  $u = \lambda v$  is a nonzero critical point of the functional  $E$ , that is,  $E'(u) = 0$ . In particular, the conditionally critical point of the functional  $\tilde{E}$  implies that  $\tilde{E}'_{\lambda}(\lambda, v) = 0$ , which takes the form

$$|\lambda|^{\frac{p+1}{p}-2} \lambda - \int_{\Omega} (\lambda v)^q v dx = 0,$$

or equivalently, for  $\lambda \neq 0$ , the equation (5.3).

We finally introduce the functional  $\hat{E}_{\pm} : S \rightarrow \mathbb{R}$  by  $\hat{E}_{\pm}(v) = \tilde{E}(\lambda_{\pm}(v), v)$  which is of class  $C^1$  and on account of (5.3) and (5.4), it becomes

$$\begin{aligned} \hat{E}_{\pm}(v) &= \frac{pq-1}{(p+1)(q+1)} |\lambda_{\pm}(v)|^{q+1} \int_{\Omega} |v|^{q+1} dx \\ &= \frac{pq-1}{(p+1)(q+1)} \left( \int_{\Omega} |v|^{q+1} dx \right)^{\frac{p+1}{1-pq}} \\ &= \frac{pq-1}{(p+1)(q+1)} |\lambda_{\pm}(v)|^{\frac{p+1}{p}}. \end{aligned} \quad (5.5)$$

Note that since  $pq > 1$ , then  $\inf_S \hat{E}_{\pm} \geq 0$ .

**Proposition 5.1** *Each critical point  $v \in S$  of  $\hat{E}_{\pm}$  provides a conditionally critical point  $(\lambda_{\pm}(v), v) \in (\mathbb{R} \setminus \{0\}) \times S$  of  $\tilde{E}$ . In particular, for a minimum point  $v \in S$  of  $\hat{E}_{\pm}$  relative to  $S$ , the point  $(\lambda_{\pm}(v), v) \in (\mathbb{R} \setminus \{0\}) \times S$  is a conditionally critical point of  $\tilde{E}$ .*

*Proof* First of all, we notice that  $\tilde{E}'_{\lambda}(\lambda_{\pm}(v), v) = 0$ , see equation (5.3). On the other hand, by assumption, we have that

$$0 \in \hat{E}'_{\pm}(v) + N_S(v). \quad (5.6)$$

Therefore, a simple calculation shows that  $-\tilde{E}'_v(\lambda_{\pm}(v), v) \in N_S(v)$ . Consequently, we have that

$$-\tilde{E}'(\lambda_{\pm}(v), v) \in \{0\} \times N_S(v) = N_{(\mathbb{R} \setminus \{0\}) \times S}(\lambda_{\pm}(v), v),$$

that is,  $(\lambda_{\pm}(v), v) \in (\mathbb{R} \setminus \{0\}) \times S$  is a conditionally critical point of the functional  $\widehat{E}$ .

Now, let  $v \in S$  be a relative minimum of  $\widehat{E}_{\pm}$  to  $S$ . In particular, this fact implies that

$$\langle \widehat{E}'_{\pm}(v), w \rangle = 0 \quad \text{for all } w \in T_v(S). \quad (5.7)$$

Note that relation (5.7) is nothing but relation (5.6), see Appendix A.  $\square$

In view of Theorem 1.32 and Proposition 5.1, it is enough to find minimum points of  $\widehat{E}_{\pm}$  on  $S$ . In the sequel, we deal only with the functional  $\widehat{E}_+$ ; analogously, it is possible to consider the functional  $\widehat{E}_-$ .

**Lemma 5.1** *Assume that (5.1) holds and  $pq > 1$ . If  $\{v_k\} \subset S$  is such that the sequence  $\{\widehat{E}_+(v_k)\}$  is bounded from above, then the sequence  $\{\lambda_+(v_k)\}$  is contained in a compact interval  $[c, C]$  for some  $c, C > 0$ .*

*Proof* Let  $\{v_k\} \subset S$  be such that  $\{\widehat{E}_+(v_k)\}$  is bounded from above. For simplicity, denote by  $\{\lambda_k\}$  the sequence  $\{\lambda_+(v_k)\}$ .

We first assume by contradiction that, up to a subsequence,  $\lambda_k \rightarrow 0$  as  $k \rightarrow \infty$ . By (5.5) and  $pq > 1$ , we have that  $\int_{\Omega} |v_k|^{q+1} dx \rightarrow \infty$  as  $k \rightarrow \infty$ . Moreover, due to (5.2), we have that  $X \subset\subset L^{q+1}(\Omega)$ ; therefore,  $\|v_k\|_X \rightarrow \infty$  as  $k \rightarrow \infty$ , a contradiction.

Now, we assume by contradiction, passing to a subsequence if necessarily, that  $\lambda_k \rightarrow +\infty$  as  $k \rightarrow \infty$ . But (5.5) and  $pq > 1$  imply that  $\widehat{E}_+(v_k) \rightarrow \infty$  as  $k \rightarrow \infty$  which contradicts our assumption.  $\square$

**Proposition 5.2** *Assume that (5.1) and  $pq > 1$ . Then  $\widehat{E}_{\pm}$  attains its infimum on  $S$ .*

*Proof* We prove the statement only for  $\widehat{E}_+$ ; in a similar way, we can prove that also the functional  $\widehat{E}_-$  attains its infimum on  $S$ .

Let  $\{v_k\} \subset S$  be a sequence such that  $\widehat{E}_+(v_k) \rightarrow m_+ = \inf_S \widehat{E}_+$ . There exists  $\bar{v}_+ \in X \setminus \{0\}$  with the property  $\|\bar{v}_+\|_X \leq 1$  such that, up to a subsequence,  $v_k \rightharpoonup \bar{v}_+$  in  $X$ , and  $v_k \rightarrow \bar{v}_+$  in  $C(\Omega)$ . Due to Lemma 5.1 the corresponding sequence  $\{\lambda_+(v_k)\}$  converges, up to subsequence, to a real number  $\bar{\lambda}_+ > 0$ . Since we have

$$|\lambda_+(v_k)|^{\frac{p+1}{p}} - |\lambda_+(v_k)|^{q+1} \int_{\Omega} |v_k|^{q+1} dx = 0,$$

see equation (5.3), by passing to the limit, we have that the pair  $(\bar{\lambda}_+, \bar{v}_+)$  still solves (5.3). Thus,  $\bar{\lambda}_+ = \lambda_+(\bar{v}_+)$  with  $\bar{v}_+ \neq 0$ . In conclusion, we have

that  $\lambda_+(v_k) \rightarrow \lambda_+(\bar{v}_+)$ . Moreover, by (5.5), we conclude that

$$\widehat{E}_+(v_k) \rightarrow \widehat{E}_+(\bar{v}_+) = m_+.$$

It remains to prove that  $\|\bar{v}_+\|_X = 1$ . To see this, we argue by contradiction, namely we assume that  $\|\bar{v}_+\|_X < 1$ . For any  $t > 0$ , we deduce that

$$\frac{d}{dt} \widehat{E}_+(t\bar{v}_+) = -\frac{1}{t} (\lambda_+(t\bar{v}_+))^{\frac{p+1}{p}} < 0.$$

Therefore, the function  $t \mapsto \widehat{E}_+(t\bar{v}_+)$  is decreasing with respect to  $t$ ,  $t > 0$ . Now, let  $t_0 > 1$  such that  $t_0 \|\bar{v}_+\|_X = 1$ . Then  $t_0 \bar{v}_+ \in S$ , but  $\widehat{E}_+(t_0 \bar{v}_+) < \widehat{E}_+(\bar{v}_+) = m_+ = \inf_S \widehat{E}_+$ , a contradiction.  $\square$

*Proof of Theorem 5.1.* On account of Propositions 5.1, 5.2 and Theorem 1.32, we conclude that  $\bar{u}_+ = \lambda_+(\bar{v}_+) \bar{v}_+$  and  $\bar{u}_- = \lambda_-(\bar{v}_-) \bar{v}_-$  are two nontrivial critical points of  $E$ , thus, they correspond to two nontrivial pairs of solutions of system  $(\bar{S})$ . Note that these pairs of solutions may coincide.

### 5.2.2 Arbitrary superlinear terms: application of the mountain pass theorem

In this subsection we are coming back to the initial system  $(\tilde{S})$  where the continuous function  $f : \mathbb{R} \rightarrow \mathbb{R}$  verifies the assumptions

(H<sub>1</sub>) there exist constants  $\theta > 1 + \frac{1}{p}$  and  $s_0 > 0$  such that

$$0 < \theta F(s) \leq f(s)s \text{ for all } |s| \geq s_0;$$

(H<sub>2</sub>)  $f(s) = o(s^{\frac{1}{p}})$  for  $s \rightarrow 0$ .

Here,  $F(s) = \int_0^s f(t)dt$ .

**Remark 5.1** A typical nonlinearity which fulfills (H<sub>1</sub>) and (H<sub>2</sub>) is  $f(s) = s^q$  with  $pq > 1$ . This particular case was handled in the previous subsection by using the fibering method.

Note that hypothesis (H<sub>1</sub>) is a sort of Ambrosetti-Rabinowitz condition, see [8]. In particular, (H<sub>1</sub>) implies that at infinity,  $|f|$  grows faster than  $s \mapsto |s|^{\theta-1}$ .

The main result of this section reads as follows.

**Theorem 5.2** *Assume that (5.1) holds and the continuous function  $f : \mathbb{R} \rightarrow \mathbb{R}$  verifies hypotheses  $H_1$  and  $H_2$ . Then, system  $(\tilde{S})$  possesses a nontrivial (strong) solution.*

*Proof* First, on account of  $(H_2)$ , to every  $\varepsilon > 0$ , there exists  $\delta > 0$  such that

$$F(s) \leq \varepsilon |s|^{1+\frac{1}{p}} \text{ for all } |s| \leq \delta.$$

Due to (5.2), for each  $u \in X$  complying with  $\|u\|_X = \kappa_0/\delta > 0$ , we have that

$$\begin{aligned} E(u) &= \frac{p}{p+1} \|u\|_X^{\frac{p+1}{p}} - \int_{\Omega} F(u(x)) dx \\ &\geq \frac{p}{p+1} \|u\|_X^{\frac{p+1}{p}} - \varepsilon m(\Omega) \kappa_0^{\frac{p+1}{p}} \|u\|_X^{\frac{p+1}{p}} \\ &= \left( \frac{p}{p+1} - \varepsilon m(\Omega) \kappa_0^{\frac{p+1}{p}} \right) (\kappa_0/\delta)^{\frac{p+1}{p}} = M_0 > 0. \end{aligned}$$

Second, due to  $(H_1)$ , there exists  $c_0 > 0$  such that

$$F(s) \geq c_0 |s|^{\theta} \text{ for all } |s| \geq s_0.$$

Fix  $w \in X \setminus \{0\}$ ,  $w \geq 0$ ; see for instance the function in (5.8) below. For every  $\tau > 0$ , we define the set  $\Omega_{\tau} = \{x \in \Omega; \tau w(x) \geq s_0\}$ . Moreover, let  $\tau_0 > 0$  such that  $m(\Omega_{\tau_0}) > 0$ . Then, for every  $\tau > \tau_0$  we have that

$$\begin{aligned} \int_{\Omega} F(\tau w(x)) dx &= \int_{\Omega_{\tau}} F(\tau w(x)) dx + \int_{\Omega \setminus \Omega_{\tau}} F(\tau w(x)) dx \\ &\geq c_0 \tau^{\theta} \int_{\Omega_{\tau_0}} (w(x))^{\theta} dx - \max_{s \in [0, s_0]} F(s) m(\Omega). \end{aligned}$$

Using this estimation and the fact that  $\theta > \frac{p+1}{p}$ , we obtain that  $E(\tau w) \rightarrow -\infty$  as  $\tau \rightarrow \infty$ .

Finally, we prove that  $E$  verifies the  $(PS)_c$ -condition for every  $c \in \mathbb{R}$ . Let  $\{u_k\} \subset X$  be a sequence such that  $E(u_k) \rightarrow c$  as  $k \rightarrow \infty$  and  $|\langle E'(u_k), v \rangle| \leq \varepsilon_k \|v\|_X$  if  $\varepsilon_k \rightarrow 0$ , for all  $v \in X$ . Then, due to  $(H_1)$ , for large  $k \in \mathbb{N}$ , we have that

$$\begin{aligned} c + \varepsilon_k \|u_k\|_X &\geq |\theta E(u_k) - \langle E'(u_k), u_k \rangle| \\ &\geq \left( \theta \frac{p}{p+1} - 1 \right) \|u_k\|_X^{\frac{p+1}{p}} \\ &\quad - \theta \int_{\Omega} F(u_k(x)) dx + \int_{\Omega} f(u_k(x)) u_k(x) dx \\ &\geq \left( \theta \frac{p}{p+1} - 1 \right) \|u_k\|_X^{\frac{p+1}{p}} - C_0, \end{aligned}$$

for some  $C_0 > 0$  independent of  $k \in \mathbb{N}$ . Consequently,  $\{u_k\}$  is bounded in  $X$ . Therefore, on account of (5.2), up to a subsequence,  $\{u_k\}$  converges strongly in  $C(\Omega)$  to an  $u \in X$ . Now, it is a standard fact to prove that  $\{u_k\}$  converges strongly in  $X$  to  $u$ .

It remains to apply Theorem 1.7 by choosing  $e_0 = 0$ ,  $e_1 = \tau w$  for  $\tau > 0$  large enough, and  $\alpha = M_0$ . Thus, there exists a critical point  $u \in X$  of  $E$  such that  $E(u) \geq \alpha = M_0 > 0$ . In particular,  $u \neq 0$ , which ends the proof.  $\square$

### 5.3 Elliptic systems with oscillatory terms

Let  $x_0 \in \Omega$  and  $R > 0$  be such that  $B(x_0, R) \subset \Omega$ ; here and in the sequel,  $B(x_0, a) = \{x \in \mathbb{R}^N : |x - x_0| < a\}$ ,  $a > 0$ . Let  $0 < r < R$  be fixed. We consider the function  $w : \Omega \rightarrow \mathbb{R}$  defined by

$$w(x) = \frac{\int_{-\infty}^{-|x-x_0|} \alpha(t) dt}{\int_{-R}^{-r} \alpha(t) dt}, \quad (5.8)$$

where  $\alpha : \mathbb{R} \rightarrow \mathbb{R}$  is given by

$$\alpha(t) = \begin{cases} e^{\frac{1}{(t+R)(t+r)}}, & \text{if } t \in ]-R, -r[; \\ 0, & \text{if } t \notin ]-R, -r[. \end{cases}$$

It is clear that  $w \in C_0^\infty(\Omega) \subset X$ ; moreover,  $w \geq 0$ ,  $\|w\|_\infty = 1$  and

$$w(x) = \begin{cases} 1, & \text{if } x \in B(x_0, r); \\ 0, & \text{if } x \in \Omega \setminus B(x_0, R). \end{cases} \quad (5.9)$$

**Lemma 5.2** *Let  $\{a_k\}, \{b_k\} \subset ]0, \infty[$  be two sequences such that  $a_k < b_k$ ,  $\lim_{k \rightarrow \infty} a_k/b_k = 0$ , and  $\text{sgn}(s)f(s) \leq 0$  for every  $|s| \in [a_k, b_k]$ . Let  $s_k = (b_k/\kappa_0)^{\frac{p+1}{p}}$ . Then,*

- a)  $\max_{[-b_k, b_k]} F = \max_{[-a_k, a_k]} F \equiv F(\bar{s}_k)$  with  $\bar{s}_k \in [-a_k, a_k]$ .  
b)  $\|\bar{s}_k w\|_{X^p}^{\frac{p+1}{p}} < s_k$  for  $k \in \mathbb{N}$  large enough.

*Proof* a) It follows from the standard mean value theorem and from the hypotheses that  $\text{sgn}(s)f(s) \leq 0$  for every  $|s| \in [a_k, b_k]$ .

b) Since  $\lim_{k \rightarrow \infty} a_k/b_k = 0$ , we may fix  $k_0 \in \mathbb{N}$  such that  $a_k/b_k < \kappa_0^{-1} \|w\|_X^{-1}$  for  $k > k_0$ . Then, one has  $\|\bar{s}_k w\|_{X^p}^{\frac{p+1}{p}} = |\bar{s}_k|^{\frac{p+1}{p}} \|w\|_{X^p}^{\frac{p+1}{p}} \leq a_k^{\frac{p+1}{p}} \|w\|_{X^p}^{\frac{p+1}{p}} < (b_k/\kappa_0)^{\frac{p+1}{p}} = s_k$ .  $\square$

As we pointed out, the results of this section are based on Ricceri's variational principle, see Theorem 1.20. In our framework concerning problem (P) (thus, system ( $\tilde{S}$ )),  $\Psi, \Phi : X \rightarrow \mathbb{R}$  are defined by

$$\Psi(u) = \|u\|_X^{\frac{p+1}{p}}, \quad \Phi(u) = -\mathcal{F}(u), \quad u \in X.$$

Standard arguments show that  $\Psi$  and  $\Phi$  are sequentially weakly lower semicontinuous. The energy functional becomes  $E = \frac{p}{p+1}\Psi + \Phi$ . Moreover the function from (1.24) takes the form

$$\varphi(s) = \inf_{\|u\|_X^{\frac{p+1}{p}} < s^p} \frac{\sup\{\mathcal{F}(v) : \|v\|_X^{\frac{p+1}{p}} \leq s^p\} - \mathcal{F}(u)}{s - \|u\|_X^{\frac{p+1}{p}}}, \quad s > 0. \quad (5.10)$$

Now, we are in the position to state our first result. Let  $f \in C(\mathbb{R}, \mathbb{R})$  and  $F(s) = \int_0^s f(t)dt$ ,  $s \in \mathbb{R}$ . We assume that:

- ( $H_0^1$ ).  $-\infty < \liminf_{s \rightarrow 0} \frac{F(s)}{|s|^{\frac{p+1}{p}}} \leq \limsup_{s \rightarrow 0} \frac{F(s)}{|s|^{\frac{p+1}{p}}} = +\infty$ ,
- ( $H_0^2$ ). there exist two sequences  $\{a_k\}$  and  $\{b_k\}$  in  $]0, \infty[$  with  $b_{k+1} < a_k < b_k$ ,  $\lim_{k \rightarrow \infty} b_k = 0$  such that

$$\text{sgn}(s)f(s) \leq 0 \quad \text{for every } |s| \in [a_k, b_k], \text{ and}$$

- ( $H_0^3$ ).  $\lim_{k \rightarrow \infty} \frac{a_k}{b_k} = 0$  and  $\lim_{k \rightarrow \infty} \frac{\max_{[-a_k, a_k]} F}{b_k^{\frac{p+1}{p}}} = 0$ .

**Remark 5.2** Hypotheses ( $H_0^1$ ) – ( $H_0^2$ ) imply an oscillatory behaviour of  $f$  near the origin while ( $H_0^3$ ) is a technical assumption which seems to be indispensable in our arguments.

In the sequel, we provide a concrete example when hypotheses ( $H_0^1$ ) – ( $H_0^3$ ) are fulfilled. Let  $a_k = k^{-k^{k+1}}$  and  $b_k = k^{-k^k}$ ,  $k \geq 2$  and  $a_1 = 1$ ,  $b_1 = 2$ . It is clear that  $b_{k+1} < a_k < b_k$ ,  $\lim_{k \rightarrow \infty} a_k/b_k = 0$ , and  $\lim_{k \rightarrow \infty} b_k = 0$ . Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be defined by

$$f(s) = \begin{cases} \varphi_k \left( \frac{s-b_{k+1}}{a_k-b_{k+1}} \right), & s \in [b_{k+1}, a_k], \quad k \geq 1; \\ 0, & s \in ]a_k, b_k[, \quad k \geq 1; \\ 0, & s \in ]-\infty, 0]; \\ g(s), & s \in [2, \infty[, \end{cases}$$

where  $g : [2, \infty[ \rightarrow \mathbb{R}$  is any continuous function with  $g(2) = 0$ , and  $\varphi_k : [0, 1] \rightarrow [0, \infty[$  is a sequence of continuous functions such that  $\varphi_k(0) = \varphi_k(1) = 0$  and there are some positive constants  $c_1$  and  $c_2$  such

that

$$c_1(b_k^{\frac{2p+2}{p}} - b_{k+1}^{\frac{2p+2}{p}})(a_k - b_{k+1})^{-1} \leq \int_0^1 \varphi_k(s) ds \leq c_2(b_k^{\frac{p+2}{p}} - b_{k+1}^{\frac{p+2}{p}})(a_k - b_{k+1})^{-1}.$$

Note that  $F(s) = 0$  for every  $s \in ]-\infty, 0]$  and  $F$  is non-decreasing on  $[0, 2]$ , while  $c_1 b_k^{\frac{2p+2}{p}} \leq F(a_k) = \max_{[-a_k, a_k]} F \leq c_2 b_k^{\frac{p+2}{p}}$ . Due to these inequalities, each hypotheses from above are verified.

**Theorem 5.3** *Assume that (5.1) holds and  $f \in C(\mathbb{R}, \mathbb{R})$  fulfills  $(H_0^1) - (H_0^3)$ . Then, system  $(\tilde{S})$  possesses a sequence  $\{(u_k, v_k)\} \subset X \times X$  of distinct (strong) solutions which satisfy*

$$\lim_{k \rightarrow \infty} \|u_k\|_X = \lim_{k \rightarrow \infty} \|v_k\|_X = \lim_{k \rightarrow \infty} \|u_k\|_\infty = \lim_{k \rightarrow \infty} \|v_k\|_\infty = 0. \quad (5.11)$$

The proof of Theorem 5.3 is based on the following two lemmas; thus, we assume the hypotheses of Theorem 5.3 are fulfilled. Let  $\{a_k\}$  and  $\{b_k\}$  be as in the hypotheses. We recall from (1.25) that  $\delta = \liminf_{s \rightarrow 0^+} \varphi(s)$  where  $\varphi$  comes from (5.10).

**Lemma 5.3**  $\delta = 0$ .

*Proof* By definition,  $\delta \geq 0$ . Suppose that  $\delta > 0$ . By the first inequality of  $(H_0^1)$ , there exist two positive numbers  $\ell_0$  and  $\varrho_0$  such that

$$F(s) > -\ell_0 |s|^{\frac{p+1}{p}} \quad \text{for every } s \in ]-\varrho_0, \varrho_0[. \quad (5.12)$$

Furthermore, let  $s_k, \bar{s}_k$  be as in Lemma 5.2 and let  $\bar{w}_k = \bar{s}_k w \in E$ , where  $w$  is defined in (5.8). By  $(H_0^3)$  and condition  $\lim_{k \rightarrow \infty} \frac{\bar{s}_k}{b_k} = 0$  ( $|\bar{s}_k| \leq a_k$ ), there exists  $k_0 \in \mathbb{N}$  such that for  $k > k_0$  we have

$$m(\Omega) \frac{F(\bar{s}_k)}{b_k^{\frac{p+1}{p}}} + \left( \frac{\delta}{2} \|w\|_X^{\frac{p+1}{p}} + m(\Omega) \ell_0 \right) \left( \frac{|\bar{s}_k|}{b_k} \right)^{\frac{p+1}{p}} < \frac{\delta}{2\kappa_0^{\frac{p+1}{p}}}. \quad (5.13)$$

Let  $v \in E$  be arbitrarily fixed with  $\|v\|_X^{\frac{p+1}{p}} \leq s_k$ . Thus, due to the embedding from (5.2), we have  $\|v\|_\infty \leq b_k$ . Due to Lemma 5.2 a), we obtain

$$F(v(x)) \leq \max_{[-b_k, b_k]} F = F(\bar{s}_k) \quad \text{for every } x \in \Omega.$$

Since  $0 \leq |\bar{w}_k(x)| \leq |\bar{s}_k| < \varrho_0$  for large  $k \in \mathbb{N}$  and for all  $x \in \Omega$ , taking

into account (5.12) and (5.13), it follows that

$$\begin{aligned} \sup_{\|v\|_X^{p+1} \leq s_k^p} \mathcal{F}(v) - \mathcal{F}(\bar{w}_k) &= \sup_{\|v\|_X^{\frac{p+1}{p}} \leq s_k} \int_{\Omega} F(v) dx - \int_{\Omega} F(\bar{w}_k) dx \\ &\leq m(\Omega)F(\bar{s}_k) + m(\Omega)\ell_0|\bar{s}_k|^{\frac{p+1}{p}} \\ &< \frac{\delta}{2}(s_k - \|\bar{w}_k\|_X^{\frac{p+1}{p}}). \end{aligned}$$

Since  $\|\bar{w}_k\|_X^{\frac{p+1}{p}} < s_k$  (cf. Lemma 5.2 b)), and  $s_k \rightarrow 0$  as  $k \rightarrow \infty$ , we obtain

$$\delta \leq \liminf_{k \rightarrow \infty} \varphi(s_k) \leq \liminf_{k \rightarrow \infty} \frac{\sup_{\|v\|_X^{p+1} \leq s_k^p} \mathcal{F}(v) - \mathcal{F}(\bar{w}_k)}{s_k - \|\bar{w}_k\|_X^{\frac{p+1}{p}}} \leq \frac{\delta}{2},$$

contradiction. This proves our claim.  $\square$

**Lemma 5.4** *0 is not a local minimum of  $E = \frac{p}{p+1}\Psi + \Phi$ .*

*Proof* Let  $\ell_0 > 0$  and  $\varrho_0 > 0$  from the proof of Lemma 5.3, and  $x_0 \in \Omega$  and  $r, R > 0$  from the definition of the function  $w$ , see (5.8). Let  $\mathcal{L}_0 > 0$  be such that

$$r^N \omega_N \mathcal{L}_0 - \frac{p}{p+1} \|w\|_X^{\frac{p+1}{p}} - (R^N - r^N) \omega_N \ell_0 > 0, \quad (5.14)$$

where  $\omega_N$  is the volume of the  $N$ -dimensional unit ball. By the right hand side of  $(H_0^1)$  we deduce the existence of a sequence  $\{s_k^0\} \subset ]-\varrho_0, \varrho_0[$  converging to zero such that

$$F(s_k^0) > \mathcal{L}_0 |s_k^0|^{\frac{p+1}{p}}. \quad (5.15)$$

Let  $w_k^0 = s_k^0 w \in E$ . Due to (5.9), (5.12), (5.14) and (5.15), we have

$$\begin{aligned} E(w_k^0) &= \frac{p}{p+1} \|w_k^0\|_X^{\frac{p+1}{p}} - \int_{\Omega} F(w_k^0) \\ &= \frac{p}{p+1} \|w\|_X^{\frac{p+1}{p}} |s_k^0|^{\frac{p+1}{p}} - \int_{B(x_0, r)} F(w_k^0) - \int_{B(x_0, R) \setminus B(x_0, r)} F(w_k^0) \\ &\leq \frac{p}{p+1} \|w\|_X^{\frac{p+1}{p}} |s_k^0|^{\frac{p+1}{p}} - F(s_k^0) m(B(x_0, r)) + \ell_0 (m(B(0, R)) \\ &\quad - m(B(0, r))) |s_k^0|^{\frac{p+1}{p}} \\ &\leq |s_k^0|^{\frac{p+1}{p}} \left( \frac{p}{p+1} \|w\|_X^{\frac{p+1}{p}} - r^N \omega_N \mathcal{L}_0 + (R^N - r^N) \omega_N \ell_0 \right) \\ &< 0 = E(0). \end{aligned}$$

Since  $\|w_k^0\|_X \rightarrow 0$  as  $k \rightarrow \infty$ , 0 is not a local minimum of  $E$ .  $\square$

*Proof of Theorem 5.3.* Applying Theorem 1.20 with  $\lambda = \frac{p}{p+1}$  (see Lemma 5.3), we can exclude condition (A1) (see Lemma 5.4). Therefore there exists a sequence  $\{u_k\} \subset X$  of pairwise distinct critical points of  $E = \frac{p}{p+1}\Psi + \Phi$  such that

$$\lim_{k \rightarrow \infty} \|u_k\|_X = 0. \quad (5.16)$$

Thus,  $\{(u_k, v_k)\} = \{(u_k, (-\Delta u_k)^{\frac{1}{p}})\} \subset X \times X$  is a sequence of distinct pairs of solutions to the system  $(\tilde{S})$ .

It remains to prove (5.11). First, due to (5.2) and (5.16), we have that  $\lim_{k \rightarrow \infty} \|u_k\|_\infty = 0$ . For every  $k \in \mathbb{N}$  let  $m_k \in [-\|u_k\|_\infty, \|u_k\|_\infty] =: J_k$  such that  $|f(m_k)| = \max_{s \in J_k} |f(s)|$ . Note that  $\text{diam} J_k \rightarrow 0$  as  $k \rightarrow \infty$ ; thus  $\lim_{k \rightarrow \infty} m_k = 0$  which implies that  $\lim_{k \rightarrow \infty} f(m_k) = 0$ . On the other hand, from the second equation of system  $(\tilde{S})$  we have that

$$\|v_k\|_X^{\frac{p+1}{p}} = \int_{\Omega} |\Delta v_k|^{\frac{p+1}{p}} dx = \int_{\Omega} |f(u_k)|^{\frac{p+1}{p}} dx \leq |f(m_k)|^{\frac{p+1}{p}} m(\Omega),$$

which implies that  $\lim_{k \rightarrow \infty} \|v_k\|_X = 0$ . Using again (5.2) we have that  $\lim_{k \rightarrow \infty} \|v_k\|_\infty = 0$ .

In the sequel, we state a perfect counterpart of Theorem 5.3 when the nonlinearity  $f$  has an oscillation at infinity. We assume that:

$$(H_\infty^1). \quad -\infty < \liminf_{|s| \rightarrow \infty} \frac{F(s)}{|s|^{\frac{p+1}{p}}} \leq \limsup_{|s| \rightarrow \infty} \frac{F(s)}{|s|^{\frac{p+1}{p}}} = +\infty,$$

$$(H_\infty^2). \quad \text{there exist two sequences } \{a_k\} \text{ and } \{b_k\} \text{ in } ]0, \infty[ \text{ with } a_k < b_k < a_{k+1} \\ \text{and } \lim_{k \rightarrow \infty} b_k = \infty \text{ such that}$$

$$\text{sgn}(s)f(s) \leq 0 \quad \text{for every } |s| \in [a_k, b_k], \text{ and}$$

$$(H_\infty^3). \quad \lim_{k \rightarrow \infty} \frac{a_k}{b_k} = 0 \text{ and } \lim_{k \rightarrow \infty} \frac{\max_{[-a_k, a_k]} F}{b_k^{\frac{p+1}{p}}} = 0.$$

**Remark 5.3** Assumptions  $(H_\infty^1) - (H_\infty^2)$  imply an oscillatory behaviour of  $f$  at infinity. A concrete example is described in the sequel when hypotheses  $(H_\infty^1) - (H_\infty^3)$  are fulfilled. Let  $a_k = k^{k^k}$  and  $b_k = k^{k^{k+1}}$  ( $k \geq 2$ ) and  $a_1 = 5$ ,  $b_1 = 10$ . Clearly, one has  $a_k < b_k < a_{k+1}$ ,  $\lim_{k \rightarrow \infty} a_k/b_k = 0$ , and  $\lim_{k \rightarrow \infty} b_k = \infty$ . Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be defined

by

$$f(s) = \begin{cases} \varphi_k \left( \frac{s-b_k}{a_{k+1}-b_k} \right), & s \in [b_k, a_{k+1}], \quad k \geq 1; \\ 0, & s \in ]a_k, b_k[, \quad k \geq 1; \\ g(s), & s \in [-5, 5]; \\ 0, & s \in ]-\infty, -5[, \end{cases}$$

where  $g : [-5, 5] \rightarrow \mathbb{R}$  is any continuous function with  $g(\pm 5) = 0$ , and  $\varphi_k : [0, 1] \rightarrow [0, \infty[$  is a sequence of continuous functions such that  $\varphi_k(0) = \varphi_k(1) = 0$  and there are some constants  $c_1, c_2 > 0$  such that

$$c_1(b_{k+1}^{\frac{3p+1}{3p}} - b_k^{\frac{3p+1}{3p}})(a_{k+1}-b_k)^{-1} \leq \int_0^1 \varphi_k(s)ds \leq c_2(b_{k+1}^{\frac{2p+1}{2p}} - b_k^{\frac{2p+1}{2p}})(a_{k+1}-b_k)^{-1}.$$

Note that  $F(s) = 0$  for every  $s \in ]-\infty, -5]$  and  $F$  is non-decreasing on  $[5, \infty[$ . Moreover, for  $k \in \mathbb{N}$  large enough we have

$$c_1(b_k^{\frac{3p+1}{3p}} - 10^{\frac{3p+1}{3p}}) + \int_0^5 g(s)ds \leq F(a_k) = \max_{[-a_k, a_k]} F \leq c_2(b_k^{\frac{2p+1}{2p}} - 10^{\frac{2p+1}{2p}}) + \int_0^5 g(s)ds.$$

Now, an easy computation shows the hypotheses from above are verified.

**Theorem 5.4** *Assume that (5.1) holds and  $f \in C(\mathbb{R}, \mathbb{R})$  fulfills  $(H_\infty^1)$ – $(H_\infty^3)$ . Then, system  $(\tilde{S})$  possesses a sequence  $\{(u_k, v_k)\} \subset X \times X$  of distinct (strong) solutions which satisfy*

$$\lim_{k \rightarrow \infty} \|u_k\|_X = \lim_{k \rightarrow \infty} \|v_k\|_X = \lim_{k \rightarrow \infty} \|u_k\|_\infty = \lim_{k \rightarrow \infty} \|v_k\|_\infty = \infty. \quad (5.17)$$

The proof of Theorem 5.4 is similar to that of Theorem 5.3. Let  $\{a_k\}$  and  $\{b_k\}$  be from Theorem 5.4 and  $\gamma = \liminf_{s \rightarrow +\infty} \varphi(s)$  from (1.25) where  $\varphi$  comes from (5.10).

**Lemma 5.5**  $\gamma = 0$ .

*Proof* It is clear that  $\gamma \geq 0$ . Suppose that  $\gamma > 0$ . Due to the left-hand side of  $(H_\infty^1)$ , one can find two positive numbers  $\ell_\infty$  and  $\varrho_\infty$  such that

$$F(s) > -\ell_\infty |s|^{\frac{p+1}{p}} \quad \text{for every } |s| > \varrho_\infty. \quad (5.18)$$

Let  $s_k, \bar{s}_k$  be as in Lemma 5.2. By the fact that  $\lim_{k \rightarrow \infty} b_k = \infty$ , hypothesis  $(H_\infty^3)$  and condition  $\lim_{k \rightarrow \infty} \frac{\bar{s}_k}{b_k} = 0$  ( $-a_k \leq \bar{s}_k \leq a_k$ ), there exists  $k_0 \in \mathbb{N}$  such that for every  $k > k_0$  we have

$$m(\Omega) \frac{\max_{[-\varrho_\infty, \varrho_\infty]} |F| + F(\bar{s}_k)}{b_k^{\frac{p+1}{p}}} + \left( \frac{\gamma}{2} \|w\|_{X^p}^{\frac{p+1}{p}} + m(\Omega)\ell_\infty \right) \left( \frac{|\bar{s}_k|}{b_k} \right)^{\frac{p+1}{p}} < \frac{\gamma}{2\kappa_0^{\frac{p+1}{p}}}. \quad (5.19)$$

Let  $\bar{w}_k = \bar{s}_k w \in X$ , where  $w$  is defined in (5.8). A similar estimation as in Lemma 5.3 gives throughout relations (5.18) and (5.19) that

$$\begin{aligned}
\sup_{\|v\|_X^{p+1} \leq s_k^p} \mathcal{F}(v) - \mathcal{F}(\bar{w}_k) &= \sup_{\|v\|_X^{\frac{p+1}{p}} \leq s_k} \int_{\Omega} F(v) dx - \int_{\Omega} F(\bar{w}_k) dx \\
&= \sup_{\|v\|_X^{\frac{p+1}{p}} \leq s_k} \int_{\Omega} F(v) dx - \int_{\{|w_k(x)| > \varrho_{\infty}\}} F(\bar{w}_k) dx \\
&\quad - \int_{\{|w_k(x)| \leq \varrho_{\infty}\}} F(\bar{w}_k) dx \\
&\leq m(\Omega) F(\bar{s}_k) + m(\Omega) \varrho_{\infty} |\bar{s}_k|^{\frac{p+1}{p}} \\
&\quad + m(\Omega) \max_{[-\varrho_{\infty}, \varrho_{\infty}]} |F| \\
&< \frac{\gamma}{2} (s_k - \|\bar{w}_k\|_X^{\frac{p+1}{p}}).
\end{aligned}$$

Since  $s_k \rightarrow +\infty$ ,

$$\gamma \leq \liminf_{k \rightarrow \infty} \varphi(s_k) \leq \liminf_{k \rightarrow \infty} \frac{\sup_{\|v\|_X^{p+1} \leq s_k^p} \mathcal{F}(v) - \mathcal{F}(\bar{w}_k)}{s_k - \|\bar{w}_k\|_X^{\frac{p+1}{p}}} \leq \frac{\gamma}{2},$$

which contradicts  $\gamma > 0$ .  $\square$

**Lemma 5.6**  $E = \frac{p}{p+1} \Psi + \Phi$  is not bounded from below on  $X$ .

*Proof* Let  $\ell_{\infty}$  and  $\varrho_{\infty}$  from the proof of Lemma 5.5, and let  $\mathcal{L}_{\infty} > 0$  be such that

$$r^N \omega_N \mathcal{L}_{\infty} - \frac{p}{p+1} \|w\|_X^{\frac{p+1}{p}} - (R^N - r^n) \omega_N \ell_{\infty} > 0, \quad (5.20)$$

where  $r$  and  $R$  are from the definition of the function  $w$ , see (5.8). By the second part of  $(H_{\infty}^1)$  we deduce the existence of a sequence  $\{s_k^{\infty}\} \subset \mathbb{R}$  with  $\lim_{k \rightarrow \infty} |s_k^{\infty}| = \infty$  and

$$F(s_k^{\infty}) > \mathcal{L}_{\infty} |s_k^{\infty}|^{\frac{p+1}{p}}. \quad (5.21)$$

Let  $w_k^{\infty} = s_k^{\infty} w \in X$ . We clearly have that

$$E(w_k^{\infty}) = \frac{p}{p+1} \|w\|_X^{\frac{p+1}{p}} |s_k^{\infty}|^{\frac{p+1}{p}} - F(s_k^{\infty}) \omega_N r^N - \int_{B(x_0, R) \setminus B(x_0, r)} F(w_k^{\infty}).$$

For abbreviation, we choose the set  $D = B(x_0, R) \setminus B(x_0, r)$ . Then, on

account of (5.18) we have

$$\begin{aligned} \int_D F(w_k^\infty) &= \int_{D \cap \{|w_k^\infty(x)| > \ell_\infty\}} F(w_k^\infty) + \int_{D \cap \{|w_k^\infty(x)| \leq \ell_\infty\}} F(w_k^\infty) \\ &\geq -\ell_\infty \int_{D \cap \{|w_k^\infty(x)| > \ell_\infty\}} |w_k^\infty|^{\frac{p+1}{p}} - (R^n - r^N) \omega_N \max_{[-\ell_\infty, \ell_\infty]} |F| \\ &\geq -(R^n - r^N) \omega_N \left( \ell_\infty |s_k^\infty|^{\frac{p+1}{p}} + \max_{[-\ell_\infty, \ell_\infty]} |F| \right). \end{aligned}$$

Consequently, due to (5.21) and the above estimation, we have

$$\begin{aligned} E(w_k^\infty) &\leq |s_k^\infty|^{\frac{p+1}{p}} \left( \frac{p}{p+1} \|w\|_X^{\frac{p+1}{p}} - r^N \omega_N \mathcal{L}_\infty + (R^N - r^n) \omega_N \ell_\infty \right) \\ &\quad + (R^N - r^n) \omega_N \max_{[-\ell_\infty, \ell_\infty]} |F|. \end{aligned}$$

Since  $\lim_{k \rightarrow \infty} |s_k^\infty| = \infty$ , due to (5.20), we have  $\lim_{k \rightarrow \infty} E(w_k^\infty) = -\infty$ ; consequently,  $\inf_X E = -\infty$ .  $\square$

*Proof of Theorem 5.4.* In Theorem 1.20 we may choose  $\lambda = \frac{p}{p+1}$  (see Lemma 5.5). On account of Lemma 5.6 the alternative (B1) can be excluded. Therefore, there exists a sequence  $\{u_k\} \subset X$  of distinct critical points of  $E = \frac{p}{p+1} \Psi + \Phi$  such that

$$\lim_{k \rightarrow \infty} \|u_k\|_X = \infty. \quad (5.22)$$

Thus,  $\{(u_k, v_k)\} = \{(u_k, (-\Delta u_k)^{\frac{1}{p}})\} \subset X \times X$  is a sequence of distinct pairs of solutions to the system  $(\tilde{S})$ .

We now prove the rest of (5.17). Assume that for every  $k \in \mathbb{N}$  we have  $\|v_k\|_\infty \leq M$  for some  $M > 0$ . In particular, from the first equation of system  $(\tilde{S})$  we obtain that

$$\|u_k\|_X^{\frac{p+1}{p}} = \int_\Omega |\Delta u_k|^{\frac{p+1}{p}} dx = \int_\Omega |v_k|^{p+1} dx \leq M^{p+1} m(\Omega),$$

which contradicts relation (5.22). Consequently,  $\lim_{k \rightarrow \infty} \|v_k\|_\infty = \infty$ . But, this fact and (5.2) give at once that  $\lim_{k \rightarrow \infty} \|v_k\|_X = \infty$  as well. Assume finally that for every  $k \in \mathbb{N}$  we have  $\|u_k\|_\infty \leq M'$  for some  $M' > 0$ . The second equation of system  $(\tilde{S})$  shows that

$$\|v_k\|_X^{\frac{p+1}{p}} = \int_\Omega |\Delta v_k|^{\frac{p+1}{p}} dx = \int_\Omega |f(u_k)|^{\frac{p+1}{p}} dx \leq m(\Omega) \max_{s \in [-M', M']} |f(s)|^{\frac{p+1}{p}},$$

which contradicts the fact that  $\lim_{k \rightarrow \infty} \|v_k\|_X = \infty$ . The proof is complete.

### 5.4 Comments and perspectives

1. Both Theorems 5.1 and 5.2 provide the existence of at least one nonzero solution to the system  $(\tilde{S})$  for  $f(s) = s^q$  with  $pq > 1$  by exploiting two different methods, that is, the fibering method and the mountain pass argument, respectively. Can we distinguish these solutions?

2. We assume (5.1) holds and consider the perturbed system

$$\begin{cases} -\Delta u = v^p & \text{in } \Omega; \\ -\Delta v = f(u) + \varepsilon g(u) & \text{in } \Omega; \\ u = v = 0 & \text{on } \partial\Omega, \end{cases} \quad (\tilde{S}_\varepsilon)$$

where  $g : \mathbb{R} \rightarrow \mathbb{R}$  is any continuous function with  $g(0) = 0$ . We predict that for every  $n \in \mathbb{N}$  there exists  $\varepsilon_n > 0$  such that for every  $\varepsilon \in [-\varepsilon_n, \varepsilon_n]$ , system  $(\tilde{S}_\varepsilon)$  has at least  $n$  distinct pairs of solutions whenever  $f : \mathbb{R} \rightarrow \mathbb{R}$  verifies the set of assumptions from Theorems 5.3 or 5.4. This statement is not unexpected taking into account the recent papers of Anello and Cordaro [13] and Kristály [170] where a prescribed number of solutions were guaranteed for certain elliptic problems of *scalar* type whenever the parameter in the front of the perturbation is small enough. In both papers [13] and [170], the uniform Lipschitz truncation function  $h_a : \mathbb{R} \rightarrow \mathbb{R}$  ( $a > 0$ ),  $h_a(s) = \min(a, \max(s, 0))$  plays a key role, fulfilling as well the so-called *Markovian property* concerning the superposition operators: for any  $u \in W_0^{1,r}(\Omega)$  one also has  $h_a \circ u \in W_0^{1,r}(\Omega)$ , where  $r > 1$ . Note however that a similar property is not available any longer replacing the space  $W_0^{1,r}(\Omega)$  by a higher order Sobolev space  $W^{2,r}(\Omega)$ ; in particular, the Markovian property is not valid for  $X = W^{2, \frac{p+1}{p}}(\Omega) \cap W_0^{1, \frac{p+1}{p}}(\Omega)$ .

3. Based on (5.1), the embedding  $X \subset\subset C(\Omega)$  is essential in our investigations, see the proof of Lemmas 5.3 and 5.5. Is it possible to obtain similar conclusions as in Theorems 5.3 and 5.4 by omitting (5.1) and considering the whole region below the critical hyperbola?

# 6

## Scalar Field Systems

The mathematics are distinguished by a particular privilege, that is, in the course of ages, they may always advance and can never recede.

---

Edward Gibbon (1737–1794),  
*Decline and Fall of the Roman Empire*

In this chapter we study the existence of multiple solutions for scalar fields systems: a double eigenvalue problem for a scalar field systems and a scalar field system with oscillatory nonlinear term. The nonlinear boundary conditions studied here recover the standard Dirichlet, Neumann, or periodic boundary conditions.

### 6.1 Introduction

Let  $h_p : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be the homeomorphism defined by  $h_p(x) = |x|^{p-2}x$  for all  $x \in \mathbb{R}^n$ , where  $p > 1$  is fixed. Let  $|\cdot|$  denote the Euclidean norm in  $\mathbb{R}^n$ . For  $T > 0$ , let  $F : [0, T] \times \mathbb{R}^n \rightarrow \mathbb{R}$  be a Carathéodory mapping satisfying:

( $F_1$ ) for a.e.  $t \in [0, T]$ , the function  $F(t, \cdot)$  is continuously differentiable;  
( $F_2$ ) the mapping  $F(\cdot, x) : [0, T] \rightarrow \mathbb{R}$  is measurable for each  $x \in \mathbb{R}^n$ ,  $F(\cdot, 0) \in L^1(0, T)$ , and for each  $M > 0$  there exists  $\alpha_M \in L^1(0, T)$  such that for all  $x \in \mathbb{R}^n$  with  $|x| < M$ ,

$$|\nabla F(t, x)| \leq \alpha_M(t) \quad \text{for a.e. } t \in [0, T];$$

Let  $j : \mathbb{R}^n \times \mathbb{R}^n \rightarrow ]-\infty, +\infty]$  be a function having the following properties:

- ( $J_1$ )  $D(j) = \{(x, y) \in \mathbb{R}^n \times \mathbb{R}^n : j(x, y) < +\infty\} \neq \emptyset$  is a closed convex cone with  $D(j) \neq \{(0, 0)\}$ ;  
 ( $J_2$ )  $j$  is a convex and lower semi-continuous (shortly, l.s.c.) function.

The first problem studied in this chapter is the following.

Let  $\gamma > 0$  be arbitrary. For  $\lambda, \mu > 0$  we consider the following double eigenvalue problem involving the  $p$ -Laplace operator:

$$(P_{\lambda, \mu}) \quad \begin{cases} -[h_p(u')] + \gamma h_p(u) = \lambda \nabla F(t, u) \text{ a.e. } t \in [0, T], \\ (h_p(u')(0), -h_p(u')(T)) \in \mu \partial j(u(0), u(T)), \end{cases}$$

where  $u : [0, T] \rightarrow \mathbb{R}^n$  is of class  $C^1$  and  $h_p(u')$  is absolutely continuous. Note, that  $\nabla F(t, \eta)$  denotes the gradient of  $F(t, \cdot)$  at  $\eta \in \mathbb{R}^n$ , while  $\partial j$  denotes the subdifferential of  $j$  in the sense of convex analysis.

Next, we assume that:

- (C1)  $F : \mathbb{R}^n \rightarrow \mathbb{R}$  is of class  $C^1$ ,  
 (C2)  $S \subset \mathbb{R}^n \times \mathbb{R}^n$  is a closed convex set with  $\{(x, x) : x \in \mathbb{R}^n\} \subset S$ ,  
 (C3)  $\gamma_1, \dots, \gamma_n \in L^\infty(]0, T[, \mathbb{R})$  are so that  $\text{essinf } \gamma_i > 0$ , for  $i = \overline{1, n}$  and put  $\gamma = (\gamma_1, \dots, \gamma_n)$ ,  
 (C4)  $\alpha \in L^1(]0, T[, \mathbb{R})$  is so that  $\alpha(t) \geq 0$  a.e. in  $]0, T[$ .

The second problem studied in this chapter is the following:

$$(S) \quad \begin{cases} -[h_p(u')] + \gamma \circ h_p(u) = \alpha(t) \nabla F(u) \\ (h_p(u')(0), -h_p(u')(T)) \in N_S(u(0), u(T)), \end{cases}$$

where  $N_S(x, y)$  denotes the normal cone of  $S$  at  $(x, y) \in S$ .

## 6.2 Multiple solutions of a double eigenvalue problem

In this section we prove a multiplicity result for the problem  $(P_{\lambda, \mu})$ . For this purpose we suppose that the following conditions are fulfilled:

- ( $F_3$ )  $\lim_{|x| \rightarrow \infty} \frac{F(t, x) - F(t, 0)}{|x|^p} \leq 0$  uniformly for a.e.  $t \in [0, T]$ .

$$(F_4) \lim_{|x| \rightarrow 0} \frac{F(t, x) - F(t, 0)}{|x|^p} \leq 0 \text{ uniformly for a.e. } t \in [0, T];$$

$$(F_5) \text{ there exists } s_0 \in \mathbb{R}^N \text{ such that } \int_0^T (F(t, s_0) - F(t, 0)) dt > 0.$$

$$(J_3) j(0, 0) = 0, j(x, y) \geq 0 \text{ for all } (x, y) \in \mathbb{R}^n \times \mathbb{R}^n.$$

We consider the Sobolev space  $W^{1,p} = W^{1,p}(0, T; \mathbb{R}^n)$  equipped with the norm

$$\|u\|_\eta = \left( \|u'\|_{L^p}^p + \eta \|u\|_{L^p}^p \right)^{1/p},$$

where  $\eta > 0$ , and denote by  $\|\cdot\|_{L^p}$  the norm of  $L^p = L^p(0, T; \mathbb{R}^n)$ , that is,

$$\|u\|_{L^p} = \left( \int_0^T |u(t)|^p dt \right)^{1/p}.$$

We consider the space of continuous functions  $C = C([0, T]; \mathbb{R}^n)$  endowed with the norm

$$\|u\|_C = \max\{|u(t)| : t \in [0, T]\}.$$

For  $\gamma > 0$ , we consider  $\varphi_\gamma : W^{1,p} \rightarrow \mathbb{R}$  defined by

$$\varphi_\gamma(u) := \frac{1}{p} \left( \|u'\|_{L^p}^p + \gamma \|u\|_{L^p}^p \right) \text{ for all } u \in W^{1,p}.$$

Note that  $\varphi_\gamma$  is convex,  $\varphi_\gamma \in C^1(W^{1,p}; \mathbb{R})$  and for all  $u, v \in W^{1,p}$ ,

$$\langle \varphi'_\gamma(u), v \rangle = \int_0^T (h_p(u'), v') dt + \gamma \int_0^T (h_p(u), v) dt, \quad (6.1)$$

where  $(\cdot, \cdot)$  denotes the usual inner product in  $\mathbb{R}^n$ .

We define the function  $J : W^{1,p} \rightarrow ]-\infty, +\infty]$  by

$$J(u) = j(u(0), u(T)) \text{ for all } u \in W^{1,p}.$$

Then  $J$  is a proper, convex and l.s.c. function. We also observe that

$$D(J) = \{u \in W^{1,p} : (u(0), u(T)) \in D(j)\}.$$

We consider the functional  $\hat{\mathcal{F}} : C \rightarrow \mathbb{R}$  defined by

$$\hat{\mathcal{F}}(v) = - \int_0^T F(t, v) dt + \int_0^T F(t, 0) dt \text{ for all } v \in C$$

and  $\mathcal{F} : W^{1,p} \rightarrow \mathbb{R}$  defined by  $\mathcal{F} = \hat{\mathcal{F}}|_{W^{1,p}}$ . The functional  $\mathcal{F}$  is sequentially weakly continuous, since the embedding  $W^{1,p} \hookrightarrow C$  is compact. We also have for all  $u, v \in W^{1,p}$ ,

$$\langle \mathcal{F}'(u), v \rangle = - \int_0^T \langle \nabla F(t, u), v \rangle dt. \quad (6.2)$$

Note that for  $1 \leq r < p$  and  $p < q < p^*$  the embeddings  $L^p \hookrightarrow L^r$ ,  $W^{1,p} \hookrightarrow L^q$ ,  $W^{1,p} \hookrightarrow C$  are continuous, hence there exist constants  $C_{r,p}, \hat{C}_{q,p}, \hat{c} > 0$  such that

$$\|u\|_{L^r} \leq C_{r,p} \|u\|_{L^p}, \quad \|u\|_{L^q} \leq \hat{C}_{q,p} \|u\|_{W^{1,p}}, \quad \|u\|_C \leq \hat{c} \|u\|_{W^{1,p}}$$

for all  $u \in W^{1,p}$ .

The energy functional  $I : [0, \infty) \times [0, \infty) \times W^{1,p} \rightarrow ]-\infty, \infty]$  associated to the problem  $(P_{\lambda, \mu})$  is defined by

$$I(\lambda, \mu, u) = \varphi_\gamma(u) + \lambda \mathcal{F}(u) + \mu J(u).$$

The functional  $I$  is of Szulkin type and  $J'$  denote the derivative in the sense of Definition A.4.

The following result is due to Jebelean and Moroşanu [150], Proposition 3.2.

**Proposition 6.1** *Assume that  $F : [0, T] \times \mathbb{R}^n \rightarrow \mathbb{R}$  satisfies  $(F_1)$  and  $(F_2)$  and  $j : \mathbb{R}^n \times \mathbb{R}^n \rightarrow ]-\infty, +\infty]$  satisfies  $(J_1)$  and  $(J_2)$ . If  $u \in W^{1,p}$  is a critical point of  $I(\lambda, \mu, \cdot)$ , then  $u$  is a solution of  $(P_{\lambda, \mu})$ .*

*Proof* Assume that  $u \in W^{1,p}$  is a critical point of  $I(\lambda, \mu, \cdot)$ . Then, for all  $v \in W^{1,p}$ ,

$$\lambda \langle \mathcal{F}'(u), v - u \rangle + \varphi_\gamma(v) - \varphi_\gamma(u) + \mu (J(v) - J(u)) \geq 0.$$

Let  $w \in W^{1,p}$  and set  $v = u + sw$ , where  $s > 0$ . Dividing by  $s$  and then letting  $s \rightarrow 0^+$ , we obtain for all  $w \in W^{1,p}$ ,

$$\lambda \langle \mathcal{F}'(u), w \rangle + \langle \varphi'_\gamma(u), w \rangle + \mu J'(u; w) \geq 0,$$

where  $J'(u; w)$  means the derivative in the sense of Definition A.4.

By the definition of  $J$  we deduce that for every  $w \in W^{1,p}$ ,

$$\lambda \langle \mathcal{F}'(u), w \rangle + \langle \varphi'_\gamma(u), w \rangle + \mu j'((u(0), u(T)); (w(0), w(T))) \geq 0. \quad (6.3)$$

Since  $C_0^\infty(0, T; \mathbb{R}^n) \subset W^{1,p}$ , it follows that for all  $w \in C_0^\infty(0, T; \mathbb{R}^n)$ ,

$$\lambda \langle \mathcal{F}'(u), w \rangle + \langle \varphi'_\gamma(u), w \rangle = 0.$$

Combining relations (6.1) and (6.2) we have for all  $w \in C_0^\infty(0, T; \mathbb{R}^n)$ ,

$$\int_0^T (h_p(u'), w') dt = \int_0^T (-\gamma h_p(u) + \nabla F(t, u), w) dt. \quad (6.4)$$

For any  $u \in W^{1,p}$ , we obtain

$$h_p(u), h_p(u') \in L^{p'}(0, T; \mathbb{R}^n) \text{ with } \frac{1}{p} + \frac{1}{p'} = 1. \quad (6.5)$$

But  $\nabla F(\cdot, u) \in L^1(0, T)$  (by  $(F_2)$ ), hence relations (6.4) and (6.5) yield  $h_p(u') \in W^{1,1}$  and

$$-\left(h_p(u')\right)' = -\gamma h_p(u) + \nabla F(t, u) \text{ for a.e. } t \in [0, T]. \quad (6.6)$$

Since  $h_p$  is a homeomorphism and  $h_p(u') \in W^{1,1}$ , it follows that  $u$  is of class  $C^1$ .

In order to show that  $u$  satisfies the boundary condition of problem  $(P_{\lambda, \mu})$  we note that by (6.3) and (6.6) we have for all  $w \in W^{1,p}$ ,

$$\mu j'((u(0), u(T)); (w(0), w(T))) \geq (h_p(u')(0), w(0)) - (h_p(u')(T), w(T)).$$

Therefore

$$\mu j'((u(0), u(T)); (x, y)) \geq (h_p(u')(0), w(0)) - (h_p(u')(T), w(T)).$$

From Lemma A.2 it follows that

$$\left(h_p(u')(0), -h_p(u')(T)\right) \in \mu \partial j(u(0), u(T)).$$

We conclude that  $u$  is a solution of problem  $(P_{\lambda, \mu})$ .  $\square$

We introduce the constant  $\eta_1 = \eta_1(p, \gamma) > 0$  by setting

$$\eta_1 = \inf \left\{ \frac{\|u'\|_{L^p}^p + \gamma \|u\|_{L^p}^p}{\|u\|_{L^p}^p} : u \in W^{1,p} \setminus \{0\}, u \in D(J) \right\}.$$

Since  $\eta_1 > 0$ , we obtain

$$2^{-1/p} \|u\|_{\eta_1} \leq (\|u'\|_{L^p}^p + \gamma \|u\|_{L^p}^p)^{1/p} \leq \|u\|_{\eta_1} \text{ for all } u \in D(J). \quad (6.7)$$

**Lemma 6.1** [150, Lemma 4.3] *Assume that  $\{u_n\} \subset D(J)$  such that  $u_n \rightharpoonup u$  weakly in  $W^{1,p}$  and*

$$\liminf_{n \rightarrow \infty} \left( \langle \varphi'_\gamma(u_n), u - u_n \rangle + \mu J'(u_n; u - u_n) \right) \geq 0. \quad (6.8)$$

*Then  $\{u_n\}$  has a subsequence which converges strongly in  $W^{1,p}$ .*

*Proof* Note that  $u \in D(J)$ , since the closed convex set  $D(J)$  is weakly closed in  $(W^{1,p}, \|\cdot\|_{\eta_1})$ . Hence,  $J'(u_n; u - u_n) < \infty$  for all  $n \in \mathbb{N}$ . Combining relation (6.8) and the lower semi-continuity of  $J$ , we have

$$\begin{aligned} & \limsup_{n \rightarrow \infty} \langle \varphi'_\gamma(u_n), u_n - u \rangle \leq \\ & - \liminf_{n \rightarrow \infty} \left( \langle \varphi'_\gamma(u_n), u - u_n \rangle + \mu J'(u_n; u - u_n) \right) \\ & + \mu \limsup_{n \rightarrow \infty} J'(u_n; u - u_n) \\ & \leq \mu \limsup_{n \rightarrow \infty} (J(u) - J(u_n)) = \mu(J(u) - \liminf_{n \rightarrow \infty} J(u_n)) \leq 0. \end{aligned}$$

Hence,

$$\limsup_{n \rightarrow \infty} \langle \varphi'_\gamma(u_n), u_n - u \rangle \leq 0. \quad (6.9)$$

From the expression (6.1) of  $\varphi'_\gamma$  we have

$$\begin{aligned} \langle \varphi'_{\eta_1}(u_n), u_n - u \rangle &= \langle \varphi'_\gamma(u_n), u_n - u \rangle \\ &+ (\eta_1 - \gamma) \int_0^T (h_p(u_n), u_n - u) dt. \end{aligned} \quad (6.10)$$

Taking into account that  $u_n \rightharpoonup u$  weakly in  $W^{1,p}$  and that the embedding  $W^{1,p} \hookrightarrow C$  is compact, we obtain for a subsequence of  $\{u_n\}$  (denoted again by  $\{u_n\}$ ) that

$$\lim_{n \rightarrow \infty} \int_0^T (h_p(u_n), u_n - u) dt = 0. \quad (6.11)$$

Then, by relations (6.9), (6.10) and (6.11) we have

$$\limsup_{n \rightarrow \infty} \langle \varphi'_{\eta_1}(u_n), u_n - u \rangle \leq 0. \quad (6.12)$$

On the other hand, for each positive integer  $n$ ,

$$0 \leq (\|u_n\|_{\eta_1}^{p-1} - \|u\|_{\eta_1}^{p-1})(\|u_n\|_{\eta_1} - \|u\|_{\eta_1}). \quad (6.13)$$

Using Hölder's inequality, we obtain

$$(\|u_n\|_{\eta_1}^{p-1} - \|u\|_{\eta_1}^{p-1})(\|u_n\|_{\eta_1} - \|u\|_{\eta_1}) \leq \langle \varphi'_{\eta_1}(u_n) - \varphi'_{\eta_1}(u), u_n - u \rangle.$$

Thus, by relation (6.13) we deduce that

$$\lim_{n \rightarrow \infty} \|u_n\|_{\eta_1} = \|u\|_{\eta_1}.$$

Since  $(W^{1,p}, \|\cdot\|_{\eta_1})$  is uniformly convex, it follows that  $\{u_n\}$  converges strongly to  $u$ .  $\square$

**Remark 6.1** Let  $\varepsilon > 0$  be arbitrary. From  $(F_1)$ ,  $(F_2)$  and  $(F_3)$  it follows that there exists  $\delta_1 > 0$  (depending on  $\varepsilon$ ) such that

$$F(t, x) - F(t, 0) \leq \varepsilon|x|^p + \alpha_{\delta_1}(t)\delta_1 \quad \text{for all } x \in \mathbb{R}^n, \text{ a.e. } t \in [0, T].$$

Then

$$\mathcal{F}(u) \geq -\varepsilon\|u\|_{L^p}^p - \delta_1\|\alpha_{\delta_1}\|_{L^1(0,T)} \quad \text{for all } u \in W^{1,p}. \quad (6.14)$$

**Proposition 6.2** Assume that  $F : [0, T] \times \mathbb{R}^n \rightarrow \mathbb{R}$  satisfies  $(F_1)$ ,  $(F_2)$  and  $(F_3)$  and that  $j : \mathbb{R}^n \times \mathbb{R}^n \rightarrow ]-\infty, +\infty]$  satisfies  $(J_1)$  and  $(J_2)$ . Then the following properties hold:

- (1).  $I(\lambda, \mu, \cdot)$  is weakly sequentially lower semi-continuous on  $W^{1,p}$  for each  $\lambda > 0$ ,  $\mu \geq 0$ ;
- (2).  $\lim_{\|u\|_{\eta_1} \rightarrow +\infty} I(\lambda, \mu, u) = +\infty$  for each  $\lambda > 0$ ,  $\mu \geq 0$ ;
- (3).  $\mathcal{E}(\lambda, \mu, \cdot)$  satisfies the (PS) condition for each  $\lambda, \mu > 0$ .

*Proof* (1) The function  $I(\lambda, \mu, \cdot)$  is weakly sequentially l.s.c on  $W^{1,p}$ , because  $\mathcal{F}$  is weakly sequentially l.s.c., while  $\varphi_\gamma$  and  $J$  are convex and l.s.c., hence they are also weakly sequentially l.s.c.

(2) We first observe that

$$\|u\|_{L^p}^p \leq \frac{1}{\eta_1}\|u\|_{\eta_1}^p \quad \text{for all } u \in W^{1,p}.$$

In (6.14) we choose  $\varepsilon < \frac{\eta_1}{2\lambda p}$ . Since the embedding  $L^p \hookrightarrow L^1$  is continuous and (6.7) holds, we have for all  $u \in D(J)$ ,

$$\begin{aligned} I(\lambda, \mu, u) &\geq \frac{1}{p} \left( \|u'\|_{L^p}^p + \gamma\|u\|_{L^p}^p \right) - \lambda\varepsilon\|u\|_{L^p}^p \\ &\quad - \lambda\delta_1\|\alpha_{\delta_1}\|_{L^1(0,T)} + \mu J(u) \\ &\geq \frac{\eta_1 - 2\varepsilon\lambda p}{2\eta_1 p} \|u\|_{\eta_1}^p - \lambda\delta_1\|\alpha_{\delta_1}\|_{L^1(0,T)} + \mu J(u). \end{aligned}$$

Since  $J$  is convex and l.s.c., it is bounded from below by an affine functional and then there exist constants  $c_1, c_2, c_3 > 0$  such that for all  $u \in D(J)$ ,

$$I(\lambda, \mu, u) \geq \frac{\eta_1 - 2\varepsilon\lambda p}{2\eta_1 p} \|u\|_{\eta_1}^p - \lambda\delta_1\|\alpha_{\delta_1}\|_{L^1(0,T)} - c_1|u(0)| - c_2|u(T)| - c_3.$$

By the continuity of the embedding  $W^{1,p} \hookrightarrow C$  we have for all  $u \in W^{1,p}$ ,

$$I(\lambda, \mu, u) \geq c_4\|u\|_{\eta_1}^p - c_5\|u\|_{\eta_1} - c_6,$$

where  $c_4, c_5, c_6$  are positive constants. Since  $p > 1$ , it follows that  $I(\lambda, \mu, u) \rightarrow +\infty$  as  $\|u\|_{\eta_1} \rightarrow +\infty$ .

(3) Let  $\{u_n\}$  in  $W^{1,p}$  be a sequence satisfying  $I(\lambda, \mu, u_n) \rightarrow c$ . Then, for all  $v \in W^{1,p}$ ,

$$\lambda \langle \mathcal{F}'(u_n), v - u_n \rangle + \varphi_\gamma(v) - \varphi_\gamma(u_n) + \mu J(v) - \mu J(u_n) \geq -\varepsilon_n \|v - u_n\|_{\eta_1},$$

where  $\{\varepsilon_n\} \subset [0, \infty)$  with  $\varepsilon_n \rightarrow 0$ . We have a subsequence  $\{u_n\} \subset D(J)$  (we just eliminate the finite number of elements of the sequence which do not belong to  $D(J)$ ), since  $\mu > 0$  and  $I(\lambda, \mu, u_n) \rightarrow c$ .

But  $I(\lambda, \mu, \cdot)$  is coercive, hence  $\{u_n\}$  is bounded in  $W^{1,p}$ . The embedding  $W^{1,p} \hookrightarrow C$  is compact, then we can find a subsequence, still denoted by  $\{u_n\}$ , which is weakly convergent to a point  $u \in W^{1,p}$  and strongly in  $C$ .

In the above inequality we take  $v = u_n + s(u - u_n)$ , with  $s > 0$ , then divide both sides of the inequality by  $s$  and let  $s \searrow 0$ . So, for all positive integer  $n$

$$\lambda \langle \mathcal{F}'(u_n), u - u_n \rangle + \langle \varphi'_\gamma(u_n), u - u_n \rangle + \mu J'(u_n; u - u_n) \geq -\varepsilon_n \|u - u_n\|_{\eta_1}.$$

By the upper semi-continuity of  $\mathcal{F}'$ , it follows that

$$\liminf_{n \rightarrow \infty} \left( \langle \varphi'_\gamma(u_n), u - u_n \rangle + \mu J'(u_n; u - u_n) \right) \geq 0.$$

By Lemma 6.1 we deduce that  $\{u_n\}$  has a subsequence that converges strongly to  $u \in W^{1,p}$ .  $\square$

**Remark 6.2** From  $(F_1)$ ,  $(F_2)$ ,  $(F_3)$  and  $(F_4)$  it follows that for each  $\varepsilon > 0$  there exist  $\delta_\varepsilon, \bar{\delta}_\varepsilon > 0$  such that

$$F(t, x) - F(t, 0) \leq \varepsilon |x|^p + \frac{\alpha_{\delta_\varepsilon}(t)}{\bar{\delta}_\varepsilon^{r-1}} |x|^r \quad \text{for all } x \in \mathbb{R}^n, \text{ a.e. } t \in [0, T],$$

where  $r \geq 1$ . Then, by using the continuity of the embedding  $W^{1,p} \hookrightarrow C$ , we obtain

$$\mathcal{F}(u) \geq -\varepsilon \|u\|_{L^p}^p - \frac{\hat{c}^r \|\alpha_{\delta_\varepsilon}\|_{L^1(0,T)}}{\bar{\delta}_\varepsilon^{r-1}} \|u\|_\gamma^r \quad \text{for all } u \in W^{1,p}. \quad (6.15)$$

**Remark 6.3** If  $F : [0, T] \times \mathbb{R}^n \rightarrow \mathbb{R}$  satisfies  $(F_1)$  and  $(F_4)$ , then  $0 = \nabla F(t, 0)$  for a.e.  $t \in [0, T]$ .

The main result of this section is the following.

**Theorem 6.1** *Let  $F : [0, T] \times \mathbb{R}^n \rightarrow \mathbb{R}$  be a function satisfying  $(F_1) - (F_5)$  and let  $j : \mathbb{R}^n \times \mathbb{R}^n \rightarrow ]-\infty, +\infty]$  be a function satisfying  $(J_1) - (J_3)$ . Then for each fixed  $\mu > 0$ , there exists an open interval  $\Lambda_\mu \subset ]0, +\infty[$*

such that for each  $\lambda \in \Lambda_\mu$ , the problem  $(P_{\lambda,\mu})$  has at least two nontrivial solutions.

*Proof* Let  $\mu > 0$  be fixed. We define the function  $g : ]0, +\infty[ \rightarrow \mathbb{R}$  by

$$g(t) = \sup \{ -\mathcal{F}(u) : \varphi_\gamma(u) + \mu J(u) \leq t \}, \text{ for all } t > 0.$$

Using (6.15) for  $r \in ]p, p^*[$  it follows that for all  $u \in W^{1,p}$  we have

$$-\mathcal{F}(u) \leq \frac{\varepsilon}{\gamma} \|u\|_\gamma^p + \frac{\hat{c}^r \|\alpha_{\delta_\varepsilon}\|_{L^1(0,T)}}{\bar{\delta}_\varepsilon^{r-1}} \|u\|_\gamma^r.$$

Since  $p < r$ , we have

$$\lim_{t \rightarrow 0^+} \frac{g(t)}{t} = 0.$$

Using  $(F_5)$  we define  $u_0(t) = s_0$  for a.e.  $t \in [0, T]$ . Then  $u_0 \in W^{1,p} \setminus \{0\}$  and  $-\mathcal{F}(u_0) > 0$ . Due to the convergence relation above, it is possible to choose a real number  $t_0$  such that  $0 < t_0 < \varphi_\gamma(u_0) + \mu J(u_0)$  and

$$\frac{g(t_0)}{t_0} < [\varphi_\gamma(u_0) + \mu J(u_0)]^{-1} \cdot (-\mathcal{F}(u_0)).$$

We choose  $\rho_0 > 0$  such that

$$g(t_0) < \rho_0 < [\varphi_\gamma(u_0) + \mu J(u_0)]^{-1} \cdot (-\mathcal{F}(u_0)) t_0. \quad (6.16)$$

We apply Theorem 1.26 to the space  $W^{1,p}$ , the interval  $\Lambda = ]0, +\infty[$  and the functions  $E_1, E_2 : W^{1,p} \rightarrow \mathbb{R}, h : \Lambda \rightarrow \mathbb{R}$  defined by

$$E_1(u) = \varphi_\gamma(u), \zeta_1(u) = \mu J(u), I_2(u) = E_2(u) = \mathcal{F}(u), h(\lambda) = \rho_0 \lambda.$$

By Proposition 6.2, the assumptions (i) and (ii) of Theorem 1.26 are fulfilled.

We prove now the minimax inequality

$$\begin{aligned} & \sup_{\lambda \in \Lambda} \inf_{u \in W^{1,p}} \left( \varphi_\gamma(u) + \mu J(u) + \lambda \mathcal{F}(u) + \rho_0 \lambda \right) < \\ & \inf_{u \in W^{1,p}} \sup_{\lambda \in \Lambda} \left( \varphi_\gamma(u) + \mu J(u) + \lambda \mathcal{F}(u) + \rho_0 \lambda \right). \end{aligned}$$

The function

$$\lambda \mapsto \inf_{u \in W^{1,p}} \left( \varphi_\gamma(u) + \mu J(u) + \lambda \mathcal{F}(u) + \rho_0 \lambda \right)$$

is upper semi-continuous on  $\Lambda$ . Since

$$\inf_{u \in W^{1,p}} \left( \varphi_\gamma(u) + \mu J(u) + \lambda \mathcal{F}(u) + \rho_0 \lambda \right) \leq \varphi_\gamma(u_0) + \mu J(u_0) + \lambda \mathcal{F}(u_0) + \rho_0 \lambda$$

and  $\rho_0 < -\mathcal{F}(u_0)$ , it follows that

$$\lim_{\lambda \rightarrow +\infty} \inf_{u \in W^{1,p}} \left( \varphi_\gamma(u) + \mu J(u) + \lambda \mathcal{F}(u) + \rho_0 \lambda \right) = -\infty.$$

Thus we can find  $\bar{\lambda} \in \Lambda$  such that

$$\begin{aligned} \beta_1 &:= \sup_{\lambda \in \Lambda} \inf_{u \in W^{1,p}} \left( \varphi_\gamma(u) + \mu J(u) + \lambda \mathcal{F}(u) + \rho_0 \lambda \right) \\ &= \inf_{u \in W^{1,p}} \left( \varphi_\gamma(u) + \mu J(u) + \bar{\lambda} \mathcal{F}(u) + \rho_0 \bar{\lambda} \right). \end{aligned}$$

In order to prove that  $\beta_1 < t_0$ , we distinguish two cases:

I. If  $0 \leq \bar{\lambda} < \frac{t_0}{\rho_0}$ , we have

$$\beta_1 \leq \varphi_\gamma(0) + \mu J(0) + \bar{\lambda} \mathcal{F}(0) + \rho_0 \bar{\lambda} = \bar{\lambda} \rho_0 < t_0.$$

II. If  $\bar{\lambda} \geq \frac{t_0}{\rho_0}$ , then we use  $\rho_0 < -\mathcal{F}(u_0)$  and the inequality (6.16) to get

$$\begin{aligned} \eta_1 &\leq \varphi_\gamma(u_0) + \mu J(u_0) + \bar{\lambda} \mathcal{F}(u_0) + \rho_0 \bar{\lambda} \\ &\leq \varphi_\gamma(u_0) + \mu J(u_0) + \frac{t_0}{\rho_0} (\rho_0 + \mathcal{F}(u_0)) < t_0. \end{aligned}$$

From  $g(t_0) < \rho_0$  it follows that for all  $u \in W^{1,p}$  with  $\varphi_\gamma(u) + \mu J(u) \leq t_0$  we have  $-\mathcal{F}(u) < \rho_0$ . Hence

$$t_0 \leq \inf \{ \varphi_\mathcal{G}(u) + \mu J(u) : -\mathcal{F}(u) \geq \rho_0 \}.$$

On the other hand,

$$\begin{aligned} \beta_2 &= \inf_{u \in W^{1,p}} \sup_{\lambda \in \Lambda} (\varphi_\mathcal{G}(u) + \mu J(u) + \lambda \mathcal{F}(u) + \rho_0 \lambda) \\ &= \inf \{ \varphi_\mathcal{G}(u) + \mu J(u) : -\mathcal{F}(u) \geq \rho_0 \}. \end{aligned}$$

We conclude that

$$\beta_1 < t_0 \leq \beta_2.$$

Hence, assumption (iii) from Theorem 1.26 holds. Thus, there exists an open interval  $\Lambda_\mu \subseteq ]0, \infty)$  such that for each  $\lambda \in \Lambda_\mu$  the function  $\varphi_\mathcal{G} + \mu J + \lambda \mathcal{F}$  has at least three critical points in  $W^{1,p}$ . By Proposition 6.1 it follows that these critical points are solutions of  $(P_{\lambda,\mu})$ . Since  $\nabla F(t, 0) = 0$  for a.e.  $t \in [0, T]$ , we get that at least two of the above solutions are nontrivial.  $\square$

**Remark 6.4** The two conditions from  $(J_3)$  can be replaced by  $(J'_3)$   $j(x, y) \geq j(0, 0)$  for all  $(x, y) \in \mathbb{R}^n \times \mathbb{R}^n$ .

Then all the proofs above can be adapted by considering

$$J(u) = j(u(0), u(T)) - j(0, 0).$$

**Corollary 6.1** *Let  $F : [0, T] \times \mathbb{R}^n \rightarrow \mathbb{R}$  be a function satisfying  $(F_1) - (F_5)$  and let  $b : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$  be a positive, convex and Gâteaux differentiable function with  $b(0, 0) = 0$ . Assume that  $S \subset \mathbb{R}^n \times \mathbb{R}^n$  is a nonempty closed convex cone with  $S \neq \{(0, 0)\}$ , whose normal cone we denote by  $N_S$ . Then, for each fixed  $\gamma, \mu > 0$ , there exists an open interval  $\Lambda_0 \subset ]0, +\infty[$  such that for each  $\lambda \in \Lambda_0$ , the following problem*

$$(\hat{P}_{\lambda, \mu}) \quad \begin{cases} -[h_p(u')] + \gamma h_p(u) = \lambda \bar{\partial} F(t, u) \text{ a.e. } t \in [0, T], \\ (u(0), u(T)) \in S, \\ \left( h_p(u')(0), -h_p(u')(T) \right) \in \mu \nabla b(u(0), u(T)) + \mu N_S(u(0), u(T)), \end{cases}$$

*has at least two nontrivial solutions.*

*Proof* The statement follows by applying Theorem 6.1 to the function  $F$  and the convex function  $j : \mathbb{R}^n \times \mathbb{R}^n \rightarrow ]-\infty, +\infty]$  defined by

$$j(x, y) = b(x, y) + I_S(x, y), \text{ for all } (x, y) \in \mathbb{R}^n \times \mathbb{R}^n,$$

where

$$I_S(x, y) = \begin{cases} 0, & \text{if } (x, y) \in S \\ +\infty, & \text{if } (x, y) \in \mathbb{R}^n \times \mathbb{R}^n \setminus S, \end{cases}$$

is the indicator function of the cone  $S$ .

Note, that in this case  $D(j) = S$  and  $j$  satisfies the conditions  $(J_1) - (J_3)$ . Moreover,

$$\partial j(x, y) = \nabla b(x, y) + \partial I_S(x, y) = \nabla b(x, y) + N_S(x, y) \text{ for all } (x, y) \in S.$$

□

**Example 6.1** Various possible choices of  $b$  and  $S$  from Corollary 6.1 recover some classical boundary conditions. For instance:

- (a)  $b = 0$  and  $S = \{(x, x) : x \in \mathbb{R}^n\}$  we get periodic boundary conditions  $u(0) = u(T), u'(0) = u'(T)$ ;
- (b)  $b = 0$  and  $S = \mathbb{R}^n \times \mathbb{R}^n$  we get Neumann type boundary conditions  $u'(0) = u'(T) = 0$ ;
- (c)  $b(z) = \frac{1}{2}(Az, z)_{\mathbb{R}^{2N}}$ ,  $z \in \mathbb{R}^{2N}$ , where  $A$  is a symmetric, positive  $2N \times 2N$  real valued matrix, and  $S = \mathbb{R}^n \times \mathbb{R}^n$ ; we get the following mixed boundary conditions

$$\begin{pmatrix} h_p(u')(0) \\ -h_p(u')(T) \end{pmatrix} = A \begin{pmatrix} u(0) \\ u(T) \end{pmatrix}.$$

For these choices of  $F$ ,  $b$  and  $S$  it follows by Corollary 6.1 that for each fixed  $\gamma, \mu > 0$ , there exists an open interval  $\Lambda_0 \subset ]0, +\infty[$  such that for each  $\lambda \in \Lambda_0$  the problem  $(\hat{P}_{\lambda, \mu})$  has at least two nontrivial solutions.

### 6.3 Scalar field systems with nonlinear oscillatory terms

To formulate the second problem we assume that:

- (C1)  $F : \mathbb{R}^n \rightarrow \mathbb{R}$  is of class  $C^1$ ,
- (C2)  $S \subset \mathbb{R}^n \times \mathbb{R}^n$  is a closed convex set with  $\{(x, x) : x \in \mathbb{R}^n\} \subset S$ ,
- (C3)  $\gamma_1, \dots, \gamma_n \in L^\infty(]0, T[, \mathbb{R})$  are so that  $\text{essinf } \gamma_i > 0$ , for all  $i = 1, \dots, n$  and put  $\gamma = (\gamma_1, \dots, \gamma_n)$ ,
- (C4)  $\alpha \in L^1(]0, T[, \mathbb{R})$  is so that  $\alpha(t) \geq 0$  a.e. in  $]0, T[$ .

We study the existence and multiplicity of the solutions of the system:

$$(S) \quad \begin{cases} -[h_p(u')] + \gamma \circ h_p(u) = \alpha(t)\nabla F(u) \\ (h_p(u')(0), -h_p(u')(T)) \in N_S(u(0), u(T)), \end{cases}$$

where  $N_S(x, y)$  denotes the normal cone of  $S$  at  $(x, y) \in S$ .

The first step in the study of  $(S)$  is to establish the corresponding energy function. For this some preparation is needed:

- For every  $i \in \{1, \dots, n\}$  denote by  $X_i$  the Sobolev space  $W^{1,p}(]0, T[, \mathbb{R})$  equipped with the norm  $\|\cdot\|_i$ , where

$$\|f\|_i = \left( \int_0^T \gamma_i(t)|f(t)|^p dt + \int_0^T |f'(t)|^p dt \right)^{\frac{1}{p}}.$$

Since  $\text{essinf } \gamma_i > 0$  it follows that  $\|\cdot\|_i$  is equivalent to the usual norm on  $W^{1,p}(]0, T[, \mathbb{R})$ .

- $X$  is the Sobolev space  $W^{1,p}(]0, T[, \mathbb{R}^n)$  equipped with the usual norm

$$\|u\| = \left( \int_0^T |u(t)|^p dt + \int_0^T |u'(t)|^p dt \right)^{1/p}.$$

We observe that  $X$  is isomorphic to  $W^{1,p}(]0, T[, \mathbb{R})^n$ .

- Let  $Y$  be the real Banach space  $C([0, T], \mathbb{R}^n)$  endowed with the supremum norm  $\|\cdot\|_s$ . For simplicity we will denote the supremum norm on every space  $C([0, T], \mathbb{R}^m)$ ,  $m \in \mathbb{N}$ , with the same symbol  $\|\cdot\|_s$ .

- Put

$$\Sigma := \{u \in X \mid (u(0), u(T)) \in S\}.$$

Note that  $\Sigma$  is a closed convex subset of  $X$  containing the constant functions.

- Define  $E_1: Y \rightarrow \mathbb{R}$  by

$$E_1(u) = \int_0^T \alpha(t)F(u(t))dt, \text{ for every } u \in Y,$$

$E_2: X \rightarrow \mathbb{R}$  by

$$E_2(u) = \frac{1}{p}(\|u_1\|_1^p + \dots + \|u_n\|_n^p), \text{ for every } u \in X,$$

and  $\zeta_1: X \rightarrow ]-\infty, +\infty]$  by

$$\zeta_1(u) = \begin{cases} 0, & u \in \Sigma \\ +\infty, & u \notin \Sigma. \end{cases}$$

The notations used above are inspired from Theorem 1.27.

**Lemma 6.2** *The following assertions hold:*

- (a) *The above defined maps  $E_1: X \rightarrow \mathbb{R}$  and  $E_2: X \rightarrow \mathbb{R}$  are of class  $C^1$  and  $\langle E_1'(u); v \rangle = \int_0^T \alpha(t)f(u(t))v(t)dt$ , for every  $u, v \in X$ .*
- (b) *The map  $\zeta_1: X \rightarrow ]-\infty, +\infty]$  is convex, proper, and lower semi-continuous.*

Define now  $I_1: X \rightarrow ]-\infty, +\infty]$  and  $I_2: X \rightarrow \mathbb{R}$  by

$$I_1(u) = E_1(u) + \zeta_1(u), \quad I_2(u) = E_2(u) \quad \text{for every } u \in X. \quad (6.17)$$

The energy functional associated to the system (S) is given by  $I_1 + I_2: X \rightarrow ]-\infty, +\infty]$  is given by  $I_1 + I_2 = (E_1 + E_2) + \zeta_1$ .

Our aim is to apply Theorem 1.27 to the maps  $I_1$  and  $I_2$  defined in (6.17). Indeed,

- $X, Y$  are real Banach spaces,  $X$  is reflexive, and  $X$  is compactly embedded in  $Y$ .
- $E_1: Y \rightarrow \mathbb{R}$ ,  $E_2: X \rightarrow \mathbb{R}$  are locally Lipschitz, and  $\zeta_1$  is convex, proper, and lower semi-continuous (according to Lemma 6.2).
- $\Psi = E_2: X \rightarrow \mathbb{R}$  is weakly sequentially lower semi-continuous (being convex and continuous) and coercive.

- Since  $\inf_X \Psi = 0$  and since  $\Sigma$  contains the constant functions, condition (1.36) is satisfied.

Furthermore we have to introduce some suitable subsets of  $\mathbb{R}^n$ . For this note that, since  $X_i$  is embedded in  $C([0, T], \mathbb{R})$ , there exist  $c_i > 0$ ,  $i = \overline{1, n}$ , such that

$$\|f\|_s \leq c_i \|f\|_i, \quad \text{for every } f \in X_i. \quad (6.18)$$

For every  $r > 0$ , let

$$\begin{aligned} A(r) &:= \left\{ x \in \mathbb{R}^n : \frac{1}{p} \sum_{i=1}^n \frac{1}{c_i^p} |x_i|^p \leq r \right\} \\ B(r) &:= \left\{ x \in \mathbb{R}^n : \frac{1}{p} \sum_{i=1}^n |x_i|^p \int_0^T \gamma_i(t) dt \leq r \right\}. \end{aligned} \quad (6.19)$$

**Remark 6.5** 1) For every  $r > 0$  the inclusion  $B(r) \subseteq A(r)$  holds. To see this, we observe that relation (6.18) implies

$$1 \leq c_i \left( \int_0^T \gamma_i(t) dt \right)^{1/p}, \quad \text{for } i = \overline{1, n}.$$

Pick now an arbitrary  $x \in B(r)$ . Then

$$\frac{1}{p} \sum_{i=1}^n \frac{1}{c_i^p} |x_i|^p \leq \frac{1}{p} \sum_{i=1}^n |x_i|^p \int_0^T \gamma_i(t) dt \leq r,$$

hence  $x \in A(r)$ .

2) Since the map  $x \in \mathbb{R}^n \mapsto \sum_{i=1}^n |x_i|^p \int_0^T \gamma_i(t) dt \in \mathbb{R}$  is convex, we have for every  $r > 0$ ,

$$\text{int } B(r) = \left\{ x \in \mathbb{R}^n : \frac{1}{p} \sum_{i=1}^n |x_i|^p \int_0^T \gamma_i(t) dt < r \right\}.$$

**Theorem 6.2** *Let  $r > 0$  be such that*

$$\min_{x \in A(r)} F(x) = \min_{x \in \text{int } B(r)} F(x).$$

*Then the following assertions hold:*

- (a)  $\varphi(r) = 0$ , where  $\varphi$  is defined by relation (1.37).
- (b) Problem (P) has a solution  $u \in \Sigma$  satisfying  $\Psi(u) < r$ .

*Proof* (a) We have  $\varphi(r) \geq 0$ , by definition. To show the converse inequality choose  $x^0 \in \text{int } B(r)$  so that

$$F(x^0) = \min_{x \in \text{int } B(r)} F(x) = \min_{x \in A(r)} F(x),$$

and let  $u^0: X \rightarrow \mathbb{R}^n$  be the function taking the constant value  $x^0$ . For every  $i \in \{1, \dots, n\}$  we have

$$\|u_i^0\|_i = |x_i^0| \left( \int_0^T \gamma_i(t) dt \right)^{1/p}.$$

Therefore

$$\psi(u^0) = \frac{1}{p} \sum_{i=1}^n |x_i^0|^p \int_0^T \gamma_i(t) dt < r,$$

hence  $u^0 \in I_2^{-1}(] - \infty, r[)$ . Since  $I_2^{-1}(] - \infty, r[)$  is convex and closed in the norm topology, it is closed also in the weak topology, hence

$$\overline{(I_2^{-1}(] - \infty, r[))}_w \subseteq I_2^{-1}(] - \infty, r[).$$

Pick now an arbitrary element  $v \in \overline{(I_2^{-1}(] - \infty, r[))}_w$ . Then  $I_2(v) \leq r$ . Thus, by (6.18), for all  $t \in [0, T]$

$$\frac{1}{p} \sum_{i=1}^n \frac{1}{c_i^p} |v_i(t)|^p \leq \frac{1}{p} \sum_{i=1}^n \frac{1}{c_i^p} \|v_i\|_s^p \leq \frac{1}{p} \sum_{i=1}^n \|v_i\|_i^p = I_2(v) \leq r.$$

We conclude that  $v(t) \in A(r)$  for every  $t \in [0, T]$ . Hence

$$F(x^0) \leq F(v(t)), \text{ for every } t \in [0, T].$$

It follows that

$$E_1(u^0) = \int_0^T \alpha(t) F(x^0) dt \leq \int_0^T \alpha(t) F(v(t)) dt = E_1(v) \leq I_1(v).$$

Since  $v \in \overline{(I_2^{-1}(] - \infty, r[))}_w$  was chosen arbitrarily, we conclude that

$$\inf_{v \in \overline{(I_2^{-1}(] - \infty, r[))}_w} I_1(v) = I_1(u^0).$$

This implies, according to the definition of  $\varphi$  in (1.37),

$$\varphi(r) \leq \frac{I_1(u^0) - I_1(u^0)}{r - I_2(u^0)} = 0,$$

hence  $\varphi(r) = 0$ .

(b) Since  $\varphi(r) = 0$ , we can apply Theorem 1.27 for  $\lambda = 1$ , and conclude

that the map  $I_1 + I_2$  has a critical point  $u$  lying in  $\Sigma$  and such that  $I_2(u) < r$ . The assertion follows from Proposition 6.1.  $\square$

**Theorem 6.3** *Assume that:*

- (1) *There exists a sequence  $(r_k)_{k \in \mathbb{N}}$  of positive reals such that  $\lim r_k = +\infty$  and*

$$\min_{x \in A(r_k)} F(x) = \min_{x \in \text{int}B(r_k)} F(x), \text{ for every } k \in \mathbb{N}.$$

- (2) *The following inequality holds*

$$\liminf_{|x| \rightarrow +\infty} \frac{F(x) \int_0^T \alpha(t) dt}{\sum_{i=1}^n |x_i|^p \int_0^T \gamma_i(t) dt} < -\frac{1}{p}.$$

*Then problem (P) has an unbounded sequence of solutions.*

*Proof* Assumption (1) implies, according to Theorem 6.2(a), that  $\varphi(r_k) = 0$ , for every  $k \in \mathbb{N}$ . Let  $\gamma$  be defined as in relation (1.38). Since  $\varphi(r) \geq 0$  for every  $r > 0$ , we conclude that

$$\gamma = \liminf_{r \rightarrow +\infty} \varphi(r) = 0.$$

Applying Theorem 1.27(a) for  $\lambda = 1$ , we conclude that either assertion (b1) or (b2) of this theorem must hold. Next we show that (b1) is not satisfied, that is, we prove that  $I_1 + I_2$  is unbounded below. For this fix a real number  $q$  such that

$$\liminf_{|x| \rightarrow +\infty} \frac{F(x) \int_0^T \alpha(t) dt}{\sum_{i=1}^n |x_i|^p \int_0^T \gamma_i(t) dt} < q < -\frac{1}{p}.$$

Choose now a sequence  $(x^k)_{k \in \mathbb{N}}$  in  $\mathbb{R}^n$  such that  $\lim |x^k| = +\infty$  and

$$\frac{F(x^k) \int_0^T \alpha(t) dt}{\sum_{i=1}^n |x_i^k|^p \int_0^T \gamma_i(t) dt} < q, \text{ for every } k \in \mathbb{N}.$$

For every  $k \in \mathbb{N}$  denote by  $u^k: X \rightarrow \mathbb{R}^n$  the constant function taking the value  $x^k$ . Then the following relations hold for every  $k \in \mathbb{N}$

$$\begin{aligned} I_1(u^k) + I_2(u^k) &= F(x^k) \int_0^T \alpha(t) dt + \frac{1}{p} \sum_{i=1}^n |x_i^k|^p \int_0^T \gamma_i(t) dt \\ &< \left( q + \frac{1}{p} \right) \sum_{i=1}^n |x_i^k|^p \int_0^T \gamma_i(t) dt. \end{aligned}$$

Since  $|x^k| \rightarrow +\infty$ ,  $\int_0^T \gamma_i(t)dt > 0$ , for every  $i \in \{1, \dots, n\}$ , and  $q + \frac{1}{p} < 0$ , we conclude that  $\lim_{k \rightarrow \infty} (I_1(u^k) + I_2(u^k)) = -\infty$ , thus  $I_1 + I_2$  is unbounded below. The assertion follows now from Theorem 1.27(b2), the definition of  $I_2$ , and Proposition 6.1.  $\square$

**Theorem 6.4** *Assume that:*

- (1) *There exists a sequence  $(r_k)_{k \in \mathbb{N}}$  of positive reals such that  $\lim r_k = 0$  and*

$$\min_{x \in A(r_k)} F(x) = \min_{x \in \text{int}B(r_k)} F(x), \text{ for every } k \in \mathbb{N}.$$

- (2) *The following inequality holds*

$$\liminf_{x \rightarrow 0_n} \frac{F(x) \int_0^T \alpha(t)dt}{\sum_{i=1}^n |x_i|^p \int_0^T \gamma_i(t)dt} < -\frac{1}{p}.$$

*Then problem (S) has a sequence of pairwise distinct solutions which converges strongly to the zero function  $\theta_X \in X$ .*

*Proof* We first observe that  $\theta_X$  is the only global minimum of  $I_2$ . Assumption (1) and Theorem 6.2(a) imply that  $\varphi(r_k) = 0$ , for every  $k \in \mathbb{N}$ . Let  $\delta$  be defined as in relation (1.39). Since  $\varphi(r) \geq 0$  for every  $r > 0$ , we conclude that

$$\delta = \liminf_{r \rightarrow 0^+} \varphi(r) = 0.$$

Applying Theorem 1.27(c) for  $\lambda = 1$ , we conclude that either assertion (c1) or (c2) of this theorem must hold. Next we show that (c1) is not satisfied, that is, we prove that  $\theta_X$  is not a local minimum of  $I_1 + I_2$ . For this fix a real number  $q$  such that

$$\liminf_{x \rightarrow 0_n} \frac{F(x) \int_0^T \alpha(t)dt}{\sum_{i=1}^n |x_i|^p \int_0^T \gamma_i(t)dt} < q < -\frac{1}{p}.$$

Choose now a sequence  $(x^k)_{k \in \mathbb{N}}$  in  $\mathbb{R}^n$  such that  $\lim x^k = 0_n$  and

$$\frac{F(x^k) \int_0^T \alpha(t)dt}{\sum_{i=1}^n |x_i^k|^p \int_0^T \gamma_i(t)dt} < q, \text{ for every } k \in \mathbb{N}.$$

For every  $k \in \mathbb{N}$  denote by  $u^k: X \rightarrow \mathbb{R}^n$  the constant function taking

the value  $x^k$ . Then the following relations hold for every  $k \in \mathbb{N}$

$$\begin{aligned} I_1(u^k) + I_2(u^k) &= F(x^k) \int_0^T \alpha(t) dt + \frac{1}{p} \sum_{i=1}^n |x_i^k|^p \int_0^T \gamma_i(t) dt \\ &< \left( q + \frac{1}{p} \right) \sum_{i=1}^n |x_i^k|^p \int_0^T \gamma_i(t) dt \\ &\leq 0 = I_1(\theta_X) + I_2(\theta_X). \end{aligned}$$

We have that  $\lim \|u^k\| = 0$ , thus  $\theta_X$  is not a local minimum of  $I_1 + I_2$ . Theorem 1.27(c2) and Proposition 6.1 imply now the existence of a sequence  $(\tilde{u}^k)$  of pairwise distinct solutions of  $(S)$  such that  $\lim I_2(\tilde{u}^k) = 0$ . Since  $I_2$  is a norm on  $X$  which is equivalent to the norm  $\|\cdot\|$ , we conclude that  $(\tilde{u}^k)$  converges strongly to  $\theta_X$  in  $X$ .  $\square$

## 6.4 Applications

We specialize now some of the data of the previous section in order to obtain applications of Theorems 6.3 and 6.4. We assume that  $n \geq 1$  is a natural number,  $T = 1$ ,  $p > 1$  is a real number,  $S \subseteq \mathbb{R}^n \times \mathbb{R}^n$  is a set satisfying condition (C2),  $\gamma_i$  is the function taking the constant value 1, for every  $i \in \{1, \dots, n\}$ , and  $\alpha \in L^1(]0, T[, \mathbb{R})$  is so that  $\int_0^1 \alpha(t) dt > 1$  and  $\alpha(t) \geq 0$  a.e. in  $]0, 1[$ . In this case every norm  $\|\cdot\|_i$ ,  $i = \overline{1, n}$ , reduces to the usual norm on  $W^{1,p}(]0, 1[, \mathbb{R})$ , and all the constants  $c_i$ ,  $i = \overline{1, n}$ , in (6.18) can be considered to be equal to a suitable real number  $c > 0$ . Furthermore, we assume in this section that  $\mathbb{R}^n$  is endowed with the  $p$ -norm

$$|x| = \left( \sum_{i=1}^n |x_i|^p \right)^{1/p}.$$

Thus, for  $r > 0$ , the sets  $A(r)$  and  $B(r)$  defined in (6.19) become

$$A(r) = \left\{ x \in \mathbb{R}^n : \frac{1}{p} \cdot \frac{1}{c^p} |x|^p \leq r \right\} \quad B(r) = \left\{ x \in \mathbb{R}^n : \frac{1}{p} |x|^p \leq r \right\}.$$

**Example 6.2** We give now an application of Theorem 6.3. In order to define  $F: \mathbb{R}^n \rightarrow \mathbb{R}$  we consider a function  $f: [0, +\infty[ \rightarrow [0, +\infty[$  with the following properties:

- (i)  $f$  is bijective,
- (ii)  $f$  is strictly increasing,

- (iii)  $\lim_{t \rightarrow +\infty} \frac{f(t + \pi)}{f(t)} > c^p$ ,  
 (iv)  $f^{-1}$  is locally Lipschitz.

Note that the above properties imply that  $f(0) = 0$ ,  $f^{-1}$  is strictly increasing, and

$$\lim_{t \rightarrow +\infty} f(t) = \lim_{t \rightarrow +\infty} f^{-1}(t) = +\infty.$$

An example of a function  $f$  satisfying properties (i)–(iv) is

$$t \in [0, +\infty[ \mapsto a^t - 1 \in [0, +\infty[,$$

where the real  $a > 1$  is chosen so that  $a^\pi > c^p$ .

Define  $F: \mathbb{R}^n \rightarrow \mathbb{R}$  by

$$F(x) = \frac{1}{p} |x|^p \sin f^{-1}(|x|^p), \text{ for every } x \in \mathbb{R}^n.$$

A straightforward argument yields that  $F$  is of class  $C^1$  (that is, satisfies condition (C1)).

We next show that the assumptions (1) and (2) of Theorem 6.3 are fulfilled. Indeed, to check (1), we observe that by (iii) and for every sufficiently large  $k \in \mathbb{N}$ ,

$$\frac{f((2k+1)\pi)}{f(2k\pi)} > c^p. \quad (6.20)$$

For these values of  $k$  put

$$r_k := \frac{f((2k+1)\pi)}{pc^p}.$$

Then  $\lim r_k = +\infty$ . Furthermore,

$$\min_{x \in A(r_k)} F(x) = \min_{|x|^p \leq f((2k+1)\pi)} F(x). \quad (6.21)$$

If  $f(2k\pi) \leq |x|^p \leq f((2k+1)\pi)$ , then

$$2k\pi \leq f^{-1}(|x|^p) \leq (2k+1)\pi,$$

hence

$$\sin(f^{-1}(|x|^p)) \geq 0,$$

This means that  $F \geq 0$ . Taking into account that  $F(0_n) = 0$ , it follows that

$$\min_{|x|^p \leq f((2k+1)\pi)} F(x) = \min_{|x|^p \leq f(2k\pi)} F(x). \quad (6.22)$$

On the other hand, if  $x \in \mathbb{R}^n$  is so that  $|x|^p \leq f(2k\pi)$  then, in view of (6.20), we have

$$\frac{1}{p}|x|^p \leq \frac{1}{p}f(2k\pi) < \frac{f((2k+1)\pi)}{pc^p} = r_k. \quad (6.23)$$

Using (6.21), (6.22), and (6.23), we conclude that

$$\min_{x \in A(r_k)} F(x) = \min_{x \in \text{int}B(r_k)} F(x),$$

hence assumption (1) of Theorem 6.3 is fulfilled. For assumption (2) we observe that

$$\begin{aligned} \liminf_{|x| \rightarrow +\infty} \frac{F(x) \int_0^1 \alpha(t) dt}{\sum_{i=1}^n |x_i|^p} &= \liminf_{|x| \rightarrow +\infty} \frac{1}{p} \sin(f^{-1}(|x|^p)) \int_0^1 \alpha(t) dt \\ &= -\frac{1}{p} \int_0^1 \alpha(t) dt < -\frac{1}{p}. \end{aligned}$$

Theorem 6.3 yields now that problem (P) has an unbounded sequence of solutions.

**Example 6.3** To get an application of Theorem 6.4 consider a function  $f: ]0, +\infty[ \rightarrow ]0, +\infty[$  with the following properties:

- (i)  $f$  is surjective,
- (ii)  $f$  is strictly increasing,
- (iii)  $f$  is differentiable on  $]0, +\infty[$ ,
- (iv) the map  $t \in ]0, +\infty[ \mapsto \frac{f'(\frac{1}{t})}{t} \in \mathbb{R}$  is bounded on every interval  $]0, a[$ ,  $a > 0$ ,
- (v)  $\lim_{t \rightarrow +\infty} \frac{f^{-1}(t + \pi)}{f^{-1}(t)} > c^p$ .

Under these assumptions,  $f^{-1}$  is strictly increasing and  $\lim_{t \rightarrow +\infty} f^{-1}(t) = +\infty$ . An example for a function satisfying properties (i)–(v) is

$$t \in ]0, +\infty[ \mapsto \log_a(t + 1) \in ]0, +\infty[,$$

where the real  $a > 1$  is chosen so that  $a^\pi > c^p$ .

Define  $g: [0, +\infty[ \rightarrow \mathbb{R}$  by

$$g(t) = \begin{cases} t \sin f(1/t), & t \neq 0 \\ 0, & t = 0. \end{cases}$$

This map has the following properties:

- $g$  is continuous on  $[0, +\infty[$ ,

- $g$  is differentiable on  $]0, +\infty[$  and  $g$  is not differentiable in  $0$ ,
- $g'$  is bounded on every interval  $]0, a[$ ,  $a > 0$ . This follows from property (iv) of  $f$  and the fact that for every  $t > 0$  we have  $g'(t) = \sin f(1/t) - \cos f(1/t) \frac{f'(1/t)}{t}$ .

Define now  $F: \mathbb{R}^n \rightarrow \mathbb{R}$  by

$$F(x) = \frac{1}{p} g(|x|^p), \quad \text{for every } x \in \mathbb{R}^n.$$

Since  $p > 1$ , it follows that  $F$  is of class  $C^1$ .

We next show that  $F$  satisfies conditions (1) and (2) of Theorem 6.4. To verify (1), we observe that (by (v)) and for every sufficiently large  $k \in \mathbb{N}$

$$\frac{f^{-1}((2k+1)\pi)}{f^{-1}(2k\pi)} > c^p. \quad (6.24)$$

For these values of  $k$  put

$$r_k := \frac{1}{pc^p} \cdot \frac{1}{f^{-1}(2k\pi)}.$$

Then  $\lim r_k = 0$ . Furthermore,

$$\min_{x \in A(r_k)} F(x) = \min_{|x|^p \leq \frac{1}{f^{-1}(2k\pi)}} F(x). \quad (6.25)$$

If  $\frac{1}{f^{-1}((2k+1)\pi)} \leq |x|^p \leq \frac{1}{f^{-1}(2k\pi)}$ , then

$$2k\pi \leq f\left(\frac{1}{|x|^p}\right) \leq (2k+1)\pi,$$

hence

$$\sin f\left(\frac{1}{|x|^p}\right) \geq 0.$$

Therefore  $F \geq 0$ . Taking into account that  $F(0_n) = 0$ , it follows that

$$\min_{|x|^p \leq \frac{1}{f^{-1}(2k\pi)}} F(x) = \min_{|x|^p \leq \frac{1}{f^{-1}((2k+1)\pi)}} F(x). \quad (6.26)$$

On the other hand, if  $x \in \mathbb{R}^n$  is so that  $|x|^p \leq \frac{1}{f^{-1}((2k+1)\pi)}$ , then, in view of (6.24), we have that

$$\frac{1}{p}|x|^p \leq \frac{1}{p} \frac{1}{f^{-1}((2k+1)\pi)} < \frac{1}{pc^p} \cdot \frac{1}{f^{-1}(2k\pi)} = r_k. \quad (6.27)$$

Using relations (6.25), (6.26), and (6.27), we conclude that

$$\min_{x \in A(r_k)} F(x) = \min_{x \in \text{int}B(r_k)} F(x),$$

hence assumption (1) of Theorem 6.4 is fulfilled. For assumption (2) we observe that

$$\begin{aligned} \liminf_{x \rightarrow 0_n} \frac{F(x) \int_0^1 \alpha(t) dt}{\sum_{i=1}^n |x_i|^p} &= \liminf_{x \rightarrow 0_n} \frac{1}{p} \sin f \left( \frac{1}{|x|^p} \right) \int_0^1 \alpha(t) dt \\ &= -\frac{1}{p} \int_0^1 \alpha(t) dt < -\frac{1}{p}. \end{aligned}$$

According to Theorem 6.4, problem (P) has a sequence of pairwise distinct solutions which converges strongly to the zero function  $\theta_X \in X$ .

### 6.5 Comments and historical notes

In the last few years, many papers were dedicated to the study of  $p$ -Laplacian systems with various types of boundary conditions. We mention here Manásevich and Mawhin [200], [201], Mawhin [208], Gasinski and Papageorgiu [126], Jebelean and Moroşanu [149], [150]. The methods used in these papers to prove the existence or multiple solutions of  $p$ -Laplacian systems are based on degree theory, minimax result, fixed point theorems, or on continuation methods of Leray-Schauder type. Jebelean [148] used the symmetric version of the mountain pass theorem to prove the existence of infinitely many solutions for problem  $(P_{\lambda, \mu})$ . Proposition 6.1 and Lemma 6.1 appear in the paper by Jebelean and Moroşanu [150], while Theorem 6.2 is due to Lisei, Moroşanu, and Varga [196]. The results contained in Section 6.3 are due to Breckner and Varga [46].

## Competition Phenomena in Dirichlet Problems

The purpose of models is not to fit the data but to sharpen the questions.



---

Samuel Karlin (1924–2007)

Having infinitely many solutions for a given equation, after a ‘small’ perturbation of it, one might be expected to find still many solutions for the perturbed equation; moreover, once the perturbation tends to zero, the number of solutions for the perturbed equation should tend to infinity. Such phenomenon is well-known in the case of the equation  $\sin x = c$  with  $c \in (-1, 1)$  fixed, and its perturbation  $\sin x = c + \varepsilon x$ ,  $x \in \mathbb{R}$ ; the perturbed equation has more and more solutions as  $|\varepsilon|$  decreases to 0. For instance, the equation  $\sin x = 1/2$  has infinitely many solutions, see Fig. 7.1. However, the perturbed equation  $\sin x = 1/2 + 0.2x$  has a finite number of solutions, which is less than the number of solutions of the equation  $\sin x = 1/2 - 0.1x$ , cf. Fig. 7.2 and Fig. 7.3.

Natural phenomena related to perturbation have been first exploited in an abstract framework by Krasnosel’skii [163]. More precisely, by using topological methods, Krasnosel’skii asserts the existence of more and more critical points of an even functional of class  $C^1$  perturbed by a non-even term tending to zero, the critical points of the perturbed functional being the solutions for the studied equation.

The purpose of this chapter is to study the number and behaviour of solutions to a Dirichlet problem which involves an *oscillatory nonlinearity* and a *pure power term*.


Figure 7.1 Equation  $\sin x = 1/2$  has infinitely many solutions.Figure 7.2 Solutions of the equation  $\sin x = 1/2 + 0.2x$ .

## 7.1 Introduction

We consider the multiplicity of solutions to the problem

$$\begin{cases} -\Delta u = \lambda a(x)u^p + f(u) & \text{in } \Omega, \\ u \geq 0, u \not\equiv 0 & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (\text{P}_\lambda)$$

where  $\Omega$  is a smooth bounded domain in  $\mathbb{R}^N$  ( $N \geq 3$ ),  $f : [0, \infty) \rightarrow \mathbb{R}$  oscillates near the origin or at infinity, and  $p > 0$ ,  $\lambda \in \mathbb{R}$ . Denoting formally  $0^+$  or  $+\infty$  by the common symbol  $L$  (standing for a limit point), we assume on the continuous nonlinearity  $f$  that

Figure 7.3 Solutions of the equation  $\sin x = 1/2 - 0.1x$ .

$$(f_1^L). \quad -\infty < \liminf_{s \rightarrow L} \frac{F(s)}{s^2} \leq \limsup_{s \rightarrow L} \frac{F(s)}{s^2} = +\infty;$$

$$(f_2^L). \quad l_L := \liminf_{s \rightarrow L} \frac{f(s)}{s} < 0,$$

where  $F(s) = \int_0^s f(t)dt$ . One can easily observe that  $f$  has an *oscillatory behaviour at  $L$* .

While oscillatory right hand sides usually produce infinitely many distinct solutions, the additional term involving  $u^p$  may alter the situation radically. Our purpose is to fully describe this phenomenon, showing that the number of distinct nontrivial solutions to problem  $(P_\lambda)$  is strongly influenced by  $u^p$  and depends on  $\lambda$  whenever one of the following two cases holds:

- $p \leq 1$  and  $f$  oscillates near the origin;
- $p \geq 1$  and  $f$  oscillates at infinity ( $p$  may be critical or even supercritical). The coefficient  $a \in L^\infty(\Omega)$  is allowed to change its sign, while its size is relevant only for the threshold value  $p = 1$  when the behaviour of  $f(s)/s$  also plays a crucial role in both cases.

## 7.2 Effects of the competition

In this section we present the main results of this chapter, establishing as well some interesting connections between the behaviour of certain algebraic equations and PDEs.

We notice that our problem  $(P_\lambda)$  can be compared with elliptic problems involving the so-called *concave-convex* nonlinearities, see Ambrosetti-




Figure 7.4 Let  $0 \leq p < 1 < q$ . For every  $\lambda_1 < \lambda^* < \lambda_2$ , the algebraic equation  $(E_{p,q}^{\lambda_1})$  has two solutions, while  $(E_{p,q}^{\lambda_2})$  has no solution.

Brézis-Cerami [6], de Figueiredo-Gossez-Ubilla [119], [118]. In such a case, the sublinear term  $u^p$  and the superlinear term  $f(u) = u^q$  compete with each other, where  $0 \leq p < 1 < q \leq (N+2)/(N-2) = 2^* - 1$ . As a consequence of this competition, problem  $(P_\lambda)$  has at least two positive solutions for small  $\lambda > 0$  and no positive solution for large  $\lambda$ . Since  $u \mapsto -\Delta u$  is a linear map, the above statement is well reflected by the algebraic equation

$$s = \lambda s^p + s^q, \quad s > 0. \quad (E_{p,q}^\lambda)$$

Indeed, there exists a  $\lambda^* > 0$  such that for every  $\lambda \in (0, \lambda^*)$  equation  $(E_{p,q}^\lambda)$  has two solutions,  $(E_{p,q}^{\lambda^*})$  has one solution, and for  $\lambda > \lambda^*$  equation  $(E_{p,q}^\lambda)$  has no solution, see Figure 7.4.

Equations involving oscillatory terms usually give infinitely many distinct solutions, see Kristály [170], Kristály-Moroşanu-Tersian [176], Omari-Zanolin [232], Saint Raymond [269]. However, surprising facts may occur even in simple cases; indeed, if  $p = 1$  and we consider the oscillatory function  $f(s) = f_\mu(s) = \mu \sin s$  ( $\mu \in \mathbb{R}$ ), problem  $(P_\lambda)$  has only the trivial solution whenever  $(|\lambda| \cdot \|a\|_{L^\infty} + |\mu|) \lambda_1(\Omega) < 1$ , where  $\lambda_1(\Omega)$  denotes the principal eigenvalue of  $-\Delta$  on  $H_0^1(\Omega)$ , and  $\|\cdot\|_{L^\infty}$  is the  $L^\infty(\Omega)$ -norm. In order to obtain infinitely many distinct solutions for  $(P_0)$  we have to

consider a class of functions having a suitable oscillatory behaviour; this class consists from functions which fulfill  $(f_1^L)$  and  $(f_2^L)$ ,  $L \in \{0^+, +\infty\}$ .

In the sequel, we state our main results, treating separately the two cases, that is, when  $f$  oscillates near the *origin*, and at *infinity*, respectively. The coefficient  $a \in L^\infty(\Omega)$  is allowed to be *indefinite* (namely, it may change its sign), suggested by several recent works, including Alama-Tarantello [4], [3], Berestycki-Cappuzzo Dolcetta-Nirenberg [36], de Figueiredo-Gossez-Ubilla [119], [118], Servadei [274].

*A. Oscillation near the origin.* Let  $f \in C([0, \infty), \mathbb{R})$  and  $F(s) = \int_0^s f(t)dt$ ,  $s \geq 0$ . We assume

$$(f_1^0). \quad -\infty < \liminf_{s \rightarrow 0^+} \frac{F(s)}{s^2}; \quad \limsup_{s \rightarrow 0^+} \frac{F(s)}{s^2} = +\infty;$$

$$(f_2^0). \quad l_0 := \liminf_{s \rightarrow 0^+} \frac{f(s)}{s} < 0.$$

**Remark 7.1** Hypotheses  $(f_1^0)$  and  $(f_2^0)$  imply an oscillatory behaviour of  $f$  near the origin. Let  $\alpha, \beta, \gamma \in \mathbb{R}$  such that  $0 < \alpha < 1 < \alpha + \beta$ , and  $\gamma \in (0, 1)$ . The function  $f_0 : [0, \infty) \rightarrow \mathbb{R}$  defined by  $f_0(0) = 0$  and  $f_0(s) = s^\alpha(\gamma + \sin s^{-\beta})$ ,  $s > 0$ , verifies  $(f_1^0)$  and  $(f_2^0)$ , respectively.

**Theorem 7.1** (Case  $p \geq 1$ ) Assume  $a \in L^\infty(\Omega)$  and let  $f \in C([0, \infty), \mathbb{R})$  satisfy  $(f_1^0)$  and  $(f_2^0)$ . If

- (i) either  $p = 1$  and  $\lambda a(x) < \lambda_0$  a.e.  $x \in \Omega$  for some  $0 < \lambda_0 < -l_0$ ,
- (ii) or  $p > 1$  and  $\lambda \in \mathbb{R}$  is arbitrary,

then there exists a sequence  $\{u_i^0\}_i \subset H_0^1(\Omega)$  of distinct weak solutions of problem  $(P_\lambda)$  such that

$$\lim_{i \rightarrow \infty} \|u_i^0\|_{H_0^1} = \lim_{i \rightarrow \infty} \|u_i^0\|_{L^\infty} = 0. \quad (7.1)$$

**Remark 7.2** (i) If  $l_0 = -\infty$ , then (i) holds for every  $\lambda \in \mathbb{R}$ . For instance, this may happen for  $f_0$  from Remark 7.1.

(ii) Notice that  $p > 1$  may be *critical* or even *supercritical* in Theorem 7.1(ii). Having a suitable nonlinearity oscillating near the origin, Theorem 7.1 roughly says that the term defined by  $s \mapsto s^p$  ( $s \geq 0$ ) does not affect the number of distinct solutions of  $(P_\lambda)$  whenever  $p > 1$ ; this is also the case for certain values of  $\lambda \in \mathbb{R}$  when  $p = 1$ . A similar relation may be stated as before for both the equation  $(E_{p,q}^\lambda)$  and the elliptic problem involving concave-convex nonlinearities. Namely, the thesis of Theorem 7.1 is nicely illustrated by the equation

$$s = \lambda s^p + f_0(s), \quad s \geq 0, \quad (E_0)$$

where  $f_0$  is the function appearing in Remark 7.1. Since  $l_0 = -\infty$ , for




Figure 7.5 For any  $\lambda \in \mathbb{R}$  and  $p = p_1 > 1$ ,  $(E_0)$  has infinitely many distinct solutions, see (a). However, for many  $\lambda \in \mathbb{R}$  and  $p = p_2 < 1$ ,  $(E_0)$  has only finitely many distinct solutions, see (b).

every  $\lambda \in \mathbb{R}$  and  $p \geq 1$ , equation  $(E_0)$  has *infinitely many* distinct positive solutions.

On the other hand, this phenomenon dramatically changes when  $p < 1$ . In this case, the term  $s \mapsto s^p$  ( $s \geq 0$ ) may compete with the function  $f_0$  near the origin such that the number of distinct solutions of  $(E_0)$  becomes *finite* for many values of  $\lambda$ ; this fact happens when  $0 < p < \alpha$  ( $\alpha$  is the number defined in Remark 7.1). However, the number of distinct solutions to  $(E_0)$  becomes greater and greater if  $|\lambda|$  gets smaller and smaller as a simple (graphical) argument shows. See Figure 7.5.

In the language of our Dirichlet problem  $(P_\lambda)$ , the latter statement is perfectly described by the following result.

**Theorem 7.2** (Case  $0 < p < 1$ ) Assume  $a \in L^\infty(\Omega)$ . Let  $f \in C([0, \infty), \mathbb{R})$  satisfy  $(f_1^0)$  and  $(f_2^0)$ , and  $0 < p < 1$ . Then, for every  $k \in \mathbb{N}$ , there exists  $\lambda_k^0 > 0$  such that  $(P_\lambda)$  has at least  $k$  distinct weak solutions  $\{u_{1,\lambda}^0, \dots, u_{k,\lambda}^0\} \subset H_0^1(\Omega)$  whenever  $\lambda \in [-\lambda_k^0, \lambda_k^0]$ . Moreover,

$$\|u_{i,\lambda}^0\|_{H_0^1} < i^{-1} \quad \text{and} \quad \|u_{i,\lambda}^0\|_{L^\infty} < i^{-1} \quad \text{for any } i = \overline{1, k}; \quad \lambda \in [-\lambda_k^0, \lambda_k^0]. \quad (7.2)$$

*B. Oscillation at infinity.* Let  $f \in C([0, \infty), \mathbb{R})$ . We assume

$$(f_1^\infty). \quad -\infty < \liminf_{s \rightarrow \infty} \frac{f(s)}{s^2}; \quad \limsup_{s \rightarrow \infty} \frac{f(s)}{s^2} = +\infty;$$

$$(f_2^\infty). \quad l_\infty := \liminf_{s \rightarrow \infty} \frac{f(s)}{s} < 0.$$

**Remark 7.3** Hypotheses  $(f_1^\infty)$  and  $(f_2^\infty)$  imply an oscillatory behaviour of  $f$  at infinity. Let  $\alpha, \beta, \gamma \in \mathbb{R}$  such that  $1 < \alpha$ ,  $|\alpha - \beta| < 1$ , and  $\gamma \in (0, 1)$ . Then, the function  $f_\infty : [0, \infty) \rightarrow \mathbb{R}$  defined by  $f_\infty(s) = s^\alpha(\gamma + \sin s^\beta)$  verifies the hypotheses  $(f_1^\infty)$  and  $(f_2^\infty)$ , respectively.

The counterpart of Theorem 7.1 can be stated as follows.

**Theorem 7.3** (Case  $p \leq 1$ ) Assume  $a \in L^\infty(\Omega)$ . Let  $f \in C([0, \infty), \mathbb{R})$  satisfy  $(f_1^\infty)$  and  $(f_2^\infty)$  with  $f(0) = 0$ . If

- (i) either  $p = 1$  and  $\lambda a(x) < \lambda_\infty$  a.e.  $x \in \Omega$  for some  $0 < \lambda_\infty < -l_\infty$ ,
- (ii) or  $p < 1$  and  $\lambda \in \mathbb{R}$  is arbitrary,

then there exists a sequence  $\{u_i^\infty\}_i \subset H_0^1(\Omega)$  of distinct, weak solutions of problem  $(P_\lambda)$  such that

$$\lim_{i \rightarrow \infty} \|u_i^\infty\|_{L^\infty} = \infty. \quad (7.3)$$

**Remark 7.4** If

$$\sup_{s \in [0, \infty)} \frac{|f(s)|}{1 + s^{2^* - 1}} < \infty \quad (7.4)$$

then we also have  $\lim_{i \rightarrow \infty} \|u_i^\infty\|_{H_0^1} = \infty$  in Theorem 7.3. For details, see Section 7.5.

**Remark 7.5** A similar observation can be made as in Remark 7.2. Indeed, when  $f$  oscillates at infinity, Theorem 7.3 shows that the term defined by  $s \mapsto s^p$  ( $s \geq 0$ ) does not affect the number of distinct solutions of  $(P_\lambda)$  whenever  $p < 1$ . This is also the case for certain values of  $\lambda \in \mathbb{R}$  when  $p = 1$ . A similar phenomenon occurs in the equation

$$s = \lambda s^p + f_\infty(s), \quad s \geq 0, \quad (E_\infty)$$

where  $f_\infty$  is the function defined in Remark 7.3. Since  $l_\infty = -\infty$ , for every  $\lambda \in \mathbb{R}$  and  $p \leq 1$ , equation  $(E_\infty)$  has *infinitely many* distinct positive solutions.

On the other hand, when  $p > 1$ , the term  $s \mapsto s^p$  ( $s \geq 0$ ) may dominate the function  $f_\infty$  at infinity. In particular, when  $\alpha < p$ , the number of distinct solutions of  $(E_\infty)$  may become *finite* for many values of  $\lambda$  (here,  $\alpha$  is the number defined in Remark 7.3). The positive finding is that the number of distinct solutions for  $(E_\infty)$  increases whenever  $|\lambda|$  decreases to zero. See Figure 7.6.

In view of this observation, we obtain a natural counterpart of Theorem 7.2.

**Theorem 7.4** (Case  $p > 1$ ) Assume  $a \in L^\infty(\Omega)$ . Let  $f \in C([0, \infty), \mathbb{R})$




Figure 7.6 For any  $\lambda \in \mathbb{R}$  and  $p = p_1 < 1$ ,  $(E_\infty)$  has infinitely many distinct solutions, see (a). However, for many  $\lambda \in \mathbb{R}$  and  $p = p_2 > 1$ ,  $(E_\infty)$  has only finitely many (or, even none) distinct solutions, see (b).

satisfy  $(f_1^\infty)$  and  $(f_2^\infty)$  with  $f(0) = 0$ , and  $p > 1$ . Then, for every  $k \in \mathbb{N}$ , there exists  $\lambda_k^\infty > 0$  such that  $(P_\lambda)$  has at least  $k$  distinct weak solutions  $\{u_{1,\lambda}^\infty, \dots, u_{k,\lambda}^\infty\} \subset H_0^1(\Omega)$  whenever  $\lambda \in [-\lambda_k^\infty, \lambda_k^\infty]$ . Moreover,

$$\|u_{i,\lambda}^\infty\|_{L^\infty} > i - 1 \quad \text{for any } i = \overline{1, k}; \quad \lambda \in [-\lambda_k^\infty, \lambda_k^\infty]. \quad (7.5)$$

**Remark 7.6** If  $f$  verifies (7.4) and  $p \leq 2^* - 1$  in Theorem 7.4, then

$$\|u_{i,\lambda}^\infty\|_{H_0^1} > i - 1 \quad \text{for any } i = \overline{1, k}; \quad \lambda \in [-\lambda_k^\infty, \lambda_k^\infty].$$

We conclude this section by stating a result for a model problem which involves concave-convex nonlinearities and an oscillatory term. We consider the problem

$$\begin{cases} -\Delta u = \lambda u^p + \mu u^q + f(u), & u \geq 0 & \text{on } \Omega, \\ u = 0 & & \text{on } \partial\Omega, \end{cases} \quad (P_{\lambda,\mu})$$

where  $0 < p < 1 < q$ , and  $\lambda, \mu \in \mathbb{R}$ . The following result proves that the number of solutions  $(P_{\lambda,\mu})$  is influenced

- (i). by the sublinear term when  $f$  oscillates near the origin (with no effect of the superlinear term); and alternatively,
- (ii). by the superlinear term when  $f$  oscillates at infinity (with no effect of the sublinear term).

More precisely, applying Theorems 7.2 and 7.4, we have

**Theorem 7.5** *Let  $f \in C([0, \infty), \mathbb{R})$  and  $0 < p < 1 < q$ .*

- (i). *If  $(f_1^0)$  and  $(f_2^0)$  hold, then for every  $k \in \mathbb{N}$  and  $\mu \in \mathbb{R}$ , there exists  $\lambda_{k,\mu} > 0$  such that  $(P_{\lambda,\mu})$  has at least  $k$  distinct weak solutions in  $H_0^1(\Omega)$  whenever  $\lambda \in [-\lambda_{k,\mu}, \lambda_{k,\mu}]$ .*
- (ii). *If  $(f_1^\infty)$  and  $(f_2^\infty)$  hold with  $f(0) = 0$ , then for every  $k \in \mathbb{N}$  and  $\lambda \in \mathbb{R}$ , there exists  $\mu_{k,\lambda} > 0$  such that  $(P_{\lambda,\mu})$  has at least  $k$  distinct weak solutions in  $H_0^1(\Omega)$  whenever  $\mu \in [-\mu_{k,\lambda}, \mu_{k,\lambda}]$ .*

### 7.3 A general location property

We consider the problem

$$\begin{cases} -\Delta u + K(x)u = h(x, u), & u \geq 0 & \text{in } \Omega, \\ u = 0 & & \text{on } \partial\Omega, \end{cases} \quad (P_h^K)$$

and assume that

- $(H_K)$ :.  $K \in L^\infty(\Omega)$ ,  $\text{essinf}_\Omega K > 0$ ;
- $(H_h^1)$ :.  $h : \Omega \times [0, \infty) \rightarrow \mathbb{R}$  is a Carathéodory function,  $h(x, 0) = 0$  for a.e.  $x \in \Omega$ , and there is  $M_h > 0$  such that  $|h(x, s)| \leq M_h$  for a.e.  $x \in \Omega$  and all  $s \in \mathbb{R}$ ;
- $(H_h^2)$ :. there are  $0 < \delta < \eta$  such that  $h(x, s) \leq 0$  for a.e.  $x \in \Omega$  and all  $s \in [\delta, \eta]$ .

We extend the function  $h$  by  $h(x, s) = 0$  for a.e.  $x \in \Omega$  and  $s \leq 0$ . We introduce the energy functional  $\mathcal{E} : H_0^1(\Omega) \rightarrow \mathbb{R}$  associated with problem  $(P_h^K)$ , defined by

$$\mathcal{E}(u) = \frac{1}{2} \|u\|_{H_0^1}^2 + \frac{1}{2} \int_\Omega K(x)u^2 dx - \int_\Omega H(x, u(x)) dx, \quad u \in H_0^1(\Omega),$$

where  $H(x, s) = \int_0^s h(x, t) dt$ ,  $s \in \mathbb{R}$ . Due to hypothesis  $(H_h^1)$ , it is easy to see that  $\mathcal{E}$  is well-defined. Moreover, standard arguments show that  $\mathcal{E}$  is of class  $C^1$  on  $H_0^1(\Omega)$ .

Finally, considering the number  $\eta \in \mathbb{R}$  from  $(H_h^2)$ , we introduce the set

$$W^\eta = \{u \in H_0^1(\Omega) : \|u\|_{L^\infty} \leq \eta\}.$$

Since  $h(x, 0) = 0$ , then 0 is clearly a solution of  $(P_h^K)$ . In the sequel, under some general assumptions, we guarantee the existence of a (possible trivial) weak solution of  $(P_h^K)$  which is indispensable in our further investigations (see Sections 7.4 and 7.5).

**Theorem 7.6** *Assume that  $(H_K)$ ,  $(H_h^1)$ ,  $(H_h^2)$  hold. Then*

- (i). *the functional  $\mathcal{E}$  is bounded from below on  $W^\eta$  and its infimum is attained at some  $\tilde{u} \in W^\eta$ ;*
- (ii).  *$\tilde{u}(x) \in [0, \delta]$  for a.e.  $x \in \Omega$ ;*
- (iii).  *$\tilde{u}$  is a weak solution of  $(P_h^K)$ .*

*Proof.* (i) Due to  $(H_h^1)$  and by using Hölder's and Poincaré's inequalities, the functional  $\mathcal{E}$  is bounded from below on the whole space  $H_0^1(\Omega)$ . In addition, one can easily see that  $\mathcal{E}$  is sequentially weak lower semi-continuous and the set  $W^\eta$  is convex and closed in  $H_0^1(\Omega)$ , thus weakly closed. Combining these facts, there is an element  $\tilde{u} \in W^\eta$  which is a minimum point of  $\mathcal{E}$  over  $W^\eta$ .

(ii) Let  $A = \{x \in \Omega : \tilde{u}(x) \notin [0, \delta]\}$  and suppose that  $m(A) > 0$ . Here and in the sequel,  $m(\cdot)$  denotes the Lebesgue measure. Define the function  $\gamma : \mathbb{R} \rightarrow \mathbb{R}$  by  $\gamma(s) = \min(s_+, \delta)$ , where  $s_+ = \max(s, 0)$ . Now, set  $w = \gamma \circ \tilde{u}$ . Since  $\gamma$  is a Lipschitz function and  $\gamma(0) = 0$ , the theorem of Marcus-Mizel [204] shows that  $w \in H_0^1(\Omega)$ . Moreover,  $0 \leq w(x) \leq \delta$  for a.e.  $\Omega$ . Consequently,  $w \in W^\eta$ .

We introduce the sets  $A_1 = \{x \in A : \tilde{u}(x) < 0\}$  and  $A_2 = \{x \in A : \tilde{u}(x) > \delta\}$ . Thus,  $A = A_1 \cup A_2$ , and we have that  $w(x) = \tilde{u}(x)$  for all  $x \in \Omega \setminus A$ ,  $w(x) = 0$  for all  $x \in A_1$ , and  $w(x) = \delta$  for all  $x \in A_2$ . Moreover, we have

$$\begin{aligned} \mathcal{E}(w) - \mathcal{E}(\tilde{u}) &= \\ &= \frac{1}{2} \left[ \|w\|_{H_0^1}^2 - \|\tilde{u}\|_{H_0^1}^2 \right] + \frac{1}{2} \int_{\Omega} K(x) [w^2 - \tilde{u}^2] - \int_{\Omega} [H(x, w) - H(x, \tilde{u})] \\ &= -\frac{1}{2} \int_A |\nabla \tilde{u}|^2 + \frac{1}{2} \int_A K(x) [w^2 - \tilde{u}^2] - \int_A [H(x, w) - H(x, \tilde{u})]. \end{aligned}$$

Since  $\text{essinf}_{\Omega} K > 0$ , one has

$$\int_A K(x) [w^2 - \tilde{u}^2] = - \int_{A_1} K(x) \tilde{u}^2 + \int_{A_2} K(x) [\delta^2 - \tilde{u}^2] \leq 0.$$

Due to the fact that  $h(x, s) = 0$  for all  $s \leq 0$ , one has

$$\int_{A_1} [H(x, w) - H(x, \tilde{u})] = 0.$$

By the mean value theorem, for a.e.  $x \in A_2$ , there exists  $\theta(x) \in [\delta, \tilde{u}(x)] \subseteq [\delta, \eta]$  such that

$$H(x, w(x)) - H(x, \tilde{u}(x)) = H(x, \delta) - H(x, \tilde{u}(x)) = h(x, \theta(x))(\delta - \tilde{u}(x)).$$

Thus, on account of  $(H_h^2)$ , one has

$$\int_{A_2} [H(x, w) - H(x, \tilde{u})] \geq 0.$$

Consequently, every term of the expression  $\mathcal{E}(w) - \mathcal{E}(\tilde{u})$  is non-positive. On the other hand, since  $w \in W^\eta$ , then  $\mathcal{E}(w) \geq \mathcal{E}(\tilde{u}) = \inf_{W^\eta} \mathcal{E}$ . So, every term in  $\mathcal{E}(w) - \mathcal{E}(\tilde{u})$  should be zero. In particular,

$$\int_{A_1} K(x) \tilde{u}^2 = \int_{A_2} K(x) [\tilde{u}^2 - \delta^2] = 0.$$

Due to  $(H_K)$ , we necessarily have  $m(A) = 0$ , contradicting our assumption.

(iii) Let us fix  $v \in C_0^\infty(\Omega)$  and let  $\varepsilon_0 = (\eta - \delta)/(\|v\|_{C_0} + 1) > 0$ . Define the function  $E : [-\varepsilon_0, \varepsilon_0] \rightarrow \mathbb{R}$  by  $E(\varepsilon) = \mathcal{E}(\tilde{u} + \varepsilon v)$  with  $\varepsilon \in [-\varepsilon_0, \varepsilon_0]$ . Due to (ii), for every  $\varepsilon \in [-\varepsilon_0, \varepsilon_0]$ , the element  $\tilde{u} + \varepsilon v$  belongs to the set  $W^\eta$ . Consequently, due to (i), one has  $E(\varepsilon) \geq E(0)$  for every  $\varepsilon \in [-\varepsilon_0, \varepsilon_0]$ . Since  $E$  is differentiable at 0 and  $E'(0) = 0$  it follows that  $\langle \mathcal{E}'(\tilde{u}), v \rangle_{H_0^1} = 0$ . Since  $v \in C_0^\infty(\Omega)$  is arbitrary, and the set  $C_0^\infty(\Omega)$  is dense in  $H_0^1(\Omega)$ , we obtain that  $\tilde{u}$  is a weak solution of  $(P_h^K)$ .  $\square$

We conclude this section by constructing a special function which will be useful in the proof of our theorems. In the sequel, let  $B(x_0, r) \subset \Omega$  be the  $N$ -dimensional ball with radius  $r > 0$  and center  $x_0 \in \Omega$ . For  $s > 0$ , define

$$z_s(x) = \begin{cases} 0, & \text{if } x \in \Omega \setminus B(x_0, r); \\ s, & \text{if } x \in B(x_0, r/2); \\ \frac{2s}{r}(r - |x - x_0|), & \text{if } x \in B(x_0, r) \setminus B(x_0, r/2). \end{cases} \quad (7.6)$$

It is clear that  $z_s \in H_0^1(\Omega)$ . Moreover, we have  $\|z_s\|_{L^\infty} = s$  and

$$\|z_s\|_{H_0^1}^2 = \int_{\Omega} |\nabla z_s|^2 = 4r^{N-2}(1 - 2^{-N})\omega_N s^2 \equiv C(r, N)s^2 > 0, \quad (7.7)$$

where  $\omega_N$  is the volume of  $B(0, 1) \subset \mathbb{R}^N$ .

*Notation.* For every  $\eta > 0$ , we define the truncation function  $\tau_\eta : [0, \infty) \rightarrow \mathbb{R}$  by  $\tau_\eta(s) = \min(\eta, s)$ ,  $s \geq 0$ .

#### 7.4 Nonlinearities with oscillation near the origin

Since parts (i) and (ii) of Theorem 7.1 will be treated simultaneously, we consider again the problem from the previous Section

$$\begin{cases} -\Delta u + K(x)u = h(x, u), & u \geq 0 & \text{in } \Omega, \\ u = 0 & & \text{on } \partial\Omega, \end{cases} \quad (\mathbf{P}_h^K)$$

where the potential  $K : \Omega \rightarrow \mathbb{R}$  fulfills  $(\mathbf{H}_K)$ . The function  $h : \Omega \times [0, \infty) \rightarrow \mathbb{R}$  is Carathéodory, and we assume

$(\mathbf{H}_0^0)$ :.  $h(x, 0) = 0$  for a.e.  $x \in \Omega$ , and there exists  $s_0 > 0$  such that

$$\sup_{s \in [0, s_0]} |h(\cdot, s)| \in L^\infty(\Omega);$$

$(\mathbf{H}_1^0)$ :.  $-\infty < \liminf_{s \rightarrow 0^+} \frac{H(x, s)}{s^2}$  and  $\limsup_{s \rightarrow 0^+} \frac{H(x, s)}{s^2} = +\infty$  uniformly for a.e.  $x \in \Omega$ ; here,  $H(x, s) = \int_0^s h(x, t)dt$ ;

$(\mathbf{H}_2^0)$ :. there are two sequences  $\{\delta_i\}, \{\eta_i\}$  with  $0 < \eta_{i+1} < \delta_i < \eta_i$ ,  $\lim_{i \rightarrow \infty} \eta_i = 0$ , and  $h(x, s) \leq 0$  for a.e.  $x \in \Omega$  and for every  $s \in [\delta_i, \eta_i]$ ,  $i \in \mathbb{N}$ .

**Theorem 7.7** *Assume  $(\mathbf{H}_K)$ ,  $(\mathbf{H}_0^0)$ ,  $(\mathbf{H}_1^0)$  and  $(\mathbf{H}_2^0)$  hold. Then there exists a sequence  $\{u_i^0\}_i \subset H_0^1(\Omega)$  of distinct weak solutions of problem  $(\mathbf{P}_h^K)$  such that*

$$\lim_{i \rightarrow \infty} \|u_i^0\|_{H_0^1} = \lim_{i \rightarrow \infty} \|u_i^0\|_{L^\infty} = 0. \quad (7.8)$$

*Proof* Without any loss of generality, we may assume that  $\{\delta_i\}_i, \{\eta_i\}_i \subset (0, s_0)$ , where  $s_0 > 0$  comes from  $(\mathbf{H}_0^0)$ . For every  $i \in \mathbb{N}$ , define the truncation function  $h_i : \Omega \times [0, \infty) \rightarrow \mathbb{R}$  by

$$h_i(x, s) = h(x, \tau_{\eta_i}(s)) \quad (7.9)$$

and let  $\mathcal{E}_i : H_0^1(\Omega) \rightarrow \mathbb{R}$  be the energy functional associated with the problem  $(\mathbf{P}_{h_i}^K)$ . Let  $H_i(x, s) = \int_0^s h_i(x, t)dt$ .

Due to hypotheses  $(\mathbf{H}_0^0)$  and  $(\mathbf{H}_2^0)$ , the function  $h_i$  verifies the assumptions of Theorem 7.6 for every  $i \in \mathbb{N}$  with  $[\delta_i, \eta_i]$ . Consequently, for every  $i \in \mathbb{N}$ , there exists  $u_i^0 \in W^{\eta_i}$  such that

$$u_i^0 \text{ is the minimum point of the functional } \mathcal{E}_i \text{ on } W^{\eta_i}, \quad (7.10)$$

$$u_i^0(x) \in [0, \delta_i] \text{ for a.e. } x \in \Omega, \tag{7.11}$$

$$u_i^0 \text{ is a weak solution of } (P_{h_i}^K). \tag{7.12}$$

Thanks to (7.9), (7.11) and (7.12),  $u_i^0$  is a weak solution not only for  $(P_{h_i}^K)$  but also for the problem  $(P_h^K)$ .

Now, we prove that there are infinitely many distinct elements in the sequence  $\{u_i^0\}_i$ . To see this, we first prove that

$$\mathcal{E}_i(u_i^0) < 0 \text{ for all } i \in \mathbb{N}; \tag{7.13}$$

$$\lim_{i \rightarrow \infty} \mathcal{E}_i(u_i^0) = 0. \tag{7.14}$$

The left part of  $(H_1^0)$  implies the existence of some  $l_0^h > 0$  and  $\zeta \in (0, \eta_1)$  such that

$$\text{essinf}_{x \in \Omega} H(x, s) \geq -l_0^h s^2 \text{ for all } s \in (0, \zeta). \tag{7.15}$$

Let  $L_0^h > 0$  be large enough so that

$$\frac{1}{2}C(r, N) + \left(\frac{1}{2}\|K\|_{L^\infty} + l_0^h\right) m(\Omega) < L_0^h (r/2)^N \omega_N, \tag{7.16}$$

where  $r > 0$  and  $C(r, N) > 0$  come from (7.7). Taking into account the right part of  $(H_1^0)$ , there is a sequence  $\{\tilde{s}_i\}_i \subset (0, \zeta)$  such that  $\tilde{s}_i \leq \delta_i$  and

$$\text{essinf}_{x \in \Omega} H(x, \tilde{s}_i) > L_0^h \tilde{s}_i^2 \text{ for all } i \in \mathbb{N}. \tag{7.17}$$

Let  $i \in \mathbb{N}$  be a fixed number and let  $z_{\tilde{s}_i} \in H_0^1(\Omega)$  be the function from (7.6) corresponding to the value  $\tilde{s}_i > 0$ . Then  $z_{\tilde{s}_i} \in W^{\eta_i}$ , and on account of (7.7), (7.17) and (7.15), one has

$$\begin{aligned} \mathcal{E}_i(z_{\tilde{s}_i}) &= \frac{1}{2}\|z_{\tilde{s}_i}\|_{H_0^1}^2 + \frac{1}{2} \int_{\Omega} K(x) z_{\tilde{s}_i}^2 - \int_{\Omega} H_i(x, z_{\tilde{s}_i}(x)) dx \\ &= \frac{1}{2}C(r, N) \tilde{s}_i^2 + \frac{1}{2} \int_{\Omega} K(x) z_{\tilde{s}_i}^2 \\ &\quad - \int_{B(x_0, r/2)} H(x, \tilde{s}_i) dx - \int_{B(x_0, r) \setminus B(x_0, r/2)} H(x, z_{\tilde{s}_i}(x)) dx \\ &\leq \left[ \frac{1}{2}C(r, N) + \frac{1}{2}\|K\|_{L^\infty} m(\Omega) - L_0^h (r/2)^N \omega_N + l_0^h m(\Omega) \right] \tilde{s}_i^2. \end{aligned}$$

Consequently, using (7.10) and (7.16), we obtain that

$$\mathcal{E}_i(u_i^0) = \min_{W^{\eta_i}} \mathcal{E}_i \leq \mathcal{E}_i(z_{\tilde{s}_i}) < 0 \tag{7.18}$$

which proves in particular (7.13). Now, let us prove (7.14). For every  $i \in \mathbb{N}$ , by using the mean value theorem, (7.9),  $(H_0^0)$  and (7.11), we have

$$\mathcal{E}_i(u_i^0) \geq - \int_{\Omega} H_i(x, u_i^0(x)) dx \geq - \left\| \sup_{s \in [0, s_0]} |h(\cdot, s)| \right\|_{L^\infty} m(\Omega) \delta_i.$$

Due to  $\lim_{i \rightarrow \infty} \delta_i = 0$ , the above inequality and (7.18) leads to (7.14).

On account of (7.9) and (7.11), we observe that

$$\mathcal{E}_i(u_i^0) = \mathcal{E}_1(u_i^0) \quad \text{for all } i \in \mathbb{N}.$$

Combining this relation with (7.13) and (7.14), we see that the sequence  $\{u_i^0\}_i$  contains infinitely many distinct elements.

It remains to prove relation (7.8). The former limit easily follows by (7.11), that is,  $\|u_i^0\|_{L^\infty} \leq \delta_i$  for all  $i \in \mathbb{N}$ , combined with  $\lim_{i \rightarrow \infty} \delta_i = 0$ . For the latter limit, we use (7.18),  $(H_0^0)$ , (7.9) and (7.11), obtaining for all  $i \in \mathbb{N}$  that

$$\begin{aligned} \frac{1}{2} \|u_i^0\|_{H_0^1}^2 &\leq \frac{1}{2} \|u_i^0\|_{H_0^1}^2 + \frac{1}{2} \int_{\Omega} K(x) (u_i^0)^2 \\ &< \int_{\Omega} H_i(x, u_i^0(x)) \leq \left\| \sup_{s \in [0, s_0]} |h(\cdot, s)| \right\|_{L^\infty} m(\Omega) \delta_i \end{aligned}$$

which concludes the proof of Theorem 7.7.  $\square$

*Proof of Theorem 7.1. (i) Case  $p = 1$ .* Let  $\lambda \in \mathbb{R}$  as in the hypothesis, namely  $\lambda a(x) < \lambda_0$  a.e.  $x \in \Omega$  for some  $0 < \lambda_0 < -l_0$ . Let us choose  $\tilde{\lambda}_0 \in (\lambda_0, -l_0)$  and

$$K(x) = \tilde{\lambda}_0 - \lambda a(x) \quad \text{and} \quad h(x, s) = \tilde{\lambda}_0 s + f(s) \quad \text{for all } (x, s) \in \Omega \times [0, \infty). \quad (7.19)$$

Note that  $\text{essinf}_{\Omega} K \geq \tilde{\lambda}_0 - \lambda_0 > 0$ , so  $(H_K)$  is satisfied. Due to  $(f_1^0)$  and  $(f_2^0)$ , we have  $f(0) = 0$ . Thus,  $(H_0^0)$  clearly holds. Moreover, since  $H(x, s)/s^2 = \tilde{\lambda}_0/2 + F(s)/s^2$ ,  $s > 0$ , hypothesis  $(f_1^0)$  implies  $(H_1^0)$ . Finally, since  $l_0 < -\tilde{\lambda}_0$ , there exists a sequence  $\{s_i\}_i \subset (0, 1)$  converging to 0 such that  $f(s_i)/s_i < -\tilde{\lambda}_0$  for all  $i \in \mathbb{N}$ . Consequently, by using the continuity of  $f$ , we may choose two sequences  $\{\delta_i\}_i, \{\eta_i\}_i \subset (0, 1)$  such that  $0 < \eta_{i+1} < \delta_i < s_i < \eta_i$ ,  $\lim_{i \rightarrow \infty} \eta_i = 0$ , and  $\tilde{\lambda}_0 s + f(s) \leq 0$  for all  $s \in [\delta_i, \eta_i]$  and  $i \in \mathbb{N}$ . Therefore,  $(H_2^0)$  holds too. It remains to apply Theorem 7.7, observing that  $(P_h^K)$  is equivalent to problem  $(P_\lambda)$  via the choice (7.19).

(ii) *Case*  $p > 1$ . Let  $\lambda \in \mathbb{R}$  be arbitrary fixed. Let us also fix a number  $\lambda_0 \in (0, -l_0)$  and choose

$$K(x) = \lambda_0 \quad \text{and} \quad h(x, s) = \lambda a(x)s^p + \lambda_0 s + f(s) \quad \text{for all } (x, s) \in \Omega \times [0, \infty). \quad (7.20)$$

Clearly,  $(H_K)$  is satisfied. Since  $a \in L^\infty(\Omega)$ , a simple argument yields that  $(H_0^0)$  also holds. Moreover, since  $p > 1$  and  $H(x, s)/s^2 = \lambda a(x)s^{p-1}/(p+1) + \lambda_0/2 + F(s)/s^2$ ,  $s > 0$ , hypothesis  $(f_1^0)$  implies  $(H_1^0)$ . Note that for a.e  $x \in \Omega$  and every  $s \in [0, \infty)$ , we have

$$h(x, s) \leq |\lambda| \cdot \|a\|_{L^\infty} s^p + \lambda_0 s + f(s) \equiv \tilde{h}_0(s). \quad (7.21)$$

Due to  $(f_2^0)$ ,

$$\liminf_{s \rightarrow 0^+} \frac{\tilde{h}_0(s)}{s} = \lambda_0 + l_0 < 0.$$

In particular, there exists a sequence  $\{s_i\}_i \subset (0, 1)$  converging to 0 such that  $\tilde{h}_0(s_i) < 0$  for all  $i \in \mathbb{N}$ . Consequently, by using the continuity of  $\tilde{h}_0$ , we can choose two sequences  $\{\delta_i\}_i, \{\eta_i\}_i \subset (0, 1)$  such that  $0 < \eta_{i+1} < \delta_i < s_i < \eta_i$ ,  $\lim_{i \rightarrow \infty} \eta_i = 0$ , and  $\tilde{h}_0(s) \leq 0$  for all  $s \in [\delta_i, \eta_i]$  and  $i \in \mathbb{N}$ . Therefore, by using (7.21), hypothesis  $(H_2^0)$  holds. Now, we can apply Theorem 7.7; problem  $(P_h^K)$  is equivalent to problem  $(P_\lambda)$  through the choice (7.20). In both cases (i) and (ii), relation (7.1) is implied by (7.8). This completes the proof of Theorem 7.1.  $\square$

*Proof of Theorem 7.2.* The proof is divided into four steps.

*Step 1.* Let  $\lambda_0 \in (0, -l_0)$ . On account of  $(f_1^0)$ , there exists a sequence  $\{s_i\}_i \subset (0, 1)$  converging to 0, such that  $f(s_i)/s_i < -\lambda_0$ . For every  $\lambda \in \mathbb{R}$  define the functions  $h^\lambda : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$  and  $\tilde{h} : \mathbb{R}^2 \rightarrow \mathbb{R}$  by

$$h^\lambda(x, s) = \lambda a(x)s^p + \lambda_0 s + f(s) \quad \text{for all } (x, s) \in \Omega \times [0, \infty);$$

$$\tilde{h}(\lambda, s) = |\lambda| \cdot \|a\|_{L^\infty} s^p + \lambda_0 s + f(s) \quad \text{for all } s \in [0, \infty).$$

Since  $\tilde{h}(0, s_i) = \lambda_0 s_i + f(s_i) < 0$  and due to the continuity of  $\tilde{h}$ , we can choose three sequences  $\{\delta_i\}_i, \{\eta_i\}_i, \{\lambda_i\}_i \subset (0, 1)$  such that  $0 < \eta_{i+1} < \delta_i < s_i < \eta_i$ ,  $\lim_{i \rightarrow \infty} \eta_i = 0$ , and for every  $i \in \mathbb{N}$ ,

$$\tilde{h}(\lambda, s) \leq 0 \quad \text{for all } \lambda \in [-\lambda_i, \lambda_i] \quad \text{and} \quad s \in [\delta_i, \eta_i]. \quad (7.22)$$

Clearly, we may assume that

$$\delta_i \leq \min\{i^{-1}, 2^{-1}i^{-2}[1 + \|a\|_{L^1} + m(\Omega) \max_{s \in [0, 1]} |f(s)|]^{-1}\}, \quad i \in \mathbb{N}. \quad (7.23)$$

Since  $h^\lambda(x, s) \leq \tilde{h}(\lambda, s)$  for a.e.  $x \in \Omega$  and all  $(\lambda, s) \in \mathbb{R} \times [0, \infty)$ , on account of (7.22), for every  $i \in \mathbb{N}$ , we have

$$h^\lambda(x, s) \leq 0 \quad \text{for a.e. } x \in \Omega \text{ and all } \lambda \in [-\lambda_i, \lambda_i], \quad s \in [\delta_i, \eta_i]. \quad (7.24)$$

For every  $i \in \mathbb{N}$  and  $\lambda \in [-\lambda_i, \lambda_i]$ , let  $h_i^\lambda : \Omega \times [0, \infty) \rightarrow \mathbb{R}$  be defined by

$$h_i^\lambda(x, s) = h^\lambda(x, \tau_{\eta_i}(s)) \quad (7.25)$$

and  $K(x) = \lambda_0$ . Let also  $\mathcal{E}_{i,\lambda} : H_0^1(\Omega) \rightarrow \mathbb{R}$  be the energy functional associated with the problem  $(P_{h_i^\lambda}^K)$ , that is,

$$\mathcal{E}_{i,\lambda}(u) = \frac{1}{2} \|u\|_{H_0^1}^2 + \frac{1}{2} \int_{\Omega} K(x) u^2 - \int_{\Omega} \left( \int_0^{u(x)} h_i^\lambda(x, s) ds \right) dx. \quad (7.26)$$

Then, for every  $i \in \mathbb{N}$  and  $\lambda \in [-\lambda_i, \lambda_i]$ , the function  $h_i^\lambda$  verifies the hypotheses of Theorem 7.6; see (7.24) for  $(H_{h_i^\lambda}^2)$ . Therefore, for every  $i \in \mathbb{N}$  and  $\lambda \in [-\lambda_i, \lambda_i]$

$$\text{there exists } u_{i,\lambda}^0 \in W^{\eta_i} \text{ such that } \mathcal{E}_{i,\lambda}(u_{i,\lambda}^0) = \min_{W^{\eta_i}} \mathcal{E}_{i,\lambda}; \quad (7.27)$$

$$u_{i,\lambda}^0(x) \in [0, \delta_i] \text{ for a.e. } x \in \Omega, \quad (7.28)$$

$$u_{i,\lambda}^0 \text{ is a weak solution of } (P_{h_i^\lambda}^K). \quad (7.29)$$

Due to the definition of the functions  $h_i^\lambda$  and  $K$ ,  $u_{i,\lambda}^0$  is a weak solution not only for  $(P_{h_i^\lambda}^K)$ , see (7.25), (7.28) and (7.29), but also for our initial problem  $(P_\lambda)$  once we guarantee that  $u_{i,\lambda}^0 \not\equiv 0$ .

*Step 2.* For  $\lambda = 0$ , the function  $h_i^\lambda = h_i^0$  verifies the hypotheses of Theorem 7.7; more precisely,  $h_i^0$  is precisely the function appearing in (7.9) and  $\mathcal{E}_i := \mathcal{E}_{i,0}$  is the energy functional associated with problem  $(P_{h_i^0}^K)$ . Consequently, besides (7.27)-(7.29), the elements  $u_i^0 := u_{i,0}^0$  also verify

$$\mathcal{E}_i(u_i^0) = \min_{W^{\eta_i}} \mathcal{E}_i \leq \mathcal{E}_i(z_{\bar{s}_i}) < 0 \text{ for all } i \in \mathbb{N}, \quad (7.30)$$

where  $z_{\bar{s}_i} \in W^{\eta_i}$  come from the proof of Theorem 7.7, see (7.18).

*Step 3.* Let  $\{\theta_i\}_i$  be a sequence with negative terms such that  $\lim_{i \rightarrow \infty} \theta_i = 0$ . On account of (7.30), up to a subsequence, we may assume that

$$\theta_i < \mathcal{E}_i(u_i^0) \leq \mathcal{E}_i(z_{\bar{s}_i}) < \theta_{i+1}. \quad (7.31)$$

Let

$$\lambda_i' = \frac{(p+1)(\theta_{i+1} - \mathcal{E}_i(z_{\bar{s}_i}))}{\|a\|_{L^1} + 1} \quad \text{and} \quad \lambda_i'' = \frac{(p+1)(\mathcal{E}_i(u_i^0) - \theta_i)}{\|a\|_{L^1} + 1}, \quad i \in \mathbb{N}.$$

Fix  $k \in \mathbb{N}$ . On account of (7.31),

$$\lambda_k^0 = \min(\lambda_1, \dots, \lambda_k, \lambda'_1, \dots, \lambda'_k, \lambda''_1, \dots, \lambda''_k) > 0.$$

Then, for every  $i \in \{1, \dots, k\}$  and  $\lambda \in [-\lambda_k^0, \lambda_k^0]$  we have

$$\begin{aligned} \mathcal{E}_{i,\lambda}(u_{i,\lambda}^0) &\leq \mathcal{E}_{i,\lambda}(z_{\tilde{s}_i}) && \text{(see (7.27))} \\ &= \frac{1}{2} \|z_{\tilde{s}_i}\|_{H_0^1}^2 - \frac{\lambda}{p+1} \int_{\Omega} a(x) z_{\tilde{s}_i}^{p+1} - \int_{\Omega} F(z_{\tilde{s}_i}(x)) dx \\ &= \mathcal{E}_i(z_{\tilde{s}_i}) - \frac{\lambda}{p+1} \int_{\Omega} a(x) z_{\tilde{s}_i}^{p+1} \\ &< \theta_{i+1}, && \text{(see the choice of } \lambda'_i \text{ and } \tilde{s}_i \leq \delta_i < 1) \end{aligned}$$

and taking into account that  $u_{i,\lambda}^0$  belongs to  $W^{\eta_i}$ , and  $u_i^0$  is the minimum point of  $\mathcal{E}_i$  over the set  $W^{\eta_i}$ , see relation (7.30), we have

$$\begin{aligned} \mathcal{E}_{i,\lambda}(u_{i,\lambda}^0) &= \mathcal{E}_i(u_{i,\lambda}^0) - \frac{\lambda}{p+1} \int_{\Omega} a(x) (u_{i,\lambda}^0)^{p+1} \\ &\geq \mathcal{E}_i(u_i^0) - \frac{\lambda}{p+1} \int_{\Omega} a(x) (u_{i,\lambda}^0)^{p+1} \\ &> \theta_i. && \text{(see the choice of } \lambda''_i \text{ and (7.28))} \end{aligned}$$

In conclusion, for every for every  $i \in \{1, \dots, k\}$  and  $\lambda \in [-\lambda_k^0, \lambda_k^0]$  we have

$$\theta_i < \mathcal{E}_{i,\lambda}(u_{i,\lambda}^0) < \theta_{i+1} < 0,$$

thus

$$\mathcal{E}_{1,\lambda}(u_{1,\lambda}^0) < \dots < \mathcal{E}_{k,\lambda}(u_{k,\lambda}^0) < 0.$$

But,  $u_{i,\lambda}^0 \in W^{\eta_i}$  for every  $i \in \{1, \dots, k\}$ , so  $\mathcal{E}_{i,\lambda}(u_{i,\lambda}^0) = \mathcal{E}_{1,\lambda}(u_{i,\lambda}^0)$ , see relation (7.25). Therefore, from above, we obtain that for every  $\lambda \in [-\lambda_k^0, \lambda_k^0]$ ,

$$\mathcal{E}_{1,\lambda}(u_{1,\lambda}^0) < \dots < \mathcal{E}_{1,\lambda}(u_{k,\lambda}^0) < 0 = \mathcal{E}_{1,\lambda}(0).$$

These inequalities show that the elements  $u_{1,\lambda}^0, \dots, u_{k,\lambda}^0$  are distinct (and non-trivial) whenever  $\lambda \in [-\lambda_k^0, \lambda_k^0]$ .

*Step 4.* It remains to prove conclusion (7.2). The former relation follows directly by (7.28) and (7.23). To check the latter, we observe that for every  $i \in \{1, \dots, k\}$  and  $\lambda \in [-\lambda_k^0, \lambda_k^0]$ ,

$$\mathcal{E}_{1,\lambda}(u_{i,\lambda}^0) = \mathcal{E}_{i,\lambda}(u_{i,\lambda}^0) < \theta_{i+1} < 0.$$

Consequently, for every  $i \in \{1, \dots, k\}$  and  $\lambda \in [-\lambda_k^0, \lambda_k^0]$ , by a mean value theorem we obtain

$$\begin{aligned} \frac{1}{2} \|u_{i,\lambda}^0\|_{H_0^1}^2 &< \frac{\lambda}{p+1} \int_{\Omega} a(x)(u_{i,\lambda}^0)^{p+1} + \int_{\Omega} F(u_{i,\lambda}^0(x)) dx \\ &\leq \left[ \frac{1}{p+1} \|a\|_{L^1} + m(\Omega) \max_{s \in [0,1]} |f(s)| \right] \delta_i \text{ (by (7.28) and } \delta_i, \lambda_k^0 \leq 1) \\ &< 2^{-1} i^{-2}, \end{aligned} \quad (\text{see (7.23)})$$

which concludes the proof of Theorem 7.2.  $\square$

## 7.5 Nonlinearities with oscillation at infinity

In order to prove Theorems 7.3 and 7.4 we follow more or less the technique of the previous Section. However, for completeness, we give all the details. We consider again the problem  $(P_h^K)$ , where the Carathéodory function  $h : \Omega \times [0, \infty) \rightarrow \mathbb{R}$  fulfills

$(H_0^\infty)$ :.  $h(x, 0) = 0$  for a.e.  $x \in \Omega$ , and for every  $s \geq 0$ ,

$$\sup_{t \in [0, s]} |h(\cdot, t)| \in L^\infty(\Omega);$$

$(H_1^\infty)$ :.  $-\infty < \liminf_{s \rightarrow \infty} \frac{H(x, s)}{s^2}$  and  $\limsup_{s \rightarrow \infty} \frac{H(x, s)}{s^2} = +\infty$  uniformly for a.e.  $x \in \Omega$ ; here,  $H(x, s) = \int_0^s h(x, t) dt$ ;

$(H_2^\infty)$ :. there are two sequences  $\{\delta_i\}, \{\eta_i\}$  with  $0 < \delta_i < \eta_i < \delta_{i+1}$ ,  $\lim_{i \rightarrow \infty} \delta_i = +\infty$ , and  $h(x, s) \leq 0$  for a.e.  $x \in \Omega$  and for every  $s \in [\delta_i, \eta_i]$ ,  $i \in \mathbb{N}$ .

**Theorem 7.8** *Assume  $(H_K)$ ,  $(H_0^\infty)$ ,  $(H_1^\infty)$  and  $(H_2^\infty)$  hold. Then there exists a sequence  $\{u_i^\infty\}_i \subset H_0^1(\Omega)$  of distinct weak solutions of problem  $(P_h^K)$  such that*

$$\lim_{i \rightarrow \infty} \|u_i^\infty\|_{L^\infty} = \infty. \quad (7.32)$$

*Proof.* For any  $i \in \mathbb{N}$ , we introduce the truncation function  $h_i : \Omega \times [0, \infty) \rightarrow \mathbb{R}$  by

$$h_i(x, s) = h(x, \tau_{\eta_i}(s)). \quad (7.33)$$

Let  $\mathcal{E}_i : H_0^1(\Omega) \rightarrow \mathbb{R}$  be the energy functional associated with problem  $(P_{h_i}^K)$ . As before, let  $H_i(x, s) = \int_0^s h_i(x, t) dt$ .

On account of hypotheses  $(H_0^\infty)$  and  $(H_2^\infty)$ ,  $h_i$  fulfills the assumptions of Theorem 7.6 for every  $i \in \mathbb{N}$  with  $[\delta_i, \eta_i]$ . Thus, for every  $i \in \mathbb{N}$ , there is an element  $u_i^\infty \in W^{\eta_i}$  such that

$$u_i^\infty \text{ is the minimum point of the functional } \mathcal{E}_i \text{ on } W^{\eta_i}, \quad (7.34)$$

$$u_i^\infty(x) \in [0, \delta_i] \text{ for a.e. } x \in \Omega, \quad (7.35)$$

$$u_i^\infty \text{ is a weak solution of } (P_{h_i}^K). \quad (7.36)$$

Due to (7.36), (7.33) and (7.35),  $u_i^\infty$  is also a weak solution to problem  $(P_h^K)$ .

We prove that there are infinitely many distinct elements in the sequence  $\{u_i^\infty\}_i$ . To this end, it is enough to show that

$$\lim_{i \rightarrow \infty} \mathcal{E}_i(u_i^\infty) = -\infty. \quad (7.37)$$

Indeed, let us assume that in the sequence  $\{u_i^\infty\}_i$  there are only finitely many distinct elements, say  $\{u_1^\infty, \dots, u_{i_0}^\infty\}$  for some  $i_0 \in \mathbb{N}$ . Consequently, due to (7.33), the sequence  $\{\mathcal{E}_i(u_i^\infty)\}_i$  reduces to at most the finite set  $\{\mathcal{E}_{i_0}(u_1^\infty), \dots, \mathcal{E}_{i_0}(u_{i_0}^\infty)\}$ , which contradicts relation (7.37).

Now, we prove (7.37). By  $(H_1^\infty)$ , there exist  $l_\infty^h > 0$  and  $\zeta > 0$  such that

$$\operatorname{ess\,inf}_{x \in \Omega} H(x, s) \geq -l_\infty^h s^2 \text{ for all } s > \zeta. \quad (7.38)$$

Fix  $L_\infty^h > 0$  large enough such that

$$\frac{1}{2}C(r, N) + \left(\frac{1}{2}\|K\|_{L^\infty} + l_\infty^h\right)m(\Omega) < L_\infty^h (r/2)^N \omega_N, \quad (7.39)$$

where  $r > 0$  and  $C(r, N) > 0$  are from (7.7). Due to the right hand side of  $(H_1^\infty)$ , one can fix a sequence  $\{\tilde{s}_i\}_i \subset (0, \infty)$  such that  $\lim_{i \rightarrow \infty} \tilde{s}_i = \infty$ , and

$$\operatorname{ess\,inf}_{x \in \Omega} H(x, \tilde{s}_i) > L_\infty^h \tilde{s}_i^2 \text{ for all } i \in \mathbb{N}. \quad (7.40)$$

Since  $\lim_{i \rightarrow \infty} \delta_i = \infty$ , see  $(H_2^\infty)$ , we can choose a subsequence  $\{\delta_{m_i}\}_i$  of  $\{\delta_i\}_i$  such that  $\tilde{s}_i \leq \delta_{m_i}$  for all  $i \in \mathbb{N}$ . Let  $i \in \mathbb{N}$  be fixed and let  $z_{\tilde{s}_i} \in H_0^1(\Omega)$  be the function from (7.6) corresponding to the value  $\tilde{s}_i > 0$ . Then  $z_{\tilde{s}_i} \in W^{\eta_{m_i}}$ , and on account of (7.7), (7.40) and (7.38), we

have

$$\begin{aligned}
\mathcal{E}_{m_i}(z_{\tilde{s}_i}) &= \frac{1}{2}\|z_{\tilde{s}_i}\|_{H_0^1}^2 + \frac{1}{2}\int_{\Omega} K(x)z_{\tilde{s}_i}^2 - \int_{\Omega} H_{m_i}(x, z_{\tilde{s}_i}(x))dx \\
&= \frac{1}{2}C(r, N)\tilde{s}_i^2 + \frac{1}{2}\int_{\Omega} K(x)z_{\tilde{s}_i}^2 - \int_{B(x_0, r/2)} H(x, \tilde{s}_i)dx \\
&\quad - \int_{(B(x_0, r) \setminus B(x_0, r/2)) \cap \{z_{\tilde{s}_i} > \zeta\}} H(x, z_{\tilde{s}_i}(x))dx \\
&\quad - \int_{(B(x_0, r) \setminus B(x_0, r/2)) \cap \{z_{\tilde{s}_i} \leq \zeta\}} H(x, z_{\tilde{s}_i}(x))dx \\
&\leq \left[ \frac{1}{2}C(r, N) + \frac{1}{2}\|K\|_{L^\infty}m(\Omega) - L_\infty^h(r/2)^N\omega_N + l_\infty^h m(\Omega) \right] \tilde{s}_i^2 \\
&\quad + \|\sup_{s \in [0, \zeta]} |h(\cdot, s)|\|_{L^\infty} m(\Omega)\zeta.
\end{aligned}$$

The above estimate, relation (7.39) and  $\lim_{i \rightarrow \infty} \tilde{s}_i = \infty$  clearly show that

$$\lim_{i \rightarrow \infty} \mathcal{E}_{m_i}(z_{\tilde{s}_i}) = -\infty. \quad (7.41)$$

On the other hand, using (7.34), we have

$$\mathcal{E}_{m_i}(u_{m_i}^\infty) = \min_{W^{\eta_{m_i}}} \mathcal{E}_{m_i} \leq \mathcal{E}_{m_i}(z_{\tilde{s}_i}). \quad (7.42)$$

Therefore, on account of (7.41), we have

$$\lim_{i \rightarrow \infty} \mathcal{E}_{m_i}(u_{m_i}^\infty) = -\infty. \quad (7.43)$$

Note that the sequence  $\{\mathcal{E}_i(u_i^\infty)\}_i$  is non-increasing. Indeed, let  $i, k \in \mathbb{N}$ ,  $i < k$ . Then, due to (7.33), we have

$$\mathcal{E}_i(u_i^\infty) = \min_{W^{\eta_i}} \mathcal{E}_i = \min_{W^{\eta_i}} \mathcal{E}_k \geq \min_{W^{\eta_k}} \mathcal{E}_k = \mathcal{E}_k(u_k^\infty).$$

Combining this fact with (7.43), we obtain (7.37).

Now, we prove (7.32). Arguing by contradiction assume there exists a subsequence  $\{u_{k_i}^\infty\}_i$  of  $\{u_i^\infty\}_i$  such that for all  $i \in \mathbb{N}$ , we have  $\|u_{k_i}^\infty\|_{L^\infty} \leq M$  for some  $M > 0$ . In particular,  $\{u_{k_i}^\infty\}_i \subset W^{\eta_l}$  for some  $l \in \mathbb{N}$ . Thus, for every  $k_i \geq l$ , we have

$$\begin{aligned}
\mathcal{E}_l(u_l^\infty) &= \min_{W^{\eta_l}} \mathcal{E}_l = \min_{W^{\eta_l}} \mathcal{E}_{k_i} \\
&\geq \min_{W^{\eta_{k_i}}} \mathcal{E}_{k_i} = \mathcal{E}_{k_i}(u_{k_i}^\infty) \\
&\geq \min_{W^{\eta_l}} \mathcal{E}_{k_i} && \text{(cf. hypothesis, } u_{k_i}^\infty \in W^{\eta_l}\text{)} \\
&= \mathcal{E}_l(u_l^\infty).
\end{aligned}$$

As a consequence,

$$\mathcal{E}_{k_i}(u_{k_i}^\infty) = \mathcal{E}_l(u_l^\infty) \quad \text{for all } i \in \mathbb{N}. \quad (7.44)$$

Since the sequence  $\{\mathcal{E}_i(u_i^\infty)\}_i$  is non-increasing, on account of (7.44), one can find a number  $i_0 \in \mathbb{N}$  such that  $\mathcal{E}_i(u_i^\infty) = \mathcal{E}_l(u_l^\infty)$  for all  $i \geq i_0$ . This fact contradicts (7.37), which concludes the proof of Theorem 7.8.  $\square$

*Proof of Theorem 7.3. (i) Case  $p = 1$ .* Let us fix  $\lambda \in \mathbb{R}$  as in the hypothesis, that is,  $\lambda a(x) < \lambda_\infty$  a.e.  $x \in \Omega$  for some  $0 < \lambda_\infty < -l_\infty$ . Fix also  $\tilde{\lambda}_\infty \in (\lambda_\infty, -l_\infty)$  and let

$$K(x) = \tilde{\lambda}_\infty - \lambda a(x) \quad \text{and} \quad h(x, s) = \tilde{\lambda}_\infty s + f(s) \quad \text{for all } (x, s) \in \Omega \times [0, \infty). \quad (7.45)$$

It is clear that  $\text{essinf}_\Omega K \geq \tilde{\lambda}_\infty - \lambda_\infty > 0$ , so  $(H_K)$  is satisfied. Since  $f(0) = 0$ ,  $(H_0^\infty)$  holds too. Note that  $H(x, s)/s^2 = \tilde{\lambda}_\infty/2 + F(s)/s^2$ ,  $s > 0$ ; thus, hypothesis  $(f_1^\infty)$  implies  $(H_1^\infty)$ . Since  $l_\infty < -\tilde{\lambda}_\infty$ , there is a sequence  $\{s_i\}_i \subset (0, \infty)$  converging to  $+\infty$  such that  $f(s_i)/s_i < -\tilde{\lambda}_\infty$  for all  $i \in \mathbb{N}$ . By using the continuity of  $f$ , we may fix two sequences  $\{\delta_i\}_i, \{\eta_i\}_i \subset (0, \infty)$  such that  $0 < \delta_i < s_i < \eta_i < \delta_{i+1}$ ,  $\lim_{i \rightarrow \infty} \delta_i = \infty$ , and  $\tilde{\lambda}_\infty s + f(s) \leq 0$  for all  $s \in [\delta_i, \eta_i]$  and  $i \in \mathbb{N}$ . Therefore,  $(H_2^\infty)$  is also fulfilled. Now, we are in the position to apply Theorem 7.8. Throughout the choice (7.45),  $(P_h^K)$  is equivalent to problem  $(P_\lambda)$  which concludes the proof.

(ii) *Case  $p < 1$ .* Let  $\lambda \in \mathbb{R}$  be fixed arbitrarily and fix also a number  $\lambda_\infty \in (0, -l_\infty)$ . Now, let us choose

$$K(x) = \lambda_\infty \quad \text{and} \quad h(x, s) = \lambda a(x)s^p + \lambda_\infty s + f(s) \quad \text{for all } (x, s) \in \Omega \times [0, \infty). \quad (7.46)$$

Hypothesis  $(H_K)$  is clearly satisfied. Due to the fact that  $a \in L^\infty(\Omega)$ , hypothesis  $(H_0^\infty)$  holds too. Since  $p < 1$  and  $H(x, s)/s^2 = \lambda a(x)s^{p-1}/(p+1) + \lambda_\infty/2 + F(s)/s^2$ ,  $s > 0$ , hypothesis  $(f_1^\infty)$  implies  $(H_1^\infty)$ . For a.e.  $x \in \Omega$  and every  $s \in [0, \infty)$ , we have

$$h(x, s) \leq |\lambda| \cdot \|a\|_{L^\infty} s^p + \lambda_\infty s + f(s) \equiv \tilde{h}_\infty(s). \quad (7.47)$$

Thanks to  $(f_2^\infty)$ , we have

$$\liminf_{s \rightarrow \infty} \frac{\tilde{h}_\infty(s)}{s} = \lambda_\infty + l_\infty < 0.$$

Therefore, one can fix a sequence  $\{s_i\}_i \subset (0, \infty)$  converging to  $+\infty$  such that  $\tilde{h}_\infty(s_i) < 0$  for all  $i \in \mathbb{N}$ . Now, by using the continuity of  $\tilde{h}_\infty$ , one can fix two sequences  $\{\delta_i\}_i, \{\eta_i\}_i \subset (0, \infty)$  such that  $0 < \delta_i < s_i < \eta_i <$

$\delta_{i+1}$ ,  $\lim_{i \rightarrow \infty} \delta_i = \infty$ , and  $\tilde{h}_\infty(s) \leq 0$  for all  $s \in [\delta_i, \eta_i]$  and  $i \in \mathbb{N}$ . Thus, by using (7.47), hypothesis  $(H_2^\infty)$  holds. Now, we can apply Theorem 7.8, observing that problem  $(P_h^K)$  is equivalent to problem  $(P_\lambda)$  through the choice (7.46). Finally, in both cases (i) and (ii), (7.32) implies relation (7.3). This concludes the proof of Theorem 7.3.  $\square$

*Proof of Remark 7.4.* Assume that (7.4) holds. By contradiction, let us assume that there exists a bounded subsequence  $\{u_{k_i}^\infty\}_i$  of  $\{u_i^\infty\}_i$  in  $H_0^1(\Omega)$ . Since  $H_0^1(\Omega)$  is continuously embedded into  $L^t(\Omega)$ ,  $t \in [1, 2^*]$ , after an elementary estimate, we obtain that the sequence  $\{\mathcal{E}_{k_i}(u_{k_i}^\infty)\}_i$  is bounded. Since the sequence  $\{\mathcal{E}_i(u_i^\infty)\}_i$  is non-increasing, it will be bounded as well, which contradicts (7.37).  $\square$

*Proof of Theorem 7.4.* The proof is divided into five steps.

*Step 1.* Let  $\lambda_\infty \in (0, -l_\infty)$ . Due to  $(f_1^\infty)$ , we may fix a sequence  $\{s_i\}_i \subset (0, \infty)$  converging to  $\infty$ , such that  $f(s_i)/s_i < -\lambda_\infty$ . For every  $\lambda \in \mathbb{R}$ , let us define two functions  $h^\lambda : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$  and  $\tilde{h} : \mathbb{R}^2 \rightarrow \mathbb{R}$  by

$$h^\lambda(x, s) = \lambda a(x)s^p + \lambda_\infty s + f(s) \text{ for all } (x, s) \in \Omega \times [0, \infty);$$

$$\tilde{h}(\lambda, s) = |\lambda| \cdot \|a\|_{L^\infty} s^p + \lambda_\infty s + f(s) \text{ for all } s \in [0, \infty).$$

Note that  $\tilde{h}(0, s_i) = \lambda_\infty s_i + f(s_i) < 0$ . Due to the continuity of  $\tilde{h}$ , we can fix three sequences  $\{\delta_i\}_i, \{\eta_i\}_i \subset (0, \infty)$  and  $\{\lambda_i\}_i \subset (0, 1)$  such that  $0 < \delta_i < s_i < \eta_i < \delta_{i+1}$ ,  $\lim_{i \rightarrow \infty} \delta_i = \infty$ , and for every  $i \in \mathbb{N}$ ,

$$\tilde{h}(\lambda, s) \leq 0 \text{ for all } \lambda \in [-\lambda_i, \lambda_i] \text{ and } s \in [\delta_i, \eta_i]. \quad (7.48)$$

Without any loss of generality, we may assume that

$$\delta_i \geq i, \quad i \in \mathbb{N}. \quad (7.49)$$

Note that  $h^\lambda(x, s) \leq \tilde{h}(\lambda, s)$  for a.e.  $x \in \Omega$  and all  $(\lambda, s) \in \mathbb{R} \times [0, \infty)$ . Taking into account of (7.48), for every  $i \in \mathbb{N}$ , we have

$$h^\lambda(x, s) \leq 0 \text{ for a.e. } x \in \Omega \text{ and all } \lambda \in [-\lambda_i, \lambda_i], \quad s \in [\delta_i, \eta_i]. \quad (7.50)$$

For any  $i \in \mathbb{N}$  and  $\lambda \in [-\lambda_i, \lambda_i]$ , let  $h_i^\lambda : \Omega \times [0, \infty) \rightarrow \mathbb{R}$  be defined by

$$h_i^\lambda(x, s) = h^\lambda(x, \tau_{\eta_i}(s)) \quad (7.51)$$

and  $K(x) = \lambda_\infty$ . Let  $\mathcal{E}_{i,\lambda} : H_0^1(\Omega) \rightarrow \mathbb{R}$  be the energy functional associated with the problem  $(P_{h_i^\lambda}^K)$ , which is formally the same as in (7.26). Note that for every  $i \in \mathbb{N}$  and  $\lambda \in [-\lambda_i, \lambda_i]$ , the function  $h_i^\lambda$  fulfills the

hypotheses of Theorem 7.6; see (7.50) for  $(H_{h_i^\lambda}^2)$ . Consequently, for every  $i \in \mathbb{N}$  and  $\lambda \in [-\lambda_i, \lambda_i]$

$$\text{there exists } \tilde{u}_{i,\lambda}^\infty \in W^{\eta_i} \text{ with } \mathcal{E}_{i,\lambda}(\tilde{u}_{i,\lambda}^\infty) = \min_{W^{\eta_i}} \mathcal{E}_{i,\lambda}; \quad (7.52)$$

$$\tilde{u}_{i,\lambda}^\infty(x) \in [0, \delta_i] \text{ for a.e. } x \in \Omega, \quad (7.53)$$

$$\tilde{u}_{i,\lambda}^\infty \text{ is a weak solution of } (P_{h_i^\lambda}^K). \quad (7.54)$$

On account of the definition of the functions  $h_i^\lambda$  and  $K$ , and relations (7.54) and (7.53),  $\tilde{u}_{i,\lambda}^\infty$  is also a weak solution for our initial problem  $(P_\lambda)$  once we have  $\tilde{u}_{i,\lambda}^\infty \not\equiv 0$ .

*Step 2.* Note that for  $\lambda = 0$ , the function  $h_i^\lambda = h_i^0$  verifies the hypotheses of Theorem 7.8; in fact,  $h_i^0$  is the function appearing in (7.33) and  $\mathcal{E}_i := \mathcal{E}_{i,0}$  is the energy functional associated with problem  $(P_{h_i^0}^K)$ . Denoting  $u_i^\infty := \tilde{u}_{i,0}^\infty$ , we also have

$$\mathcal{E}_{m_i}(u_{m_i}^\infty) = \min_{W^{\eta_{m_i}}} \mathcal{E}_{m_i} \leq \mathcal{E}_{m_i}(z_{\tilde{s}_i}); \quad (7.55)$$

$$\lim_{i \rightarrow \infty} \mathcal{E}_{m_i}(u_{m_i}^\infty) = -\infty, \quad (7.56)$$

where the special subsequence  $\{u_{m_i}^\infty\}_i$  of  $\{u_i^\infty\}_i$  and  $z_{\tilde{s}_i} \in W^{\eta_{m_i}}$  appear in the proof of Theorem 7.8, see relations (7.42) and (7.43), respectively.

*Step 3.* Let us fix a sequence  $\{\theta_i\}_i$  with negative terms such that  $\lim_{i \rightarrow \infty} \theta_i = -\infty$ . Due to (7.55) and (7.56), up to a subsequence, we may assume that

$$\theta_{i+1} < \mathcal{E}_{m_i}(u_{m_i}^\infty) \leq \mathcal{E}_{m_i}(z_{\tilde{s}_i}) < \theta_i. \quad (7.57)$$

For any  $i \in \mathbb{N}$ , define

$$\lambda'_i = \frac{(p+1)(\theta_i - \mathcal{E}_{m_i}(z_{\tilde{s}_i}))}{\delta_{m_i}^{p+1}(\|a\|_{L^1} + 1)} \quad \text{and} \quad \lambda''_i = \frac{(p+1)(\mathcal{E}_{m_i}(u_{m_i}^\infty) - \theta_{i+1})}{\delta_{m_i}^{p+1}(\|a\|_{L^1} + 1)}.$$

Fix  $k \in \mathbb{N}$ . Thanks to (7.57),

$$\lambda_k^\infty = \min(\lambda_1, \dots, \lambda_k, \lambda'_1, \dots, \lambda'_k, \lambda''_1, \dots, \lambda''_k) > 0.$$

Therefore, for every  $i \in \{1, \dots, k\}$  and  $\lambda \in [-\lambda_k^\infty, \lambda_k^\infty]$  we have

$$\begin{aligned} \mathcal{E}_{m_i,\lambda}(\tilde{u}_{m_i,\lambda}^\infty) &\leq \mathcal{E}_{m_i,\lambda}(z_{\tilde{s}_i}) && \text{(see (7.52))} \\ &= \frac{1}{2} \|z_{\tilde{s}_i}\|_{H_0^1}^2 - \frac{\lambda}{p+1} \int_{\Omega} a(x) z_{\tilde{s}_i}^{p+1} - \int_{\Omega} F(z_{\tilde{s}_i}(x)) dx \\ &= \mathcal{E}_{m_i}(z_{\tilde{s}_i}) - \frac{\lambda}{p+1} \int_{\Omega} a(x) z_{\tilde{s}_i}^{p+1} \\ &< \theta_i, && \text{(see the choice of } \lambda'_i \text{ and } \tilde{s}_i \leq \delta_{m_i}) \end{aligned}$$

and since  $\tilde{u}_{m_i, \lambda}^\infty$  belongs to  $W^{\eta_{m_i}}$ , and  $u_{m_i}^\infty$  is the minimum point of  $\mathcal{E}_{m_i}$  over the set  $W^{\eta_{m_i}}$ , see relation (7.55), we have

$$\begin{aligned} \mathcal{E}_{m_i, \lambda}(\tilde{u}_{m_i, \lambda}^\infty) &= \mathcal{E}_{m_i}(\tilde{u}_{m_i, \lambda}^\infty) - \frac{\lambda}{p+1} \int_{\Omega} a(x)(\tilde{u}_{m_i, \lambda}^\infty)^{p+1} \\ &\geq \mathcal{E}_{m_i}(u_{m_i}^\infty) - \frac{\lambda}{p+1} \int_{\Omega} a(x)(\tilde{u}_{m_i, \lambda}^\infty)^{p+1} \\ &> \theta_{i+1}. \end{aligned} \quad (\text{see the choice of } \lambda_i' \text{ and (7.53)})$$

Consequently, for every  $i \in \{1, \dots, k\}$  and  $\lambda \in [-\lambda_k^\infty, \lambda_k^\infty]$  we have

$$\theta_{i+1} < \mathcal{E}_{m_i, \lambda}(\tilde{u}_{m_i, \lambda}^\infty) < \theta_i < 0, \quad (7.58)$$

therefore

$$\mathcal{E}_{m_k, \lambda}(\tilde{u}_{m_k, \lambda}^\infty) < \dots < \mathcal{E}_{m_1, \lambda}(\tilde{u}_{m_1, \lambda}^\infty) < 0. \quad (7.59)$$

Note that  $\tilde{u}_{m_i, \lambda}^\infty \in W^{\eta_{m_k}}$  for every  $i \in \{1, \dots, k\}$ , so  $\mathcal{E}_{m_i, \lambda}(\tilde{u}_{m_i, \lambda}^\infty) = \mathcal{E}_{m_k, \lambda}(\tilde{u}_{m_i, \lambda}^\infty)$ , see relation (7.51). From above, for every  $\lambda \in [-\lambda_k^\infty, \lambda_k^\infty]$ , we have

$$\mathcal{E}_{m_k, \lambda}(\tilde{u}_{m_k, \lambda}^\infty) < \dots < \mathcal{E}_{m_k, \lambda}(\tilde{u}_{m_1, \lambda}^\infty) < 0 = \mathcal{E}_{m_k, \lambda}(0).$$

In particular, the elements  $\tilde{u}_{m_1, \lambda}^\infty, \dots, \tilde{u}_{m_k, \lambda}^\infty$  are distinct (and non-trivial) whenever  $\lambda \in [-\lambda_k^\infty, \lambda_k^\infty]$ .

*Step 4.* Assume that  $k \geq 2$  and fix  $\lambda \in [-\lambda_k^\infty, \lambda_k^\infty]$ . We prove that

$$\|\tilde{u}_{m_i, \lambda}^\infty\|_{L^\infty} > \delta_{m_{i-1}} \quad \text{for all } i \in \{2, \dots, k\}. \quad (7.60)$$

Let us assume that there exists an element  $i_0 \in \{2, \dots, k\}$  such that  $\|\tilde{u}_{m_{i_0}, \lambda}^\infty\|_{L^\infty} \leq \delta_{m_{i_0-1}}$ . Since  $\delta_{m_{i_0-1}} < \eta_{m_{i_0-1}}$ , then  $\tilde{u}_{m_{i_0}, \lambda}^\infty \in W^{\eta_{m_{i_0-1}}}$ . Thus, on account of (7.52) and (7.51), we have

$$\begin{aligned} \mathcal{E}_{m_{i_0-1}, \lambda}(\tilde{u}_{m_{i_0}, \lambda}^\infty) &= \min_{W^{\eta_{m_{i_0-1}}}} \mathcal{E}_{m_{i_0-1}, \lambda} \leq \mathcal{E}_{m_{i_0-1}, \lambda}(\tilde{u}_{m_{i_0}, \lambda}^\infty) \\ &= \mathcal{E}_{m_{i_0}, \lambda}(\tilde{u}_{m_{i_0}, \lambda}^\infty), \end{aligned}$$

which contradicts (7.59). Therefore, relation (7.60) holds true.

*Step 5.* Let  $u_{i, \lambda}^\infty := \tilde{u}_{m_i, \lambda}^\infty$  for any  $i \in \{1, \dots, k\}$  and  $\lambda \in [-\lambda_k^\infty, \lambda_k^\infty]$ ; these elements verify all the requirements of Theorem 7.4. Indeed, since  $\mathcal{E}_{m_1, \lambda}(u_{1, \lambda}^\infty) = \mathcal{E}_{m_1, \lambda}(\tilde{u}_{m_1, \lambda}^\infty) < 0 = \mathcal{E}_{m_1, \lambda}(0)$ , then  $\|u_{1, \lambda}^\infty\|_{L^\infty} > 0$ , which proves (7.5) for  $i = 1$ . If  $k \geq 2$ , then on account of Step 4, (7.49) and  $m_i \geq i$ , for every  $i \in \{2, \dots, k\}$ , we have

$$\|u_{i, \lambda}^\infty\|_{L^\infty} > \delta_{m_{i-1}} \geq m_{i-1} \geq i - 1,$$

that is, relation (7.5) holds true. This ends the proof of Theorem 7.4.  $\square$

*Proof of Remark 7.6.* Due to (7.4), there exists a  $C > 0$  such that  $|f(s)| \leq C(1 + s^{2^*-1})$  for all  $s \geq 0$ . We denote by  $S_t > 0$  the Sobolev embedding constant of the continuous embedding  $H_0^1(\Omega) \hookrightarrow L^t(\Omega)$ ,  $t \in [1, 2^*]$ . Without any loss of generality, we may assume that for every  $i \in \mathbb{N}$ ,

$$\theta_i < -\frac{1}{p+1} \|a\|_{L^\infty} S_{p+1}^{p+1} (i-1)^{p+1} - C[S_1(i-1) + S_{2^*}^{2^*} (i-1)^{2^*}], \quad (7.61)$$

where the sequence  $\{\theta_i\}_i$  comes from Step 3 of the proof of Theorem 7.4.

Fix  $\lambda \in [-\lambda_k^\infty, \lambda_k^\infty]$  and assume that there exists  $i_0 \in \{1, \dots, k\}$  such that  $\|u_{i_0, \lambda}^\infty\|_{H_0^1} \leq i_0 - 1$ . On account of (7.58), we have in particular  $\mathcal{E}_{m_{i_0}, \lambda}(u_{i_0, \lambda}^\infty) < \theta_{i_0}$ . Consequently, we have

$$\begin{aligned} \frac{1}{2} \|u_{i_0, \lambda}^\infty\|_{H_0^1}^2 &= \mathcal{E}_{m_{i_0}, \lambda}(u_{i_0, \lambda}^\infty) + \frac{\lambda}{p+1} \int_{\Omega} a(x) (u_{i_0, \lambda}^\infty)^{p+1} + \int_{\Omega} F(u_{i_0, \lambda}^\infty(x)) dx \\ &< \theta_{i_0} + \frac{|\lambda|}{p+1} \|a\|_{L^\infty} S_{p+1}^{p+1} \|u_{i_0, \lambda}^\infty\|_{H_0^1}^{p+1} \\ &\quad + C[S_1 \|u_{i_0, \lambda}^\infty\|_{H_0^1} + S_{2^*}^{2^*} \|u_{i_0, \lambda}^\infty\|_{H_0^1}^{2^*}] \\ &\leq \theta_{i_0} + \frac{1}{p+1} \|a\|_{L^\infty} S_{p+1}^{p+1} (i_0 - 1)^{p+1} \\ &\quad + C[S_1 (i_0 - 1) + S_{2^*}^{2^*} (i_0 - 1)^{2^*}] \quad (\lambda_k^\infty \leq 1) \\ &< 0, \quad (\text{see (7.61)}) \end{aligned}$$

contradiction.  $\square$

## 7.6 Perturbation from symmetry

For details related to the main notions and properties used in this section we refer to Appendix D.

Let  $\Omega \subset \mathbb{R}^N$  be a bounded open set. For some fixed  $r > 0$ , consider the following nonlinear eigenvalue problem: find  $(u, \lambda) \in H_0^1(\Omega) \times \mathbb{R}$  such that

$$\begin{cases} f(x, u) \in L_{\text{loc}}^1(\Omega), \\ -\Delta u = \lambda f(x, u) \quad \text{in } \Omega, \\ \int_{\Omega} |\nabla u|^2 dx = r^2, \end{cases} \quad (7.62)$$

where  $f : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$  is a Carathéodory function such that the following conditions hold:

- (f1)  $f(x, -s) = -f(x, s)$ , for a.e.  $x \in \Omega$  and every  $s \in \mathbb{R}$ ;  
 (f2) there exist  $a \in L^1(\Omega)$ ,  $b \in \mathbb{R}$  and  $0 \leq p < \frac{2N}{N-2}$  (if  $N > 2$ ) such that

$$0 < sf(x, s) \leq a(x) + b|s|^p, \quad F(x, s) \leq a(x) + b|s|^p,$$

for a.e.  $x \in \Omega$  and every  $s \in \mathbb{R} \setminus \{0\}$ , where  $F(x, s) = \int_0^s f(x, t)dt$ ;

- (f3)  $\sup_{|s| \leq t} |f(x, s)| \in L^1_{\text{loc}}(\Omega)$ , for every  $t > 0$ .

We point out that if  $N = 1$ , then in condition (f2) the term  $b|s|^p$  can be substituted by any continuous function  $\varphi(s)$  of  $s$ , while, if  $N = 2$ , the same term can be substituted by  $\exp(\varphi(s))$ , with  $\varphi(s)s^{-2} \rightarrow 0$  as  $|s| \rightarrow \infty$ .

Our first result in this section is the following.

**Theorem 7.9** *Assume that hypotheses (f1) – (f3) hold. Then problem (7.62) admits a sequence  $(\pm u_n, \lambda_n)$  of distinct solutions.*

Our main purpose is to study what happens when the energy functional is affected by an *arbitrary* perturbation which destroys the symmetry.

Consider the problem: find  $(u, \lambda) \in H_0^1(\Omega) \times \mathbb{R}$  such that

$$\begin{cases} f(x, u), g(x, u) \in L^1_{\text{loc}}(\Omega), \\ -\Delta u = \lambda(f(x, u) + g(x, u)) & \text{in } \Omega, \\ \int_{\Omega} |\nabla u|^2 dx = r^2, \end{cases} \quad (7.63)$$

where  $g : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$  is a Carathéodory function. We make no symmetry assumption on  $g$ , but we impose only

- (g1)  $0 < sg(x, s) \leq a(x) + b|s|^p$  for a.e.  $x \in \Omega$  and every  $s \in \mathbb{R} \setminus \{0\}$ ;  
 (g2)  $\sup_{|s| \leq t} |g(x, s)| \in L^1_{\text{loc}}(\Omega)$ , for every  $t > 0$ ;  
 (g3)  $G(x, s) \leq C_g(1 + |s|^p)$ , for a.e.  $x \in \Omega$  and every  $s \in \mathbb{R}$ , for some  $C_g > 0$ , where  $G(x, s) = \int_0^s g(x, t)dt$ .

The next result shows that the number of solutions of problem (7.63) becomes greater and greater, as the perturbation tends to zero.

**Theorem 7.10** *Assume that hypotheses (f1) – (f3) and (g1) – (g3) hold. Then, for every positive integer  $n$ , there exists  $\varepsilon_n > 0$  such that problem (7.63) admits at least  $n$  distinct solutions, provided that (g3) holds for  $C_g = \varepsilon_n$ .*

We prove theorems 7.9 and 7.10 by a variational argument. First we set

$$S_r = \left\{ u \in H_0^1(\Omega) : \int_{\Omega} |\nabla u|^2 dx = r^2 \right\}$$

and we study the critical points on  $S_r$  of the even continuous functional  $I : H_0^1(\Omega) \rightarrow \mathbb{R}$  defined by

$$I(u) = - \int_{\Omega} F(x, u) dx.$$

We point out that if conditions (f2), (f3) are substituted by the more standard condition  $0 < sf(x, s) \leq a_1(x)|s| + b|s|^p$  with  $a_1 \in L^{\frac{2N}{N+2}}(\Omega)$ , then  $I$  is of class  $C^1$  and Theorem 7.9 can be found in [253, Theorem 8.17]. Under our assumptions,  $f$  could have the form  $f(x, s) = \alpha(x)\gamma(s)$  with  $\alpha \in L^1(\Omega)$ ,  $\alpha \geq 0$ ,  $\gamma \in C_c(\mathbb{R})$ ,  $\gamma$  odd and  $s\gamma(s) \geq 0$  for any  $s \in \mathbb{R}$ . In such a case,  $I$  is clearly continuous, but not locally Lipschitz.

When  $f$  and  $g$  are subjected to the standard condition we have mentioned, results like Theorem 7.10 go back to Krasnoselskii [163]. For perturbation results, quite different from ours, where the perturbed problem still has infinitely many solutions, we refer the reader to [253, 284]. In a nonsmooth setting, a result in the line of Theorem 7.10 has been proved in [83] when  $f$  and  $g$  satisfy the standard condition, but the function  $u$  is subjected to an obstacle, so that the equation becomes a variational inequality.

We first observe that from our assumption (f2) it follows that  $I(u) < 0$  and  $\sup I_r(u) = 0$ , where  $I_r = I|_{S_r}$ .

We start with the following preliminary result.

**Lemma 7.1** *The following properties hold true:*

- (a) *if  $u \in S_r$  satisfies  $|dI_r|(u) < +\infty$ , then  $f(x, u) \in L_{\text{loc}}^1(\Omega) \cap H^{-1}(\Omega)$  and there exists  $\mu \in \mathbb{R}$  such that*

$$\|\mu \Delta u + f(x, u)\|_{H^{-1}} \leq |dI_r|(u);$$

- (b) *the functional  $I_r$  satisfies  $(PS)_c$  for any  $c < 0$ ;*  
 (c) *if  $u \in S_r$  is a critical point of  $I_r$ , then there exists  $\lambda > 0$  such that  $(u, \lambda)$  is a solution of problem (7.62).*

*Proof* (a) Set

$$I_{r, \text{est}}(w) = \begin{cases} I(w) & \text{if } w \in S_r, \\ +\infty & \text{if } w \in H_0^1(\Omega) \setminus S_r. \end{cases}$$

Then  $|dI_{r,est}|(u) = |dI_r|(u)$ . We also deduce that there exists  $\alpha \in \partial I_{r,est}(u)$  with  $\|\alpha\|_{H^{-1}} \leq |dI_{r,est}|(u)$ , where  $\partial$  stands for the generalized gradient of  $I_{r,est}$ . Taking into account (f2), we obtain

$$I^0(u; 0) \leq 0, \quad I^0(u; 2u) \leq -2 \int_{\Omega} f(x, u) u dx < +\infty.$$

Actually, the same proof shows a stronger fact, namely that

$$\bar{I}^0(u; 0) \leq 0, \quad \bar{I}^0(u; 2u) \leq -2 \int_{\Omega} f(x, u) u dx < +\infty.$$

Thus, there are  $\beta \in \partial I(u)$  and  $\mu \in \mathbb{R}$  such that  $\alpha = \beta - \mu \Delta u$ . We conclude that  $f(x, u) \in L^1_{loc}(\Omega) \cap H^{-1}(\Omega)$  and  $\beta = -f(x, u)$ . Then (a) follows.

(b) Let  $c < 0$  and let  $(u_n)$  be a  $(PS)_c$ -sequence for  $I_r$ . By the previous point, we have  $f(x, u_n) \in L^1_{loc}(\Omega) \cap H^{-1}(\Omega)$  and there exists a sequence  $(\mu_n)$  in  $\mathbb{R}$  with

$$\|\mu_n \Delta u_n + f(x, u_n)\|_{H^{-1}} \rightarrow 0.$$

Up to a subsequence,  $(u_n)$  is convergent to some  $u$  weakly in  $H^1_0(\Omega)$  and a.e. From (f2) it follows  $I(u) = c < 0$ , hence  $u \neq 0$ . Again by (f2) and Lebesgue's dominated convergence theorem, we deduce that

$$0 < \int_{\Omega} f(x, u) u dx = \lim_n \int_{\Omega} f(x, u_n) u_n dx = \lim_n \mu_n \int_{\Omega} |\nabla u_n|^2 dx.$$

Therefore, up to a further subsequence,  $(\mu_n)$  is convergent to some  $\mu > 0$  and

$$\left\| \Delta u_n + \frac{1}{\mu} f(x, u_n) \right\|_{H^{-1}} \rightarrow 0.$$

This shows that  $(u_n)$  is precompact in  $H^1_0(\Omega)$  and (b) follows.

(c) Arguing as in (b), we find that  $f(x, u) \in L^1_{loc}(\Omega) \cap H^{-1}(\Omega)$  and that there exists  $\mu > 0$  with  $\mu \Delta u + f(x, u) = 0$ . This concludes the proof.  $\square$

**Lemma 7.2** *There exists a sequence  $(b_n)$  of essential values of  $I_r$  strictly increasing to 0.*

*Proof* Let  $\psi : ]-\infty, 0[ \rightarrow \mathbb{R}$  be an increasing diffeomorphism. From Lemma 7.1 it follows that  $\psi \circ I_r$  satisfies condition  $(PS)_c$  for every  $c \in \mathbb{R}$ . Then the set  $\{u \in S_r : \psi \circ I_r(u) \leq b\}$  has finite genus for every  $b \in \mathbb{R}$ . For any integer  $n \geq 1$ , set

$$c_n := \inf_{S \in \Gamma_n} \sup_{u \in S} \psi \circ I_r(u),$$

where  $\Gamma_n := \{S \subset S_r; S \in \mathcal{F}, \gamma(S) \geq n\}$  and  $\mathcal{F}$  denotes the family of closed and symmetric subsets of  $S_r$  with respect to the origin. Since  $c_n \rightarrow +\infty$  as  $n \rightarrow \infty$ , it follows that there exists a sequence  $(b'_n)$  of essential values of  $\psi \circ I_r$  strictly increasing to  $+\infty$ . Then  $b_n = \psi^{-1}(b'_n)$  has the required properties.  $\square$

*Proof of Theorem 7.9.* Using Lemma 7.1, we deduce that each  $b_n$  is a critical value of  $I_r$ . Again from Lemma 7.1, we conclude that there exists a sequence  $(\pm u_n, \lambda_n)$  of solutions of problem (7.62) with  $I(u_n) = b_n$  strictly increasing to 0. This concludes the proof of Theorem 7.9.

Let us now introduce the continuous functional  $J_r : S_r \rightarrow \mathbb{R}$  defined by

$$J(u) = I(u) - \int_{\Omega} G(x, u) dx.$$

**Lemma 7.3** *For every  $\eta > 0$ , there exists  $\varepsilon > 0$  such that  $\sup_{u \in S_r} |I_r(u) - J_r(u)| < \eta$ , provided that (g3) holds for  $C_g = \varepsilon$ .*

*Proof* By Sobolev inclusions, we have for any  $u \in S_r$ ,

$$0 \leq I_r(u) - J_r(u) = \int_{\Omega} G(x, u) dx \leq C_g \int_{\Omega} (1 + |u|^p) dx < \eta,$$

if  $g$  is chosen as in the hypothesis.  $\square$

*Proof of Theorem 7.10.* As in the proof of Theorem 7.9, let us consider a strictly increasing sequence  $(b_n)$  of essential values of  $I_r$  such that  $b_n \rightarrow 0$  as  $n \rightarrow \infty$ . Given  $n \geq 1$ , take some  $\delta > 0$  with  $b_n + \delta < 0$  and  $2(b_j - b_{j-1}) < \delta$  for  $j = 2, \dots, n$ . Thus, for any  $j = 1, \dots, n$ , there exists  $\eta_j > 0$  such that  $\sup_{u \in S_r} |I_r(u) - J_r(u)| < \eta_j$  implies the existence of an essential value  $c_j \in ]b_j - \delta, b_j + \delta[$  of  $J_r$ . We now apply Lemma 7.3 for  $\eta = \min\{\eta_1, \dots, \eta_n\}$ . Thus we obtain  $\varepsilon_n > 0$  such that  $\sup_{u \in S_r} |I_r(u) - J_r(u)| < \eta$ , if (g3) holds with  $C_g = \varepsilon_n$ . It follows that  $J_r$  has at least  $n$  distinct essential values  $c_1, \dots, c_n$  in the interval  $] -\infty, 0[$ .

Now Lemma 1 can be clearly adapted to the functional  $J_r$ . Then we find  $u_1, \dots, u_n \in S_r$  and  $\lambda_1, \dots, \lambda_n > 0$  such that each  $(u_j, \lambda_j)$  is a solution of Problem (7.62) with  $J_r(u_j) = c_j$ .

## 7.7 Historical notes, comments

A. *Historical notes.* Let us consider the equation

$$\begin{cases} -\Delta u = f(x, u) + \lambda g(x, u) & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (\text{E}_\lambda)$$

where  $\Omega \subseteq \mathbb{R}^N$  is a bounded open domain, while  $f, g : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$  are Carathéodory functions. We assume the unperturbed equation  $(\text{E}_0)$  has *infinitely many* distinct solutions. Then, the main question is:

- (q). *Fixing  $k \in \mathbb{N}$ , can one find a number  $\lambda_k > 0$  such that the perturbed equation  $(\text{E}_\lambda)$  has at least  $k$  distinct solutions whenever  $\lambda \in [-\lambda_k, \lambda_k]$ ?*

Two different classes of results are available in the literature answering affirmatively question (q):

1) *Perturbation of symmetric problems.* Assume  $f(x, s) = -f(x, -s)$  for every  $(x, s) \in \Omega \times \mathbb{R}$ . It is well-known that if the energy functional has the mountain pass geometry, problem  $(\text{E}_0)$  has infinitely many solutions, due to the symmetric version of the Mountain Pass theorem, see Ambrosetti-Rabinowitz [8]. Furthermore, question (q) was fully answered by Li-Liu [189] for arbitrarily continuous nonlinearity  $g$ , following the topological approach developed by Degiovanni-Lancelotti [83] and Degiovanni-Rădulescu [87];

2) *Perturbation of oscillatory problems.* Assume  $f(x, \cdot)$  oscillates near the origin or at infinity, uniformly with respect to  $x \in \Omega$ . Special kinds of oscillations produce infinitely many solutions for  $(\text{E}_0)$ , as shown by Omari-Zanolin [232], and Saint Raymond [269]. Concerning the perturbed problem, Anello-Cordaro [13] answered question (q), by using Theorem 1.21.

B. *Comments.* In this chapter we presented a third, *direct method* for answering question (q) whenever the nonlinear term  $f(x, \cdot)$  belongs to a wide class of oscillatory functions and  $g(x, s) = s^p$ . We fully described the effect of the term  $g$  to the oscillatory function  $f$ . Namely, the number of distinct solutions of  $(\text{E}_\lambda)$  is sensitive when:

- $p \leq 1$  and  $f$  oscillates near the origin;
- $p \geq 1$  and  $f$  oscillates at infinity.

Notice that  $g$  can be any continuous function with  $g(x, 0) = 0$ , and our method works also for other elliptic problems.

## 8

### Problems to Part I

I hope that seeing the excitement of solving this problem will make young mathematicians realize that there are lots and lots of other problems in mathematics which are going to be just as challenging in the future.

---

Andrew Wiles (b. 1953)

**Problem 8.1** Show that any function  $f : \mathbb{R} \rightarrow \mathbb{R}$  of class  $C^1$  having two local minima has necessarily a third critical point.

**Problem 8.2** Show that the  $(PS)_c$ -condition implies the  $(C)_c$ -condition for any function  $f : X \rightarrow \mathbb{R}$  of class  $C^1$  and  $c \in \mathbb{R}$ , see Remark 1.3. Construct a function which shows that the converse does not hold in general.

**Problem 8.3** Prove that every function  $f : T^2 \rightarrow \mathbb{R}$  of class  $C^1$  has at least 3 critical points. Here  $T^2 = S^1 \times S^1$  is the 2-dimensional torus. [Hint: Prove that  $\text{cat}(T^2) = 3$  and apply Theorem 1.11.]

**Problem 8.4** Let  $X$  be a real Banach space and assume that  $f$  is a nonnegative  $C^1$  function on  $X$  satisfying the Palais–Smale condition or even a weaker form: every sequence  $(u_n)$  in  $X$  such that  $\sup_n f(u_n) < \infty$  and  $\|f'(u_n)\| \rightarrow 0$ , is bounded in  $X$ . Prove that  $f$  is coercive, that is,

$$f(u) \rightarrow +\infty \quad \text{as } \|u\| \rightarrow \infty.$$

**Problem 8.5** Let  $X$  be a real Banach space and assume that  $f$  is

a nonnegative  $C^1$  function on  $X$  satisfying the Palais–Smale condition. Prove that every minimizing sequence of  $f$  has a convergent subsequence.

The following exercise complements the results of Corollary 1.3.

**Problem 8.6** Let  $f$  be a real-valued lower semicontinuous, Gâteaux differentiable, and bounded from below functional on a Hilbert space  $X$ . Assume that  $(u_n)$  is a minimizing sequence of  $f$ . Prove that there exists a minimizing sequence  $(v_n)$  of  $f$  such that the following properties hold true:

$$\lim_{n \rightarrow \infty} \|u_n - v_n\| = 0; \quad \lim_{n \rightarrow \infty} \|f'(v_n)\| = 0.$$

Furthermore, one has the second-order inequality

$$0 \leq \liminf_{n \rightarrow \infty} \left[ \liminf_{x \rightarrow v_n} \frac{f(x) - f(v_n) - f'(v_n)(x - v_n)}{\|u_n - v_n\|^2} \right].$$

**Problem 8.7** (Brézis–Nirenberg, [51]). Let  $X$  be a real Banach space and assume that  $f$  is a  $C^1$  function on  $X$  which is bounded below and satisfies the Palais–Smale condition. Suppose that all the critical points of  $f$  lie in  $\{u \in X; \|u\| < R\}$ . Set

$$M(r) = \inf\{f(u); \|u\| = r\}.$$

Prove that for  $r > R$ ,  $M(r)$  is strictly increasing and continuous from the right.

**Problem 8.8** Let  $\Omega$  be a bounded open set in  $\mathbb{R}^N$ . For every  $f \in L^2(\Omega)$  and all  $\lambda \in \mathbb{R}$ , define the functional

$$E_\lambda(u) = \frac{1}{2} \int_{\Omega} (|\nabla u|^2 - \lambda u^2) dx - \int_{\Omega} f u dx, \quad u \in H_0^1(\Omega).$$

a) Prove that if  $\lambda$  is not an eigenvalue of  $(-\Delta)$  in  $H_0^1(\Omega)$ , then  $E_\lambda$  satisfies the Palais–Smale condition in  $H_0^1(\Omega)$ .

b) Prove that if  $\lambda$  is an eigenvalue of  $(-\Delta)$  in  $H_0^1(\Omega)$  and if  $f = 0$ , then  $E_\lambda$  does not satisfy the Palais–Smale condition in  $H_0^1(\Omega)$ .

**Problem 8.9** Let  $X$  be a real Banach space and let  $U$  be an open subset of  $X$ . Let  $\Phi$  be a real  $C^1$  bounded below function on  $\bar{U}$ . Set

$$a = \inf_{\bar{U}} \Phi.$$

Assume  $F$  is a closed subset of  $U$  such that  $\text{dist}(F, \partial U) > 0$  (no assumption if  $U = X$ ) and

$$a = \inf_F \Phi.$$

Prove that there exists a sequence  $(x_n)$  in  $U$  such that

$$\Phi(x_n) \rightarrow a, \quad \Phi'(x_n) \rightarrow 0 \text{ in } X^* \quad \text{and} \quad \text{dist}(x_n, F) \rightarrow 0.$$

**Problem 8.10** In Theorem 1.3, instead of the assumption that  $f$  is bounded from below, assume that

$$f(x) + \varphi(d(x, x_0)) \geq 0 \quad \text{for every } x \in X,$$

for some  $x_0 \in X$  and some function  $\varphi(t) = o(t)$  as  $t \rightarrow +\infty$ . Prove that the conclusion of Theorem 1.3 still holds.

**Problem 8.11** (Caristi's fixed point theorem). Let  $(X, d)$  be a complete metric space and let  $f : X \rightarrow X$  be a function. Assume that  $\Psi : X \rightarrow \mathbb{R}$  is a lower semicontinuous function such that

$$d(x, f(x)) \leq \Psi(x) - \Psi(f(x)) \quad \text{for all } x \in X.$$

Prove that  $f$  has a fixed point.

**Problem 8.12** Let  $\Phi$  be a  $C^1$  function on  $\mathbb{R}^N$  such that

$$m = \liminf_{\|x\| \rightarrow \infty} \Phi(x) \quad \text{is finite.}$$

Prove that there exists a sequence of points  $(x_m)$  in  $\mathbb{R}^N$  such that  $\|x_m\| \rightarrow \infty$ ,  $\Phi(x_m) \rightarrow m$ ,  $\|\Phi'(x_m)\| \rightarrow 0$  and  $\Phi'(x_m)$  is parallel to  $x_m$  ( $\Phi'(x_m)$  may be zero).

**Problem 8.13** Let  $X$  be a real Banach space and assume that  $f : X \rightarrow \mathbb{R}$  is a function of class  $C^1$  such that

$$\inf_{\|u - e_0\| = \rho} f(u) \geq \alpha > \max\{f(e_0), f(e_1)\}$$

for some  $\alpha \in \mathbb{R}$  and  $e_0 \neq e_1 \in X$  with  $0 < \rho < \|e_0 - e_1\|$ . Set

$$c = \inf_{\gamma \in \Gamma} \max_{t \in [0, 1]} f(\gamma(t)),$$

where

$$\Gamma = \{\gamma \in C([0, 1], X) : \gamma(0) = e_0, \gamma(1) = e_1\}.$$

Prove that the conclusion of Theorem 1.7 can be strengthened as follows: there exists a sequence  $(u_n)$  in  $X$  such that  $f(u_n) \rightarrow c$  and  $(1 + \|u_n\|) \|f'(u_n)\| \rightarrow 0$  as  $n \rightarrow \infty$ .

**Problem 8.14** Prove the same result if  $(1 + \|u\|)$  is replaced by  $\varphi(\|u\|)$ , where  $\varphi \geq 1$  and  $\int_0^\infty \frac{dt}{\varphi(t)} = +\infty$ .

**Problem 8.15** Let  $X$  be a real Banach space and assume that  $f, g : X \rightarrow \mathbb{R}$  are functions of class  $C^1$  satisfying the Palais–Smale condition with  $f \geq g$  on  $X$ . Let  $K$  be a compact metric space and let  $K^*$  be a nonempty closed subset of  $K$ ,  $K^* \neq K$ . Define

$$\mathcal{P} = \{p \in C(K, X); p = p^* \text{ on } K^*\},$$

where  $p^*$  is a fixed continuous map on  $K$ . Assume that

$$c = \inf_{p \in \mathcal{P}} \max_{t \in K} f(p(t)) = \inf_{p \in \mathcal{P}} \max_{t \in K} g(p(t)) > \inf_{p \in \mathcal{P}} \max_{t \in K^*} f(p^*(t)).$$

Prove that  $f$  and  $g$  have a common critical point where both functions equal  $c$ .

**Problem 8.16** Let  $\Omega \subset \mathbb{R}^N$  be a bounded domain with smooth boundary and assume that  $p$  is a real number such that  $1 < p < (N+2)/(N-2)$  if  $N \geq 3$  and  $1 < p < \infty$  if  $N \in \{1, 2\}$ .

Prove that the nonlinear problem

$$\begin{cases} -\Delta u + u = u^p & \text{in } \Omega \\ u > 0 & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega. \end{cases}$$

has a solution.

**Problem 8.17** Establish the same existence result for the problem

$$\begin{cases} -\Delta u + u = f(x, u) & \text{in } \Omega \\ u > 0 & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

where

$$f(x, u) = o\left(|u|^{(N+2)/(N-2)}\right) \quad \text{as } u \rightarrow +\infty, \text{ uniformly in } x \in \bar{\Omega}.$$

**Problem 8.18** Let  $\Omega \subset \mathbb{R}^N$  be a bounded domain with smooth boundary and assume that  $f$  is a  $C^1$  function on  $[0, 1]$  such that  $f > 0$  on  $(0, 1)$  and  $f(0) = f'(0) = f(1) = 0$ . Suppose that  $\underline{u} \in C_0^1(\bar{\Omega})$  satisfies  $0 < \underline{u} \leq 1$  in  $\Omega$  and  $-\Delta \underline{u} \leq f(\underline{u})$  in  $\Omega$ .

Consider the nonlinear elliptic problem

$$\begin{cases} -\Delta u = f(u) & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega. \end{cases} \quad (8.1)$$

Prove that if  $\underline{u}$  is not a solution of (8.1), then there are at least two solutions  $u_1$  and  $u_2$  of problem (8.1) with  $0 < u_1 < u_2 < 1$  in  $\Omega$ .

**Problem 8.19** Let  $\Omega \subset \mathbb{R}^N$  be a bounded domain with smooth boundary,  $\lambda \in \mathbb{R}$ , and  $p > 1$ . For any integer  $j \geq 1$ , set

$$\Gamma_j := \{A \subset H_0^1(\Omega); A \text{ is compact, symmetric, } 0 \notin A, \gamma(A) \geq j\}.$$

Define the functional

$$E(u) := \frac{1}{2} \int_{\Omega} [|\nabla u|^2 - \lambda u^2] dx - \frac{1}{p+1} \int_{\Omega} |u|^{p+1} dx, \quad u \in H_0^1(\Omega).$$

Prove that

$$\inf_{A \in \Gamma_j} \max_{u \in A} E(u) = -\infty \quad \text{and} \quad \sup_{A \in \Gamma_j} \min_{u \in A} E(u) = +\infty.$$

**Problem 8.20** Let  $(X, \|\cdot\|)$  be a real Banach space with  $\|\cdot\|$  of class  $C^1$  on  $X \setminus \{0\}$ . Denote by  $S = \{v \in X \mid \|v\| = 1\}$  the unit sphere and let  $J \subset \mathbb{R}$  a nonempty open set. Let  $f : X \rightarrow \mathbb{R}$  be a function of class  $C^1$  on  $X \setminus \{0\}$ . We associate to  $f$  a function  $\tilde{f} : J \times X \rightarrow \mathbb{R}$  defined by

$$\tilde{f}(t, v) = f(tv),$$

where  $(t, v) \in J \times S$  and  $J \subset \mathbb{R}$  is a nonempty set. Let  $H : \mathbb{R} \times X \rightarrow \mathbb{R}$  be a  $C^1$  function satisfying

$$\langle H'_v, v \rangle \neq tH'_t \quad \text{if} \quad H(t, v) = c.$$

If  $(t, v) \in J \times X$  with  $tv \neq 0$  is a critical point of  $\tilde{f}$ , then  $u = tv$  is a critical point of  $f$ .

**Problem 8.21** Let  $(X, \|\cdot\|)$  be a real Banach space with  $\|\cdot\|$  of class  $C^1$  on  $X \setminus \{0\}$ . Denote by  $S = \{v \in X : \|v\| = 1\}$  the unit sphere and let  $J \subset \mathbb{R}$  a nonempty open set. Let  $f : X \rightarrow \mathbb{R}$  be a function of class  $C^1$  on  $X \setminus \{0\}$ . Suppose that for all  $v \in S$  the number

$$\hat{f}(v) = \max_{t \in J} f(tv)$$

is finite, and  $\hat{f}(v) > f(0)$  if  $0 \in J$ . Assume that  $\hat{f} : S \rightarrow \mathbb{R}$  is of class  $C^1$ . Then to every critical point  $v \in S$  of  $\hat{f}$ , there correspond a critical point  $u = tv$  of  $f$  with  $t \in J \setminus \{0\}$  such that  $f(u) = \hat{f}(v)$ .

## P A R T II

### Variational Principles in Geometry

## 9

# Sublinear Problems on Riemannian Manifolds

If only I had the theorems!  
Then I should find the proofs  
easily enough.

---

Bernhard Riemann  
(1826–1866)

This chapter deals with elliptic problems defined on compact Riemannian manifolds. This study is motivated by the Emden-Fowler equation arising in Mathematical Physics; after a suitable transformation, one obtains a new problem defined on the standard unit sphere  $\mathbb{S}^d$ ,  $d \geq 3$ . This problem has been extensively studied in the case when the nonlinear term is superlinear and (sub)critical at infinity. Our aim is to complete these results by considering nonlinear terms of *sublinear* type at infinity.

### 9.1 Introduction

Consider the parametrized Emden-Fowler equation

$$(EF)_\lambda \quad -\Delta v = \lambda |x|^{\alpha-2} K(x/|x|) f(|x|^{-\alpha} v), \quad x \in \mathbb{R}^{d+1} \setminus \{0\},$$

where  $f : \mathbb{R} \rightarrow \mathbb{R}$  is a continuous function,  $K$  is smooth on the  $d$ -dimensional unit sphere  $\mathbb{S}^d$ ,  $d \geq 3$ ,  $\alpha \in \mathbb{R}$ , and  $\lambda > 0$  is a parameter. The equation  $(EF)_\lambda$  has been extensively studied in the *pure superlinear* case, i.e., when  $f$  has the form  $f(s) = |s|^{p-1}s$ ,  $p > 1$ , see for instance Cotsiolis-Iliopoulos [80], Vázquez-Véron [292]. In these papers, the authors obtained existence and multiplicity of solutions for  $(EF)_\lambda$ , applying either minimization or minimax methods. Note that in the pure superlinear case the presence of the parameter  $\lambda > 0$  is not relevant due to

the rescaling technique. The solutions of  $(\text{EF})_\lambda$  are being sought in the particular form

$$v(x) = v(|x|, x/|x|) = u(r, \sigma) = r^\alpha u(\sigma), \quad (9.1)$$

where  $(r, \sigma) \in (0, \infty) \times \mathbb{S}^d$  are the spherical coordinates in  $\mathbb{R}^{d+1} \setminus \{0\}$ . This type of transformation is also used by Bidaut-Véron and Véron [40], where the asymptotics of a special form of  $(\text{EF})_\lambda$  has been studied. Throughout (9.1), the equation  $(\text{EF})_\lambda$  reduces to

$$(9.1)_\lambda \quad -\Delta_h u + \alpha(1 - \alpha - d)u = \lambda K(\sigma) f(u), \quad \sigma \in \mathbb{S}^d,$$

where  $\Delta_h$  denotes the Laplace-Beltrami operator on  $(\mathbb{S}^d, h)$  and  $h$  is the canonical metric induced from  $\mathbb{R}^{d+1}$ .

Note that when  $\alpha = -d/2$  or  $\alpha = -d/2 + 1$ , and  $f(s) = |s|^{\frac{4}{d-2}} s$ , the existence of a smooth solution  $u > 0$  of  $(9.1)_\lambda$  can be viewed as an affirmative answer to the famous Yamabe problem on  $\mathbb{S}^d$ , see Chapter 11 or Aubin [20], Cotsiolis and Iliopoulos [79], Hebey [138], and references therein. In these cases the right hand side of  $(9.1)_\lambda$  involves the Sobolev critical exponent.

The purpose of this chapter is to guarantee *multiple* solutions of  $(\text{EF})_\lambda$  for certain  $\lambda > 0$  when the nonlinearity  $f : \mathbb{R} \rightarrow \mathbb{R}$  has a sublinear growth at infinity.

## 9.2 Existence of two solutions

Since  $1 - d < \alpha < 0$  implies the coercivity of the operator  $u \mapsto -\Delta_h u + \alpha(1 - \alpha - d)u$ , the form of  $(9.1)_\lambda$  motivates the study of the following general *eigenvalue problem*, which constitutes the main objective of this section: Find  $\lambda \in (0, \infty)$  and  $u \in H_1^2(M)$  such that

$$(P_\lambda) \quad -\Delta_g u + \alpha(\sigma)u = \tilde{K}(\lambda, \sigma) f(u), \quad \sigma \in M,$$

where we assume

- (A<sub>1</sub>).  $(M, g)$  is a smooth compact  $d$ -dimensional Riemannian manifold without boundary,  $d \geq 3$ ;
- (A<sub>2</sub>).  $\alpha \in C^\infty(M)$  and  $\tilde{K} \in C^\infty((0, \infty) \times M)$  are positive functions;
- (f<sub>1</sub>).  $f : \mathbb{R} \rightarrow \mathbb{R}$  is locally Hölder continuous and sublinear at infinity, that is,

$$\lim_{|s| \rightarrow \infty} \frac{f(s)}{s} = 0. \quad (9.2)$$

A typical case when (9.2) holds is

$(f_1^{q,c})$ . There exist  $q \in (0, 1)$  and  $c > 0$  such that  $|f(s)| \leq c|s|^q$  for every  $s \in \mathbb{R}$ .

As usual,  $\Delta_g$  is the Laplace-Beltrami operator on  $(M, g)$ ; its expression in local coordinates is

$$\Delta_g u = g^{ij}(\partial_{ij} u - \Gamma_{ij}^k \partial_k u).$$

For every  $u \in C^\infty(M)$ , set

$$\|u\|_{H_\alpha^2}^2 = \int_M \langle \nabla u, \nabla u \rangle_g d\sigma_g + \int_M \alpha(\sigma) u^2 d\sigma_g,$$

where  $\langle \cdot, \cdot \rangle_g$  is the inner product on covariant tensor fields associated to  $g$ ,  $\nabla u$  is the covariant derivative of  $u$ , and  $d\sigma_g$  is the Riemannian measure. The Sobolev space  $H_\alpha^2(M)$  is defined as the completion of  $C^\infty(M)$  with respect to the norm  $\|\cdot\|_{H_\alpha^2}$ . Clearly,  $H_\alpha^2(M)$  is a Hilbert space endowed with the inner product

$$\langle u_1, u_2 \rangle_{H_\alpha^2} = \int_M \langle \nabla u_1, \nabla u_2 \rangle_g d\sigma_g + \int_M \alpha(\sigma) u_1 u_2 d\sigma_g, \quad u_1, u_2 \in H_\alpha^2(M).$$

Since  $\alpha$  is positive, the norm  $\|\cdot\|_{H_\alpha^2}$  is equivalent with the standard norm  $\|\cdot\|_{H_1^2}$ ; actually, the latter norm is nothing but  $\|\cdot\|_{H_\alpha^2}$  with  $\alpha = 1$ . Moreover, for every  $u \in H_\alpha^2(M)$ , we have

$$\min\{1, \min_M \alpha^{1/2}\} \|u\|_{H_1^2} \leq \|u\|_{H_\alpha^2} \leq \max\{1, \|\alpha\|_{L^\infty}^{1/2}\} \|u\|_{H_1^2}. \quad (9.3)$$

Note that  $H_\alpha^2(M)$  is compactly embedded into  $L^p(M)$  for every  $p \in [1, 2d/(d-2))$ ; the Sobolev embedding constant will be denoted by  $S_p > 0$ .

The presence of the parameter  $\lambda > 0$  in  $(P_\lambda)$  is indispensable. Indeed, if we consider a sublinear function at infinity which is, in addition, uniformly Lipschitz (with Lipschitz constant  $L > 0$ ), and  $\tilde{K}(\lambda, \sigma) = \lambda K(\sigma)$ , with  $K \in C^\infty(M)$  positive, one can prove that for  $0 < \lambda < \frac{1}{L} \frac{\min_M \alpha}{\max_M K} := \lambda_L$  we have only the  $u = u_\lambda = 0$  solution of  $(P_\lambda)$ , as the standard contraction principle on the Hilbert space  $H_1^2(M)$  shows. For a concrete example, let us consider the function  $f(s) = \ln(1 + s^2)$  and assume that  $\frac{K(\sigma)}{\alpha(\sigma)} = \text{const.} = \mu_0 \in (0, \infty)$ . Then, for every  $0 < \lambda < \frac{\min_M \alpha}{\mu_0 \max_M \alpha}$ , problem  $(P_\lambda)$  has only the trivial solution; however, when  $\lambda > \frac{5}{4\mu_0}$ , problem  $(P_\lambda)$  has *three* distinct constant solutions which are precisely the fixed points of the function  $s \mapsto \lambda \mu_0 \ln(1 + s^2)$ . Note that one of the solutions is the trivial one.

Let  $\lambda > 0$ . The energy functional  $\mathcal{E}_\lambda : H_1^2(M) \rightarrow \mathbb{R}$  associated with problem  $(P_\lambda)$  is

$$\mathcal{E}_\lambda(u) = \frac{1}{2} \|u\|_{H_\alpha^2}^2 - \int_M \tilde{K}(\lambda, \sigma) F(u(\sigma)) d\sigma_g, \quad (9.4)$$

where  $F(s) = \int_0^s f(\tau) d\tau$ . Due to our initial assumptions  $(A_1)$ ,  $(A_2)$  and  $(f_1)$ , the functional  $\mathcal{E}_\lambda$  is well-defined, it belongs to  $C^1(H_1^2(M), \mathbb{R})$ , and its critical points are precisely the weak (so, classical) solutions of problem  $(P_\lambda)$ . By  $(f_1)$ , for every  $\varepsilon > 0$  sufficiently small there is  $c(\varepsilon) > 0$  such that  $|f(s)| \leq \varepsilon|s| + c(\varepsilon)$  for every  $s \in \mathbb{R}$ . Thus, for every  $u \in H_1^2(M)$ , we have

$$\mathcal{E}_\lambda(u) \geq \frac{1}{2} (1 - \varepsilon \|\tilde{K}(\lambda, \cdot)\|_{L^\infty} S_2^2) \|u\|_{H_\alpha^2}^2 - c(\varepsilon) \|\tilde{K}(\lambda, \cdot)\|_{L^\infty} S_1 \|u\|_{H_\alpha^2}.$$

Therefore, the functional  $\mathcal{E}_\lambda$  is coercive and bounded from below on  $H_1^2(M)$ . Moreover, it satisfies the standard  $(PS)$ -condition.

Our first result concerns the case when  $f : \mathbb{R} \rightarrow \mathbb{R}$  fulfills the hypothesis

$(f_2)$ . There exists  $\nu_0 > 1$  such that  $\lim_{s \rightarrow 0} \frac{f(s)}{|s|^{\nu_0}} = 0$ .

**Theorem 9.1** Assume that  $f : \mathbb{R} \rightarrow \mathbb{R}$  fulfills  $(f_1)$ ,  $(f_2)$  and  $\sup_{s \in \mathbb{R}} F(s) > 0$ . Assume also that  $(A_1)$  and  $(A_2)$  are verified with  $\tilde{K}(\lambda, \sigma) = \lambda K(\sigma)$ , and  $K \in C^\infty(M)$  is positive. Then


- (i). for every  $\lambda > \lambda^* := \frac{1}{2} \frac{\|\alpha\|_{L^1}}{\|K\|_{L^1}} \left( \max_{s \neq 0} \frac{F(s)}{s^2} \right)^{-1}$  problem  $(P_\lambda)$  has at least two distinct, nontrivial solutions  $u_\lambda^1, u_\lambda^2 \in H_1^2(M)$ , where  $u_\lambda^1$  is the global minimum of the energy functional associated with  $(P_\lambda)$ ;
- (ii). if  $(f_1^{q,c})$  holds then  $\|u_\lambda^1\|_{H_1^2} = o(\lambda^{\frac{1}{1-\tau}})$  for every  $r \in (q, 1)$ , but  $\|u_\lambda^1\|_{H_1^2} \neq O(\lambda^{\frac{1}{1-\mu}})$  for any  $\mu \in (1, (d+2)/(d-2))$  as  $\lambda \rightarrow \infty$ .

A direct consequence of Theorem 9.1 applied for  $(9.1)_\lambda$  is the following

**Theorem 9.2** Assume that  $1-d < \alpha < 0$ . Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be a function as in Theorem 9.1 and  $K \in C^\infty(\mathbb{S}^d)$  positive. Then, for every  $\lambda > \lambda^*$  problem  $(EF)_\lambda$  has at least two distinct, nontrivial solutions.

**Example 9.1** Let  $f(s) = \ln(1 + s^2)$ , see Figure 9.1. One can apply Theorems 9.1–9.2.

*Proof of Theorem 9.1.* Due to (9.2) and  $(f_2)$ , the term  $F(s)/s^2$  tends to 0 as  $|s| \rightarrow \infty$  and  $s \rightarrow 0$ , respectively. Since there exists  $s_0 \in \mathbb{R}$  such that  $F(s_0) > 0$ , the number  $\lambda^*$  is well-defined. Now, fix a number  $s^* \in \mathbb{R} \setminus \{0\}$  such that  $\frac{F(s^*)}{(s^*)^2} = \max_{s \neq 0} \frac{F(s)}{s^2}$ . Therefore,  $\lambda^* = \frac{1}{2} \frac{\|\alpha\|_{L^1}}{\|K\|_{L^1}} \frac{(s^*)^2}{F(s^*)}$ .

Figure 9.1 The function  $f(s) = \ln(1 + s^2)$ .

Fix  $\lambda > \lambda^*$ . Let us first consider the constant function  $u_{s^*}(\sigma) = s^*$ ,  $\sigma \in M$ . One has

$$\begin{aligned} \mathcal{E}_\lambda(u_{s^*}) &= \frac{1}{2} \|u_{s^*}\|_{H_\alpha^2}^2 - \lambda \int_M K(\sigma) F(u_{s^*}(\sigma)) d\sigma_g \\ &= \frac{1}{2} \|\alpha\|_{L^1} (s^*)^2 - \lambda \|K\|_{L^1} F(s^*) = (\lambda^* - \lambda) \|K\|_{L^1} F(s^*) \\ &< 0. \end{aligned}$$

Thus,  $\inf_{H_1^2(M)} \mathcal{E}_\lambda \leq \mathcal{E}_\lambda(u_{s^*}) < 0$ . Since  $\mathcal{E}_\lambda$  verifies the  $(PS)$ -condition and it is bounded from below, one can find  $u_\lambda^1 \in H_1^2(M)$  such that  $\mathcal{E}_\lambda(u_\lambda^1) = \inf_{H_1^2(M)} \mathcal{E}_\lambda$ , see Theorem 1.4. Therefore,  $u_\lambda^1 \in H_1^2(M)$  is the first solution of  $(P_\lambda)$  and  $u_\lambda^1 \neq 0$ , since  $\mathcal{E}_\lambda(0) = 0$ .

Now, we prove that for every  $\lambda > \lambda^*$  the functional  $\mathcal{E}_\lambda$  has the standard Mountain Pass geometry. First, due to  $(f_1)$  and  $(f_2)$ , one can fix two numbers  $C > 0$  and  $1 < \nu < \frac{d+2}{d-2}$  such that

$$|F(s)| \leq C|s|^{\nu+1}, \quad s \in \mathbb{R}.$$

Since  $\nu + 1 < \frac{2d}{d-2}$  (thus  $H_\alpha^2(M) \hookrightarrow L^{\nu+1}(M)$  is continuous), one has

$$\mathcal{E}_\lambda(u) \geq \frac{1}{2} \|u\|_{H_\alpha^2}^2 - \lambda C \|K\|_{L^\infty} S_{\nu+1}^{\nu+1} \|u\|_{H_\alpha^2}^{\nu+1}, \quad u \in H_1^2(M). \quad (9.5)$$

Let us take  $\rho_\lambda > 0$  so small such that

$$\rho_\lambda < \min \left\{ (2\lambda C \|K\|_{L^\infty} S_{\nu+1}^{\nu+1})^{\frac{1}{1-\nu}} \min\{1, \|\alpha\|^{-1/2}\}, |s^*| (\text{Vol}_g(M))^{1/2} \right\}.$$

Consequently, by (9.5), for every  $u \in H_1^2(M)$  complying with  $\|u\|_{H_1^2} = \rho_\lambda$ , we have

$$\begin{aligned} \mathcal{E}_\lambda(u) &\geq \left( \frac{1}{2} - \lambda C \|K\|_{L^\infty} S_{\nu+1}^{\nu+1} \|u\|_{H_\alpha^2}^{\nu-1} \right) \|u\|_{H_\alpha^2}^2 \\ &\geq \left( \frac{1}{2} - \lambda C \|K\|_{L^\infty} S_{\nu+1}^{\nu+1} \max\{1, \|\alpha\|_{L^\infty}^{(\nu-1)/2}\} \rho_\lambda^{\nu-1} \right) \min\{1, \min_M \alpha\} \rho_\lambda^2 \\ &\equiv \eta(\rho_\lambda) > 0. \end{aligned}$$

By construction, one has  $\|u_{s^*}\|_{H_1^2} = |s^*|(\text{Vol}_g(M))^{1/2} > \rho_\lambda$  and from above  $\mathcal{E}_\lambda(u_{s^*}) < 0 = \mathcal{E}_\lambda(0)$ . Since  $\mathcal{E}_\lambda$  satisfies the  $(PS)$ -condition, one can apply the Mountain Pass Theorem, see Theorem 1.7. Thus, there exists an element  $u_\lambda^2 \in H_1^2(M)$  such that  $\mathcal{E}'_\lambda(u_\lambda^2) = 0$  and  $\mathcal{E}_\lambda(u_\lambda^2) \geq \eta(\rho_\lambda) > 0$ . In particular,  $u_\lambda^2 \neq 0$ , and the elements  $u_\lambda^1$  and  $u_\lambda^2$  are distinct. This ends (i).

Now, we assume that  $(f_1^{c,q})$  holds. Since  $\mathcal{E}_\lambda(u_\lambda^1) < 0$ , then

$$\frac{1}{2} \|u_\lambda^1\|_{H_\alpha^2}^2 - \lambda \frac{c}{q+1} \|K\|_{L^\infty} S_2^{q+1} (\text{Vol}_g(M))^{(1-q)/2} \|u_\lambda^1\|_{H_\alpha^2}^{q+1} < 0.$$

In particular,  $\|u_\lambda^1\|_{H_1^2} = O(\lambda^{\frac{1}{1-q}})$  as  $\lambda \rightarrow \infty$ . Therefore, for every  $r \in (q, 1)$ , one has  $\|u_\lambda^1\|_{H_1^2} = o(\lambda^{\frac{1}{1-r}})$  as  $\lambda \rightarrow \infty$ .

Let us assume that  $\|u_\lambda^1\|_{H_1^2} = O(\lambda^{\frac{1}{1-\mu}})$  for some  $\mu \in (1, (d+2)/(d-2))$  as  $\lambda \rightarrow \infty$ . Consequently,  $\|u_\lambda^1\|_{H_\alpha^2} \rightarrow 0$  as  $\lambda \rightarrow \infty$ . On the other hand, since  $\mathcal{E}_\lambda(u_\lambda^1) \leq (\lambda^* - \lambda) \|K\|_{L^1} F(s^*)$ , then  $\mathcal{E}_\lambda(u_\lambda^1) \rightarrow -\infty$  as  $\lambda \rightarrow \infty$ . On account of (9.5), we have

$$\left( \frac{1}{2} - \lambda C \|K\|_{L^\infty} S_{\mu+1}^{\mu+1} \|u_\lambda^1\|_{H_\alpha^2}^{\mu-1} \right) \|u_\lambda^1\|_{H_\alpha^2}^2 \rightarrow -\infty$$

as  $\lambda \rightarrow \infty$ . Therefore, the expression  $\lambda \|u_\lambda^1\|_{H_\alpha^2}^{\mu-1}$  necessarily tends to  $\infty$  as  $\lambda \rightarrow \infty$ . But this contradicts the initial assumption. This completes (ii).  $\square$

*Proof of Theorem 9.2.* Let us choose  $(M, g) = (\mathbb{S}^d, h)$ , and  $\alpha(\sigma) := \alpha(1 - \alpha - d)$  for every  $\sigma \in \mathbb{S}^d$  in Theorem 9.1. Thus, for every  $\lambda > \lambda^*$ , problem  $(9.1)_\lambda$  has at least two distinct, nontrivial solutions  $u_\lambda^1, u_\lambda^2 \in H_1^2(\mathbb{S}^d)$ . On account of (9.1), the elements  $v_\lambda^i(x) = |x|^\alpha u_\lambda^i(x/|x|)$ ,  $i \in \{1, 2\}$ , are solutions of  $(EF)_\lambda$ .  $\square$

Now, instead of  $(f_2)$  we consider a weaker assumption, namely:

$$(f'_2). \quad \lim_{s \rightarrow 0} \frac{f(s)}{s} = 0.$$




Figure 9.2 The function from Example 9.2.

The following result shows that  $(f'_2)$  is still enough to prove a similar multiplicity result as Theorem 9.1, but we lose the precise location of the eigenvalues. More precisely, we have

**Theorem 9.3** *Assume that  $f : \mathbb{R} \rightarrow \mathbb{R}$  fulfills  $(f_1)$ ,  $(f'_2)$  and  $\sup_{s \in \mathbb{R}} F(s) > 0$ . Assume also that  $(A_1)$  and  $(A_2)$  are verified with  $\tilde{K}(\lambda, \sigma) = \lambda K(\sigma)$ .*

*Then there exist a nonempty open interval  $\Lambda \subset (0, \infty)$  and a number  $\gamma > 0$  such that for every  $\lambda \in \Lambda$  problem  $(P_\lambda)$  has at least two distinct, nontrivial solutions  $u_\lambda^1, u_\lambda^2 \in H_1^2(M)$  and  $\|u_\lambda^i\|_{H_1^2} < \gamma$ ,  $i \in \{1, 2\}$ .*

Similarly as in Theorem 9.2, we have

**Theorem 9.4** *Assume that  $1 - d < \alpha < 0$ . Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be a function as in Theorem 9.3 and  $K \in C^\infty(\mathbb{S}^d)$  positive. Then, there exists a nonempty open interval  $\Lambda \subset (0, \infty)$  such that for every  $\lambda \in \Lambda$  problem  $(EF)_\lambda$  has at least two distinct, nontrivial solutions.*

**Example 9.2** Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be defined by  $f(s) = 0$  for  $s \leq 0$ ,  $f(s) = \frac{s}{\ln s}$  for  $s \in (0, e^{-1}]$  and  $f(s) = -e^{-1}$  for  $s > e^{-1}$ , see Figure 9.2. Note that  $f$  satisfies  $(f'_2)$  but not  $(f_2)$  for any  $\nu_0 > 1$ . Therefore, one can apply Theorems 9.3 and 9.4 (but not Theorems 9.1 and 9.2).

In order to prove Theorems 9.3 and 9.4, we introduce the functionals  $\mathcal{N}, \mathcal{F} : H_1^2(M) \rightarrow \mathbb{R}$  defined by

$$\mathcal{N}(u) = \frac{1}{2} \|u\|_{H_1^2}^2 \quad \text{and} \quad \mathcal{F}(u) = \int_M K(\sigma) F(u(\sigma)) d\sigma_g, \quad u \in H_1^2(M). \quad (9.6)$$

**Proposition 9.1**  $\lim_{\rho \rightarrow 0^+} \frac{\sup\{\mathcal{F}(u) : \mathcal{N}(u) < \rho\}}{\rho} = 0$ .

*Proof.* Due to  $(f'_2)$ , for an arbitrarily small  $\varepsilon > 0$  there exists  $\delta(\varepsilon) > 0$  such that  $|f(s)| < \varepsilon(2\|K\|_{L^\infty}S_2^2)^{-1}|s|$  for every  $|s| < \delta(\varepsilon)$ . On account of  $(f_1)$ , one may fix  $1 < \nu < \frac{d+2}{d-2}$  and  $c(\varepsilon) > 0$  such that  $|f(s)| < c(\varepsilon)|s|^\nu$  for every  $|s| \geq \delta(\varepsilon)$ . Combining these two facts, after an integration, we obtain

$$|F(s)| \leq \varepsilon(4\|K\|_{L^\infty}S_2^2)^{-1}s^2 + c(\varepsilon)(\nu+1)^{-1}|s|^{\nu+1} \quad \text{for every } s \in \mathbb{R}.$$

Fix a  $\rho > 0$  and any element  $u \in H_\alpha^2(M)$  complying with  $\mathcal{N}(u) < \rho$ . Due to the above estimation, we have

$$\begin{aligned} \mathcal{F}(u) &\leq \frac{\varepsilon}{4}\|u\|_{H_\alpha^2}^2 + \frac{c(\varepsilon)}{\nu+1}\|K\|_{L^\infty}S_{\nu+1}^{\nu+1}\|u\|_{H_\alpha^2}^{\nu+1} \\ &< \frac{\varepsilon}{2}\rho + \frac{c(\varepsilon)}{\nu+1}\|K\|_{L^\infty}S_{\nu+1}^{\nu+1}(2\rho)^{\frac{\nu+1}{2}} = \frac{\varepsilon}{2}\rho + c'(\varepsilon)\rho^{\frac{\nu+1}{2}}. \end{aligned}$$

Thus there exists  $\rho(\varepsilon) > 0$  such that for every  $0 < \rho < \rho(\varepsilon)$ , we have

$$0 \leq \frac{\sup\{\mathcal{F}(u) : \mathcal{N}(u) < \rho\}}{\rho} \leq \frac{\varepsilon}{2} + c'(\varepsilon)\rho^{\frac{\nu-1}{2}} < \varepsilon,$$

which completes the proof.  $\square$

*Proof of Theorem 9.3.* Let  $X = H_1^2(M)$ , and the functionals  $\mathcal{N}, \mathcal{F}$  defined in (9.6). Note that  $\mathcal{E}_\lambda = \mathcal{N} - \lambda\mathcal{F}$ . We know already that for every  $\lambda > 0$  the functional  $\mathcal{E}_\lambda = \mathcal{N} - \lambda\mathcal{F}$  is coercive and satisfies the (PS)-condition. Moreover, since the embedding  $H_1^2(M) \hookrightarrow L^p(M)$  is compact,  $1 \leq p < 2d/(d-2)$ , the functional  $\mathcal{F}$  is sequentially weakly continuous; thus,  $\mathcal{E}_\lambda$  is sequentially weakly lower semicontinuous.

Due to (9.2) and  $(f'_2)$ , the term  $F(s)/s^2$  tends to 0 as  $|s| \rightarrow \infty$  and  $s \rightarrow 0$ , respectively. Since there exists  $s_0 \in \mathbb{R}$  such that  $F(s_0) > 0$ , we may fix a number  $s^* \neq 0$  such that  $\frac{F(s^*)}{(s^*)^2} = \max_{s \neq 0} \frac{F(s)}{s^2}$ . Therefore,  $\lambda^*$  appearing in Theorem 9.1 is well-defined and  $\lambda^* = \frac{1}{2} \frac{\|\alpha\|_{L^1} (s^*)^2}{\|K\|_{L^1} F(s^*)}$ .

Now, let us choose  $u_0 = 0$ , and  $u_1(\sigma) = u_{s^*}(\sigma) = s^*$  for every  $\sigma \in M$ . Fixing  $\varepsilon \in (0, 1)$ , due to Proposition 9.1, one may choose  $\rho > 0$  such that

$$\frac{\sup\{\mathcal{F}(u) : \mathcal{N}(u) < \rho\}}{\rho} < \frac{\varepsilon}{\lambda^*}; \quad (9.7)$$

$$\rho < \frac{1}{2}(s^*)^2\|\alpha\|_{L^1}.$$

Note that  $\frac{\varepsilon}{\lambda^*} < \frac{1}{\lambda^*} = \frac{\mathcal{F}(u_1)}{\mathcal{N}(u_1)}$  and  $\frac{1}{2}(s^*)^2 \|\alpha\|_{L^1} = \mathcal{N}(u_1)$ . Therefore, by choosing

$$\bar{a} = \frac{1 + \varepsilon}{\frac{\mathcal{F}(u_1)}{\mathcal{N}(u_1)} - \frac{\sup\{\mathcal{F}(u) : \mathcal{N}(u) < \rho\}}{\rho}}, \tag{9.8}$$

all the hypotheses of Theorem 1.16 are verified. Consequently, there is a nonempty open interval  $\Lambda \subset [0, \bar{a}]$  and a number  $\gamma > 0$  such that for every  $\lambda \in \Lambda$ , the functional  $\mathcal{E}_\lambda$  has at least three distinct critical points in  $H_1^2(M)$  having  $\|\cdot\|_{H_1^2}$ -norm less than  $\gamma$ . This ends the proof of Theorem 9.3.  $\square$

*Proof of Theorem 9.4.* Similar to Theorem 9.2.  $\square$

### 9.3 Existence of many global minima

In order to obtain a new kind of multiplicity result concerning  $(P_\lambda)$  (specially,  $(9.1)_\lambda$  and  $(EF)_\lambda$ ), we require:

- ( $f_3$ ). *There exists  $\mu_0 \in (0, \infty)$  such that the global minima of the function  $s \mapsto \tilde{F}_{\mu_0}(s) := \frac{1}{2}s^2 - \mu_0 F(s)$  has at least  $m \geq 2$  connected components.*

Note that ( $f_3$ ) implies that the function  $s \mapsto \tilde{F}_{\mu_0}(s)$  has at least  $m - 1$  local maxima. Thus, the function  $s \mapsto \mu_0 f(s)$  has at least  $2m - 1$  fixed points. In particular, if for some  $\lambda > 0$  one has  $\frac{\tilde{K}(\lambda, \sigma)}{\alpha(\sigma)} = \mu_0$  for every  $\sigma \in M$ , then problem  $(P_\lambda)$  has at least  $2m - 1 \geq 3$  constant solutions. On the other hand, the following general result can be shown.

**Theorem 9.5** *Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be a function which fulfills ( $f_1$ ) and ( $f_3$ ). Assume that  $(A_1)$  and  $(A_2)$  are verified with  $\tilde{K}(\lambda, \sigma) = \lambda K(\sigma) + \mu_0 \alpha(\sigma)$ , and  $K \in C^\infty(M)$  is positive. Then*

- (i). *for every  $\eta > \max\{0, \|\alpha\|_{L^1} \min_{s \in \mathbb{R}} \tilde{F}_{\mu_0}(s)\}$  there exists a number  $\tilde{\lambda}_\eta > 0$  such that for every  $\lambda \in (0, \tilde{\lambda}_\eta)$  problem  $(P_\lambda)$  has at least  $m + 1$  solutions  $u_\lambda^{1, \eta}, \dots, u_\lambda^{m+1, \eta} \in H_1^2(M)$ ;*
- (ii). *if  $(f_1^{q,c})$  holds then for each  $\lambda \in (0, \tilde{\lambda}_\eta)$  there is a set  $I_\lambda \subset \{1, \dots, m + 1\}$  with  $\text{card}(I_\lambda) = m$  such that*

$$\|u_\lambda^{i, \eta}\|_{H_1^2} < \frac{s_{\eta, q, c}}{\min\{1, \min_M \alpha^{1/2}\}}, \quad i \in I_\lambda,$$




Figure 9.3 The function  $f(s) = \min\{s_+ - \sin(\pi s_+), 2(m-1)\}$  from Example 9.3 for  $m = 5$ .

where  $s_{\eta,q,c} > 0$  is the greatest solution of the equation

$$\frac{1}{2}s^2 - \frac{\mu_0 c \|\alpha\|_{L^1}^{(1-q)/2}}{q+1} s^{q+1} - \eta = 0, \quad s > 0.$$

A consequence of Theorem 9.5 in the context of  $(\text{EF})_\lambda$  reads as follows.

**Theorem 9.6** *Assume that  $1-d < \alpha < 0$ . Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be a function as in Theorem 9.5 and  $K \in C^\infty(\mathbb{S}^d)$  be a positive function. Then, there exists  $\lambda_0 > 0$  such that for every  $\lambda \in (0, \lambda_0)$  problem*

$$-\Delta v = |x|^{\alpha-2} [\lambda K(x/|x|) + \mu_0 \alpha (1-\alpha-d)] f(|x|^{-\alpha} v), \quad x \in \mathbb{R}^{d+1} \setminus \{0\}$$

has at least  $m+1$  solutions.

**Example 9.3** Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be the function defined by

$$f(s) = \min\{s_+ - \sin(\pi s_+), 2(m-1)\},$$

where  $m \in \mathbb{N} \setminus \{1\}$  is fixed and we use the notation  $s_+ = \max\{s, 0\}$ . Clearly,  $(f_1)$  is verified, while for  $\mu_0 = 1$ , the assumption  $(f_3)$  is also fulfilled. Indeed, the function  $s \mapsto \tilde{F}_1(s)$  has precisely  $m$  global minima; they are  $0, 2, \dots, 2(m-1)$ . Moreover,  $\min_{s \in \mathbb{R}} \tilde{F}_1(s) = 0$ . Therefore, one can apply Theorems 9.5 and 9.6.

In order to prove Theorems 9.5 and 9.6, throughout relation (9.6), we define the functional  $\mathcal{N}_{\mu_0} : H_1^2(M) \rightarrow \mathbb{R}$  by

$$\mathcal{N}_{\mu_0}(u) = \mathcal{N}(u) - \mu_0 \int_M \alpha(\sigma) F(u(\sigma)) d\sigma_g, \quad u \in H_1^2(M).$$

**Proposition 9.2** *The set of all global minima of the functional  $\mathcal{N}_{\mu_0}$  has at least  $m$  connected components in the weak topology on  $H_1^2(M)$ .*

*Proof.* First, for every  $u \in H_1^2(M)$  we have

$$\begin{aligned} \mathcal{N}_{\mu_0}(u) &= \frac{1}{2} \|u\|_{H_\alpha^2}^2 - \mu_0 \int_M \alpha(\sigma) F(u(\sigma)) d\sigma_g \\ &= \frac{1}{2} \int_M |\nabla u|_g^2 d\sigma_g + \int_M \alpha(\sigma) \tilde{F}_{\mu_0}(u(\sigma)) d\sigma_g \\ &\geq \|\alpha\|_{L^1} \inf_{s \in \mathbb{R}} \tilde{F}_{\mu_0}(s). \end{aligned}$$

Moreover, if we consider  $u(\sigma) = u_{\tilde{s}}(\sigma) = \tilde{s}$  for a.e.  $\sigma \in M$ , where  $\tilde{s} \in \mathbb{R}$  is a minimum point of the function  $s \mapsto \tilde{F}_{\mu_0}(s)$ , then we have equality in the previous estimation. Thus,

$$\inf_{u \in H_1^2(M)} \mathcal{N}_{\mu_0}(u) = \|\alpha\|_{L^1} \inf_{s \in \mathbb{R}} \tilde{F}_{\mu_0}(s).$$

Moreover, if  $u \in H_1^2(M)$  is not a constant function, then  $|\nabla u|_g^2 = g^{ij} \partial_i u \partial_j u > 0$  on a positive measured set of the manifold  $M$ . In this case, we have

$$\mathcal{N}_{\mu_0}(u) = \frac{1}{2} \int_M |\nabla u|_g^2 d\sigma_g + \int_M \alpha(\sigma) \tilde{F}_{\mu_0}(u(\sigma)) d\sigma_g > \|\alpha\|_{L^1} \inf_{s \in \mathbb{R}} \tilde{F}_{\mu_0}(s).$$

Consequently, between the sets

$$\text{Min}(\mathcal{N}_{\mu_0}) = \{u \in H_1^2(M) : \mathcal{N}_{\mu_0}(u) = \inf_{u \in H_1^2(M)} \mathcal{N}_{\mu_0}(u)\}$$

and

$$\text{Min}(\tilde{F}_{\mu_0}) = \{s \in \mathbb{R} : \tilde{F}_{\mu_0}(s) = \inf_{s \in \mathbb{R}} \tilde{F}_{\mu_0}(s)\}$$

there is a one-to-one correspondence. Indeed, let  $\theta$  be the function that associates to every number  $t \in \mathbb{R}$  the equivalence class of those functions which are a.e. equal to  $t$  in the whole manifold  $M$ . Then  $\theta : \text{Min}(\tilde{F}_{\mu_0}) \rightarrow \text{Min}(\mathcal{N}_{\mu_0})$  is actually a homeomorphism between  $\text{Min}(\tilde{F}_{\mu_0})$  and  $\text{Min}(\mathcal{N}_{\mu_0})$ , where the set  $\text{Min}(\mathcal{N}_{\mu_0})$  is considered with the relativization of the weak topology on  $H_1^2(M)$ . On account of the hypothesis  $(f_3)$ , the set  $\text{Min}(\tilde{F}_{\mu_0})$  contains at least  $m \geq 2$  connected components. Therefore, the same is true for the set  $\text{Min}(\mathcal{N}_{\mu_0})$ , which completes the proof.  $\square$

*Proof of Theorem 9.5.* Let us choose  $X = H_1^2(M)$ ,  $\mathcal{N} = \mathcal{N}_{\mu_0}$  and  $\mathcal{G} = -\mathcal{F}$  in Theorem 1.22. Due to Proposition 9.2 and to basic properties of the functions  $\mathcal{N}_{\mu_0}$ ,  $\mathcal{F}$ , all the hypotheses of Theorem 1.22 are satisfied.

Then, for every

$$\eta > \max\{0, \|\alpha\|_{L^1} \min_{s \in \mathbb{R}} \tilde{F}_{\mu_0}(s)\} \left( \geq \inf_{u \in H_1^2(M)} \mathcal{N}_{\mu_0}(u) \right)$$

there is a number  $\tilde{\lambda}_\eta > 0$  such that for every  $\lambda \in (0, \tilde{\lambda}_\eta)$  the function  $\mathcal{N}_{\mu_0} - \lambda \mathcal{F}$  has at least  $m + 1$  critical points; let us denote them by  $u_\lambda^{1,\eta}, \dots, u_\lambda^{m+1,\eta} \in H_1^2(M)$ . Clearly, they are solutions of problem  $(P_\lambda)$ , which concludes (i).

We know in addition that  $m$  elements from  $u_\lambda^{1,\eta}, \dots, u_\lambda^{m+1,\eta}$  belong to the set  $\mathcal{N}_{\mu_0}^{-1}((-\infty, \eta))$ . Let  $\tilde{u}$  be such an element, i.e.,

$$\mathcal{N}_{\mu_0}(\tilde{u}) = \frac{1}{2} \|\tilde{u}\|_{H_\alpha^2}^2 - \mu_0 \int_M \alpha(\sigma) F(\tilde{u}(\sigma)) d\sigma_g < \eta. \tag{9.9}$$

Assume that  $(f_1^{q,c})$  holds. Then  $|F(s)| \leq \frac{c}{q+1} |s|^{q+1}$  for every  $s \in \mathbb{R}$ . By using the Hölder inequality, one has

$$\int_M \alpha(\sigma) |\tilde{u}(\sigma)|^{q+1} d\sigma_g \leq \|\alpha\|_{L^1}^{(1-q)/2} \|\tilde{u}\|_{H_\alpha^2}^{q+1}. \tag{9.10}$$

Since  $\eta > 0$ , the equation

$$\frac{1}{2} s^2 - \frac{\mu_0 c \|\alpha\|_{L^1}^{(1-q)/2}}{q+1} |s|^{q+1} - \eta = 0, \tag{9.11}$$

has always a positive solution. On account of (9.9) and (9.10), the number  $\|\tilde{u}\|_{H_\alpha^2}$  is less then the greatest solution  $s_{\eta,q,c} > 0$  of the equation (9.11). It remains to apply (9.3), which concludes (ii).  $\square$

*Proof of Theorem 9.6.* It follows directly by Theorem 9.5.  $\square$

### 9.4 Historical notes and comments

*A. Historical notes.* An important class of elliptic problems defined on compact manifolds arises from the Emden-Fowler equation, studied for superlinear nonlinearities via minimization or minimax methods, see Cotsiolis and Iliopoulos [80], Vázquez and Véron [292], Bidaut-Véron and Véron [40].

In this chapter we presented some contributions related to Emden-Fowler equations when the nonlinearity has a sublinear growth at infinity.

*B. Comments.* The main results of the present chapter are based on the paper of Kristály and Rădulescu [178].

As we mentioned at the beginning of Section 9.2, when the nonlinearity  $f$  is a uniformly Lipschitz function (with Lipschitz constant  $L > 0$ ), we have extra information on the eigenvalues:

- (a) problem  $(P_\lambda)$  has only the trivial solution whenever  $\lambda \in (0, \lambda_L)$ ;
- (b) problem  $(P_\lambda)$  has at least two nontrivial solutions whenever  $\lambda > \lambda^*$ . Clearly, we have  $\lambda^* \geq \lambda_L$ , and usually these two numbers do not coincide. (For instance, if  $f(s) = \ln(1 + s^2)$ ,  $K(\sigma) = \alpha(\sigma) = \text{const.}$ , then  $\lambda_L = 1$  while  $\lambda^* \approx 1.32$ .) Unfortunately, we have no any precise result when the parameter belongs to the 'gap-interval'  $[\lambda_L, \lambda^*]$ .

# 10

## Asymptotically Critical Problems on Spheres

Facts are the air of scientists.  
Without them you can never  
fly.

---

Linus Pauling (1901–1994)

In this chapter we study a nonlinear elliptic problem defined on the unit sphere  $(\mathbb{S}^d, h)$ , involving an asymptotically critical oscillatory term and the Laplace-Beltrami operator  $\Delta_h u = \operatorname{div}_h(\nabla u)$  where  $h$  is the canonical metric induced from  $\mathbb{R}^{d+1}$ . By a group-theoretical argument, the existence of infinitely many sign-changing solutions for the studied problem are guaranteed, giving also a lower estimate of the number of those sequences of solutions whose elements in different sequences are mutually symmetrically distinct.

### 10.1 Introduction

We consider the nonlinear elliptic problem

$$-\Delta_h u = f(u) \quad \text{on } \mathbb{S}^d, \quad (\text{P})$$

where  $\Delta_h u = \operatorname{div}_h(\nabla u)$  denotes the Laplace-Beltrami operator acting on  $u : \mathbb{S}^d \rightarrow \mathbb{R}$ ,  $(\mathbb{S}^d, h)$  is the unit sphere,  $h$  being the canonical metric induced from  $\mathbb{R}^{d+1}$ .

Consider a continuous nonlinearity  $f : \mathbb{R} \rightarrow \mathbb{R}$  that satisfies  $(f_1^L)$  and  $(f_2^L)$  from Section 7.1, where  $L$  denotes  $0^+$  or  $+\infty$ . As we pointed out before, these assumptions imply that  $f$  has an *oscillatory behaviour at  $L$* . In particular, a whole sequence of distinct, *constant* solutions for (P) appears as zeros of the function  $s \mapsto f(s)$ ,  $s > 0$ .

The purpose of this chapter is to investigate the existence of *non-constant* solutions for (P) under the assumptions  $(f_1^L)$  and  $(f_2^L)$ . This problem will be achieved by constructing sign-changing solutions for (P). We prove two multiplicity results corresponding to  $L = 0^+$  and  $L = +\infty$ , respectively; the 'piquancy' is that not only infinitely many sign-changing solutions for (P) are guaranteed but we also give a lower estimate of the number of those sequences of solutions for (P) whose elements in different sequences are mutually symmetrically distinct.

In order to handle this problem, solutions for (P) are being sought in the standard Sobolev space  $H_1^2(\mathbb{S}^d)$ . We say that  $u \in H_1^2(\mathbb{S}^d)$  is a weak solution for (P) if

$$\int_{\mathbb{S}^d} \langle \nabla u, \nabla v \rangle_h d\sigma_h = \int_{\mathbb{S}^d} f(u) v d\sigma_h \quad \text{for all } v \in H_1^2(\mathbb{S}^d).$$

Since we are interested in the existence of infinitely many sign-changing solutions, it seems some kind of symmetry hypothesis on the nonlinearity  $f$  is indispensable; namely, we assume that  $f$  is odd in an arbitrarily small neighborhood of the origin whenever  $L = 0^+$ , and  $f$  is odd on the whole  $\mathbb{R}$  whenever  $L = +\infty$ . In the case  $L = 0^+$  no further assumption on  $f$  is needed at infinity (neither symmetry nor growth of  $f$ ; in particular,  $f$  may have even a *supercritical* growth). However, when  $L = +\infty$ , we have to control the growth of  $f$ ; we assume  $f(s) = O(s^{\frac{d+2}{d-2}})$  as  $s \rightarrow \infty$ , that is,  $f$  has an *asymptotically critical growth at infinity*. In both cases ( $L = 0^+$  and  $L = +\infty$ ), the energy functional  $\mathcal{E} : H_1^2(\mathbb{S}^d) \rightarrow \mathbb{R}$  associated with (P) is well-defined, which is the key tool in order to achieve our results.

The first task is to construct certain subspaces of  $H_1^2(\mathbb{S}^d)$  containing invariant functions under special actions defined by means of carefully chosen subgroups of the orthogonal group  $O(d+1)$ . A particular form of this construction has been first exploited by Ding [91]. In our case, every nontrivial element from these subspaces of  $H_1^2(\mathbb{S}^d)$  changes the sign. The main feature of these subspaces of  $H_1^2(\mathbb{S}^d)$  is based on the symmetry properties of their elements: no nontrivial element from one subspace can belong to another subspace, i.e., elements from distinct subspaces are distinguished by their symmetries. Consequently, guaranteeing nontrivial solutions for (P) in distinct subspaces of  $H_1^2(\mathbb{S}^d)$  of the above type, these elements cannot be compared with each other. In the next section we show by an explicit construction that the minimal number of these subspaces of  $H_1^2(\mathbb{S}^d)$  is  $s_d = [d/2] + (-1)^{d+1} - 1$ . Here,  $[ \cdot ]$  denotes the integer function.

## 10.2 Group-theoretical argument

Let  $d \geq 5$  and  $s_d = [d/2] + (-1)^{d+1} - 1$ . For every  $i \in \{1, \dots, s_d\}$ , we define

$$G_{d,i} = \begin{cases} O(i+1) \times O(d-2i-1) \times O(i+1), & \text{if } i \neq \frac{d-1}{2}, \\ O(\frac{d+1}{2}) \times O(\frac{d+1}{2}), & \text{if } i = \frac{d-1}{2}. \end{cases}$$

Let us denote by  $\langle G_{d,i}; G_{d,j} \rangle$  the group generated by  $G_{d,i}$  and  $G_{d,j}$ . The key result of this section is

**Proposition 10.1** *For every  $i, j \in \{1, \dots, s_d\}$  with  $i \neq j$ , the group  $\langle G_{d,i}; G_{d,j} \rangle$  acts transitively on  $\mathbb{S}^d$ .*

*Proof.* Without losing the generality, we may assume that  $i < j$ . The proof is divided into three steps. For abbreviation, we introduce the notation  $0_k = (0, \dots, 0) \in \mathbb{R}^k$ ,  $k \in \{1, \dots, d+1\}$ .

*Step 1.* The group  $\langle G_{d,i}; G_{d,j} \rangle$  acts transitively on  $\mathbb{S}^{d-j-1} \times \{0_{j+1}\}$ . When  $j = \frac{d-1}{2}$ , the proof is trivial since  $O(\frac{d+1}{2})$  acts transitively on  $\mathbb{S}^{\frac{d-1}{2}}$ . Assume so that  $j \neq \frac{d-1}{2}$ . We show that for every  $\sigma = (\sigma_1, \sigma_2, \sigma_3) \in \mathbb{S}^{d-j-1}$  with  $\sigma_1 \in \mathbb{R}^{i+1}$ ,  $\sigma_2 \in \mathbb{R}^{j-i}$ ,  $\sigma_3 \in \mathbb{R}^{d-2j-1}$ , and  $\omega \in \mathbb{S}^j$  fixed arbitrarily, there exists  $g_{ij} \in \langle G_{d,i}; G_{d,j} \rangle$  such that

$$g_{ij}(\omega, 0_{d-j}) = (\sigma, 0_{j+1}). \quad (10.1)$$

Since  $O(j+1)$  acts transitively on  $\mathbb{S}^j$ , for every  $\tilde{\sigma}_2 \in \mathbb{R}^{j-i}$  with the property that  $(\sigma_1, \tilde{\sigma}_2) \in \mathbb{S}^j$ , there exists an element  $g_j \in O(j+1)$  such that

$$g_j \omega = (\sigma_1, \tilde{\sigma}_2). \quad (10.2)$$

Note that  $|\sigma_1|^2 + |\tilde{\sigma}_2|^2 = 1$  and  $|\sigma_1|^2 + |\sigma_2|^2 + |\sigma_3|^2 = 1$ ; so  $|\tilde{\sigma}_2|^2 = |\sigma_2|^2 + |\sigma_3|^2$ .

If  $\tilde{\sigma}_2 = 0_{j-i}$  then  $\sigma_2 = 0_{j-i}$  and  $\sigma_3 = 0_{d-2j-1}$ ; thus,  $\sigma = (\sigma_1, 0_{d-j-i-1})$ . Let  $g_{ij} := g_j \times id_{\mathbb{R}^{d-j}} \in G_{d,j}$ . Then, due to (10.2), we have

$$g_{ij}(\omega, 0_{d-j}) = (g_j \omega, 0_{d-j}) = (\sigma_1, 0_{j-i}, 0_{d-j}) = (\sigma_1, 0_{d-i}) = (\sigma, 0_{j+1}),$$

which proves (10.1).

If  $\tilde{\sigma}_2 \neq 0_{j-i}$ , let  $r = |\tilde{\sigma}_2| > 0$ . Since  $O(d-2i-1)$  acts transitively on  $\mathbb{S}^{d-2i-2}$  (thus, also on the sphere  $r\mathbb{S}^{d-2i-2}$ ), then there exists  $g_i \in O(d-2i-1)$  such that  $g_i(\tilde{\sigma}_2, 0_{d-j-i-1}) = (\sigma_2, \sigma_3, 0_{j-i}) \in r\mathbb{S}^{d-2i-2}$ . Let

$$\tilde{g}_i = id_{\mathbb{R}^{i+1}} \times g_i \times id_{\mathbb{R}^{i+1}} \in G_{d,i} \quad \text{and} \quad \tilde{g}_j = g_j \times id_{\mathbb{R}^{d-j}} \in G_{d,j}.$$

Then  $g_{ij} := \tilde{g}_i \tilde{g}_j \in \langle G_{d,i}; G_{d,j} \rangle$  and on account of (10.2) and  $i+1 < d-j$

(since  $i < j \leq s_d$ ), we have

$$\begin{aligned} \tilde{g}_i \tilde{g}_j(\omega, 0_{d-j}) &= \tilde{g}_i(g_j \omega, 0_{d-j}) = \tilde{g}_i(\sigma_1, \tilde{\sigma}_2, 0_{d-j}) \\ &= (\sigma_1, g_i(\tilde{\sigma}_2, 0_{d-j-i-1}), 0_{i+1}) \\ &= (\sigma_1, \sigma_2, \sigma_3, 0_{j-i}, 0_{i+1}) = (\sigma, 0_{j+1}), \end{aligned}$$

i.e., relation (10.1).

Now, let  $\bar{\sigma}, \tilde{\sigma} \in \mathbb{S}^{d-j-1}$ . Then, fixing  $\omega \in \mathbb{S}^j$ , on account of (10.1), there are  $g_1, g_2 \in \langle G_{d,i}; G_{d,j} \rangle$  such that  $g_1(\omega, 0_{d-j}) = (\bar{\sigma}, 0_{j+1})$  and  $g_2(\omega, 0_{d-j}) = (\tilde{\sigma}, 0_{j+1})$ . Consequently,  $g_2 g_1^{-1} \in \langle G_{d,i}; G_{d,j} \rangle$  and

$$g_2 g_1^{-1}(\bar{\sigma}, 0_{j+1}) = (\tilde{\sigma}, 0_{j+1}),$$

i.e., the group  $\langle G_{d,i}; G_{d,j} \rangle$  acts transitively on  $\mathbb{S}^{d-j-1} \times \{0_{j+1}\}$ .

*Step 2.* The group  $\langle G_{d,i}; G_{d,j} \rangle$  acts transitively on  $\mathbb{S}^{d-i-1} \times \{0_{i+1}\}$ .

We can proceed in a similar way as in Step 1; however, we sketch the proof. We show that for every  $\sigma = (\sigma_1, \sigma_2, \sigma_3) \in \mathbb{S}^{d-i-1}$  with  $\sigma_1 \in \mathbb{R}^{i+1}$ ,  $\sigma_2 \in \mathbb{R}^{d-j-i-1}$ ,  $\sigma_3 \in \mathbb{R}^{j-i}$ , and  $\omega \in \mathbb{S}^{d-j-1}$  fixed arbitrarily, there is  $g_{ij} \in \langle G_{d,i}; G_{d,j} \rangle$  such that

$$g_{ij}(\omega, 0_{j+1}) = (\sigma, 0_{i+1}). \quad (10.3)$$

Let  $\tilde{\sigma}_2 \in \mathbb{R}^{d-j-i-1}$  be such that  $|\sigma_1|^2 + |\tilde{\sigma}_2|^2 = 1$ . Then, due to Step 1, there exists  $\tilde{g}_{ij} \in \langle G_{d,i}; G_{d,j} \rangle$  such that  $\tilde{g}_{ij}(\omega, 0_{j+1}) = (\sigma_1, \tilde{\sigma}_2, 0_{j+1})$ .

If  $\tilde{\sigma}_2 = 0_{d-j-i-1}$  then  $\sigma_2 = 0_{d-j-i-1}$  and  $\sigma_3 = 0_{j-i}$ ; thus, (10.3) is verified with the choice  $g_{ij} := \tilde{g}_{ij} \in \langle G_{d,i}; G_{d,j} \rangle$ .

If  $\tilde{\sigma}_2 \neq 0_{d-j-i-1}$  then let  $r = |\tilde{\sigma}_2| > 0$ . Since  $O(d-2i-1)$  acts transitively on  $\mathbb{S}^{d-2i-2}$  (thus, also on the sphere  $r\mathbb{S}^{d-2i-2}$ ), then there exists  $g_i \in O(d-2i-1)$  such that  $g_i(\tilde{\sigma}_2, 0_{j-i}) = (\sigma_2, \sigma_3) \in r\mathbb{S}^{d-2i-2}$ . Let  $\tilde{g}_i = id_{\mathbb{R}^{i+1}} \times g_i \times id_{\mathbb{R}^{i+1}} \in G_{d,i}$ . Then

$$\tilde{g}_i \tilde{g}_{ij}(\omega, 0_{j+1}) = \tilde{g}_i(\sigma_1, \tilde{\sigma}_2, 0_{j+1}) = (\sigma_1, g_i(\tilde{\sigma}_2, 0_{j-i}), 0_{i+1}) = (\sigma, 0_{i+1}).$$

Consequently,  $g_{ij} := \tilde{g}_i \tilde{g}_{ij} \in \langle G_{d,i}; G_{d,j} \rangle$  verifies (10.3). Now, following the last part of Step 1, our claim follows.

*Step 3.* (Proof concluded) The group  $\langle G_{d,i}; G_{d,j} \rangle$  acts transitively on  $\mathbb{S}^d$ .

We show that for every  $\sigma = (\sigma_1, \sigma_2, \sigma_3) \in S^d$  with  $\sigma_1 \in \mathbb{R}^{i+1}$ ,  $\sigma_2 \in \mathbb{R}^{d-j-i-1}$ ,  $\sigma_3 \in \mathbb{R}^{j+1}$ , and  $\omega \in S^{d-i-1}$  fixed arbitrarily, there is  $g_{ij} \in \langle G_{d,i}; G_{d,j} \rangle$  such that

$$g_{ij}(\omega, 0_{i+1}) = \sigma. \quad (10.4)$$

Let  $\tilde{\sigma}_3 \in \mathbb{R}^{j-i}$  such that  $|\tilde{\sigma}_3| = |\sigma_3|$ . Then, due to Step 2, there exists  $\tilde{g}_{ij} \in \langle G_{d,i}; G_{d,j} \rangle$  such that  $\tilde{g}_{ij}(\omega, 0_{i+1}) = (\sigma_1, \sigma_2, \tilde{\sigma}_3, 0_{i+1})$ .

If  $\tilde{\sigma}_3 = 0_{j-i}$  then  $\sigma_3 = 0_{j+1}$  and (10.4) is verified by choosing  $g_{ij} := \tilde{g}_{ij} \in \langle G_{d,i}; G_{d,j} \rangle$ .

If  $\tilde{\sigma}_3 \neq 0_{j-i}$ , let  $r = |\tilde{\sigma}_3| = |\sigma_3| > 0$ . Since  $O(j+1)$  acts transitively on  $\mathbb{S}^j$ , there exists  $g_j \in O(j+1)$  such that  $g_j(\tilde{\sigma}_3, 0_{i+1}) = \sigma_3 \in r\mathbb{S}^j$ . Let us fix the element  $\tilde{g}_j = id_{\mathbb{R}^{d-j}} \times g_j \in G_{d,j}$ . Then

$$\begin{aligned} \tilde{g}_j \tilde{g}_{ij}(\omega, 0_{i+1}) &= \tilde{g}_j(\sigma_1, \sigma_2, \tilde{\sigma}_3, 0_{i+1}) = (\sigma_1, \sigma_2, g_j(\tilde{\sigma}_3, 0_{i+1})) \\ &= (\sigma_1, \sigma_2, \sigma_3) = \sigma. \end{aligned}$$

Consequently,  $g_{ij} := \tilde{g}_j \tilde{g}_{ij} \in \langle G_{d,i}; G_{d,j} \rangle$  verifies (10.4).

Now, let  $\bar{\sigma}, \tilde{\sigma} \in \mathbb{S}^d$ . Then, fixing  $\omega \in \mathbb{S}^{d-i-1}$ , on account of (10.4), there are  $g_1, g_2 \in \langle G_{d,i}; G_{d,j} \rangle$  such that  $g_1(\omega, 0_{i+1}) = \bar{\sigma}$  and  $g_2(\omega, 0_{i+1}) = \tilde{\sigma}$ . Consequently,  $g_2 g_1^{-1} \in \langle G_{d,i}; G_{d,j} \rangle$  and  $g_2 g_1^{-1}(\bar{\sigma}) = \tilde{\sigma}$ , i.e., the group  $\langle G_{d,i}; G_{d,j} \rangle$  acts transitively on  $\mathbb{S}^d$ . This completes the proof.  $\square$

Let  $d \geq 5$  and fix  $G_{d,i}$  for some  $i \in \{1, \dots, s_d\}$ . We define the function  $\tau_i : \mathbb{S}^d \rightarrow \mathbb{S}^d$  associated to  $G_{d,i}$  by

$$\tau_i(\sigma) = \begin{cases} (\sigma_3, \sigma_2, \sigma_1), & \text{if } i \neq \frac{d-1}{2}, \text{ and } \sigma = (\sigma_1, \sigma_2, \sigma_3) \text{ with } \sigma_1, \sigma_3 \in \mathbb{R}^{i+1}, \\ & \sigma_2 \in \mathbb{R}^{d-2i-1}; \\ (\sigma_3, \sigma_1), & \text{if } i = \frac{d-1}{2}, \text{ and } \sigma = (\sigma_1, \sigma_3) \text{ with } \sigma_1, \sigma_3 \in \mathbb{R}^{\frac{d+1}{2}}. \end{cases}$$

The explicit form of the groups  $G_{d,i}$  and the functions  $\tau_i$  can be seen in Table 1 for dimensions  $d = 5, \dots, 12$ . It is clear by construction that  $\tau_i \notin G_{d,i}$ ,  $\tau_i G_{d,i} \tau_i^{-1} = G_{d,i}$  and  $\tau_i^2 = id_{\mathbb{R}^{d+1}}$ .

$d$	$s_d$	$G_{d,i}; i \in \{1, \dots, s_d\}$	$\tau_i; i \in \{1, \dots, s_d\}$
5	2	$G_{5,1} = O(2) \times O(2) \times O(2)$ $G_{5,2} = O(3) \times O(3)$	$\tau_1(\sigma_1, \sigma_2, \sigma_3) = (\sigma_3, \sigma_2, \sigma_1); \sigma_1, \sigma_2, \sigma_3 \in \mathbb{R}^2$ $\tau_2(\sigma_1, \sigma_2) = (\sigma_2, \sigma_1); \sigma_1, \sigma_2 \in \mathbb{R}^3$
6	1	$G_{6,1} = O(2) \times O(3) \times O(2)$	$\tau_1(\sigma_1, \sigma_2, \sigma_3) = (\sigma_3, \sigma_2, \sigma_1); \sigma_1, \sigma_3 \in \mathbb{R}^2, \sigma_2 \in \mathbb{R}^3$
7	3	$G_{7,1} = O(2) \times O(4) \times O(2)$ $G_{7,2} = O(3) \times O(2) \times O(3)$ $G_{7,3} = O(4) \times O(4)$	$\tau_1(\sigma_1, \sigma_2, \sigma_3) = (\sigma_3, \sigma_2, \sigma_1); \sigma_1, \sigma_3 \in \mathbb{R}^2, \sigma_2 \in \mathbb{R}^4$ $\tau_2(\sigma_1, \sigma_2, \sigma_3) = (\sigma_3, \sigma_2, \sigma_1); \sigma_1, \sigma_3 \in \mathbb{R}^3, \sigma_2 \in \mathbb{R}^2$ $\tau_3(\sigma_1, \sigma_2) = (\sigma_2, \sigma_1); \sigma_1, \sigma_2 \in \mathbb{R}^4$
8	2	$G_{8,1} = O(2) \times O(5) \times O(2)$ $G_{8,2} = O(3) \times O(3) \times O(3)$	$\tau_1(\sigma_1, \sigma_2, \sigma_3) = (\sigma_3, \sigma_2, \sigma_1); \sigma_1, \sigma_3 \in \mathbb{R}^2, \sigma_2 \in \mathbb{R}^5$ $\tau_2(\sigma_1, \sigma_2, \sigma_3) = (\sigma_2, \sigma_1, \sigma_3); \sigma_1, \sigma_2, \sigma_3 \in \mathbb{R}^3$
9	4	$G_{9,1} = O(2) \times O(6) \times O(2)$ $G_{9,2} = O(3) \times O(4) \times O(3)$ $G_{9,3} = O(4) \times O(2) \times O(4)$ $G_{9,4} = O(5) \times O(5)$	$\tau_1(\sigma_1, \sigma_2, \sigma_3) = (\sigma_3, \sigma_2, \sigma_1); \sigma_1, \sigma_3 \in \mathbb{R}^2, \sigma_2 \in \mathbb{R}^6$ $\tau_2(\sigma_1, \sigma_2, \sigma_3) = (\sigma_3, \sigma_2, \sigma_1); \sigma_1, \sigma_3 \in \mathbb{R}^3, \sigma_2 \in \mathbb{R}^4$ $\tau_3(\sigma_1, \sigma_2, \sigma_3) = (\sigma_3, \sigma_2, \sigma_1); \sigma_1, \sigma_3 \in \mathbb{R}^4, \sigma_2 \in \mathbb{R}^2$ $\tau_4(\sigma_1, \sigma_2) = (\sigma_2, \sigma_1); \sigma_1, \sigma_2 \in \mathbb{R}^5$
10	3	$G_{10,1} = O(2) \times O(7) \times O(2)$ $G_{10,2} = O(3) \times O(5) \times O(3)$ $G_{10,3} = O(4) \times O(3) \times O(4)$	$\tau_1(\sigma_1, \sigma_2, \sigma_3) = (\sigma_3, \sigma_2, \sigma_1); \sigma_1, \sigma_3 \in \mathbb{R}^2, \sigma_2 \in \mathbb{R}^7$ $\tau_2(\sigma_1, \sigma_2, \sigma_3) = (\sigma_3, \sigma_2, \sigma_1); \sigma_1, \sigma_3 \in \mathbb{R}^3, \sigma_2 \in \mathbb{R}^5$ $\tau_3(\sigma_1, \sigma_2, \sigma_3) = (\sigma_3, \sigma_2, \sigma_1); \sigma_1, \sigma_3 \in \mathbb{R}^4, \sigma_2 \in \mathbb{R}^3$
11	5	$G_{11,1} = O(2) \times O(8) \times O(2)$ $G_{11,2} = O(3) \times O(6) \times O(3)$ $G_{11,3} = O(4) \times O(4) \times O(4)$ $G_{11,4} = O(5) \times O(2) \times O(5)$ $G_{11,5} = O(6) \times O(6)$	$\tau_1(\sigma_1, \sigma_2, \sigma_3) = (\sigma_3, \sigma_2, \sigma_1); \sigma_1, \sigma_3 \in \mathbb{R}^2, \sigma_2 \in \mathbb{R}^8$ $\tau_2(\sigma_1, \sigma_2, \sigma_3) = (\sigma_3, \sigma_2, \sigma_1); \sigma_1, \sigma_3 \in \mathbb{R}^3, \sigma_2 \in \mathbb{R}^6$ $\tau_3(\sigma_1, \sigma_2, \sigma_3) = (\sigma_2, \sigma_1, \sigma_3); \sigma_1, \sigma_2, \sigma_3 \in \mathbb{R}^4$ $\tau_4(\sigma_1, \sigma_2, \sigma_3) = (\sigma_3, \sigma_2, \sigma_1); \sigma_1, \sigma_3 \in \mathbb{R}^5, \sigma_2 \in \mathbb{R}^2$ $\tau_5(\sigma_1, \sigma_2) = (\sigma_2, \sigma_1); \sigma_1, \sigma_2 \in \mathbb{R}^6$
12	4	$G_{12,1} = O(2) \times O(9) \times O(2)$ $G_{12,2} = O(3) \times O(7) \times O(3)$ $G_{12,3} = O(4) \times O(5) \times O(4)$ $G_{12,4} = O(5) \times O(3) \times O(5)$	$\tau_1(\sigma_1, \sigma_2, \sigma_3) = (\sigma_3, \sigma_2, \sigma_1); \sigma_1, \sigma_3 \in \mathbb{R}^2, \sigma_2 \in \mathbb{R}^9$ $\tau_2(\sigma_1, \sigma_2, \sigma_3) = (\sigma_3, \sigma_2, \sigma_1); \sigma_1, \sigma_3 \in \mathbb{R}^3, \sigma_2 \in \mathbb{R}^7$ $\tau_3(\sigma_1, \sigma_2, \sigma_3) = (\sigma_3, \sigma_2, \sigma_1); \sigma_1, \sigma_3 \in \mathbb{R}^4, \sigma_2 \in \mathbb{R}^5$ $\tau_4(\sigma_1, \sigma_2, \sigma_3) = (\sigma_3, \sigma_2, \sigma_1); \sigma_1, \sigma_3 \in \mathbb{R}^5, \sigma_2 \in \mathbb{R}^3$

TABLE 1.

Inspired by [31], [33], we introduce the action of the group  $G_{d,i}^{\tau_i} = \langle G_{d,i}, \tau_i \rangle \subset O(d+1)$  on the space  $H_1^2(\mathbb{S}^d)$ . Due to the above properties of  $\tau_i$ , only two types of elements in  $G_{d,i}^{\tau_i}$  can be distinguished; namely,  $\tilde{g} = g \in G_{d,i}$ , and  $\tilde{g} = \tau_i g \in G_{d,i}^{\tau_i} \setminus G_{d,i}$  (with  $g \in G_{d,i}$ ), respectively. Therefore, the action  $G_{d,i}^{\tau_i} \times H_1^2(\mathbb{S}^d) \rightarrow H_1^2(\mathbb{S}^d)$  given by

$$gu(\sigma) = u(g^{-1}\sigma), \quad (\tau_i g)u(\sigma) = -u(g^{-1}\tau_i^{-1}\sigma), \quad (10.5)$$

for  $g \in G_{d,i}$ ,  $u \in H_1^2(\mathbb{S}^d)$  and  $\sigma \in \mathbb{S}^d$ , is well-defined, continuous and linear. We define the subspace of  $H_1^2(\mathbb{S}^d)$  containing all symmetric points with respect to the compact group  $G_{d,i}^{\tau_i}$ , i.e.,

$$H_{G_{d,i}^{\tau_i}}(\mathbb{S}^d) = \{u \in H_1^2(\mathbb{S}^d) : \tilde{g}u = u \text{ for every } \tilde{g} \in G_{d,i}^{\tau_i}\}.$$

For further use, we also introduce

$$H_{G_{d,i}}(\mathbb{S}^d) = \{u \in H_1^2(\mathbb{S}^d) : gu = u \text{ for every } g \in G_{d,i}\},$$

where the action of the group  $G_{d,i}$  on  $H_1^2(\mathbb{S}^d)$  is defined by the first relation of (10.5).

**Remark 10.1** Every nonzero element of the space  $H_{G_{d,i}^{\tau_i}}(\mathbb{S}^d)$  changes sign. To see this, let  $u \in H_{G_{d,i}^{\tau_i}}(\mathbb{S}^d) \setminus \{0\}$ . Due to the  $G_{d,i}^{\tau_i}$ -invariance of  $u$  and (10.5) we have  $u(\sigma) = -u(\tau_i^{-1}\sigma)$  for every  $\sigma \in \mathbb{S}^d$ . Since  $u \neq 0$ , it should change the sign.

**Theorem 10.1** For every  $i, j \in \{1, \dots, s_d\}$  with  $i \neq j$ , one has

- (i).  $H_{G_{d,i}}(\mathbb{S}^d) \cap H_{G_{d,j}}(\mathbb{S}^d) = \{\text{constant functions on } \mathbb{S}^d\}$ ;
- (ii).  $H_{G_{d,i}^{\tau_i}}(\mathbb{S}^d) \cap H_{G_{d,j}^{\tau_j}}(\mathbb{S}^d) = \{0\}$ .

*Proof.* (i) Let  $u \in H_{G_{d,i}}(\mathbb{S}^d) \cap H_{G_{d,j}}(\mathbb{S}^d)$ . In particular,  $u$  is both  $G_{d,i}$ - and  $G_{d,j}$ -invariant, i.e.  $g_i u = g_j u = u$  for every  $g_i \in G_{d,i}$  and  $g_j \in G_{d,j}$ , respectively. Consequently,  $u$  is also  $\langle G_{d,i}, G_{d,j} \rangle$ -invariant; thus,  $u(\sigma) = u(g_{ij}\sigma)$  for every  $g_{ij} \in \langle G_{d,i}, G_{d,j} \rangle$  and  $\sigma \in \mathbb{S}^d$ . Due to Proposition 10.1, for every fixed  $\sigma \in \mathbb{S}^d$ , the orbit of  $g_{ij}\sigma$  is the whole sphere  $\mathbb{S}^d$  whenever  $g_{ij}$  runs through  $\langle G_{d,i}, G_{d,j} \rangle$ . Therefore,  $u$  should be constant.

(ii) Let  $u \in H_{G_{d,i}^{\tau_i}}(\mathbb{S}^d) \cap H_{G_{d,j}^{\tau_j}}(\mathbb{S}^d)$ . The second relation of (10.5) shows that  $u(\sigma) = -u(\tau_i^{-1}\sigma) = -u(\tau_j^{-1}\sigma)$ ,  $\sigma \in \mathbb{S}^d$ . But, due to (i),  $u$  is constant. Thus,  $u$  should be 0.  $\square$

To conclude this section, we construct explicit functions belonging to  $H_{G_{d,i}^{\tau_i}}(\mathbb{S}^d)$  which is of interest in its own right as well. Before we give the class of functions we are speaking about, we say that a set  $D \subset \mathbb{S}^d$  is  $G_{d,i}^{\tau_i}$ -invariant, if  $\tilde{g}D \subseteq D$  for every  $\tilde{g} \in G_{d,i}^{\tau_i}$ .

**Proposition 10.2** Let  $i \in \{1, \dots, s_d\}$  and  $s > 0$  be fixed. Then there exist a number  $C_i > 0$  and a  $G_{d,i}^{\tau_i}$ -invariant set  $D_i \subset \mathbb{S}^d$  with  $\text{Vol}_h(D_i) > 0$ , both independent on the number  $s$ , and a function  $w \in H_{G_{d,i}^{\tau_i}}(\mathbb{S}^d)$  such that

- (i).  $\|w\|_{L^\infty} \leq s$ ;
- (ii).  $|\nabla w(\sigma)|_h \leq C_i s$  for a.e.  $\sigma \in \mathbb{S}^d$ ;
- (iii).  $|w(\sigma)| = s$  for every  $\sigma \in D_i$ .

An explicit function  $w : \mathbb{S}^d \rightarrow \mathbb{R}$  fulfilling all the requirements of Proposition 10.2 is given by

$$w(\sigma) = \frac{8s}{(R-r)} \text{sgn}(|\sigma_1| - |\sigma_3|) \max \left( 0, \min \left( \frac{R-r}{8}, \frac{R-r}{4} - \right. \right. \quad (10.6) \\ \left. \left. - \max \left( \left| |\sigma_1| + |\sigma_3| - \frac{R+3r}{4} \right|, \left| |\sigma_1| - |\sigma_3| - \frac{R+3r}{4} \right| \right) \right) \right)$$




Figure 10.1 The image of the function  $w : \mathbb{S}^d \rightarrow \mathbb{R}$  from (10.6) with parameters  $r = 0.2, R = 1.5, s = 0.4$ ; the value  $w(\sigma)$  is represented (radially) on the line determined by  $0 \in \mathbb{R}^{d+1}$  and  $\sigma \in \mathbb{S}^d$ , the 'zero altitude' being  $c\sigma$ , i.e., the sphere  $c\mathbb{S}^d$ , with  $c = 1.3$ . The union of those 8 disconnected holes on the sphere  $\mathbb{S}^d$  where the function  $w$  takes values  $s$  and  $(-s)$  corresponds to the  $G_{d,i}^r$ -invariant set  $D_i$ . (Note that the figure describes the case  $i \neq \frac{d-1}{2}$ . When  $i = \frac{d-1}{2}$  the coordinate  $\sigma_2$  vanishes and the figure becomes simpler.)

where  $R > r$ , and  $\sigma = (\sigma_1, \sigma_2, \sigma_3) \in \mathbb{S}^d$  with  $\sigma_1, \sigma_3 \in \mathbb{R}^{i+1}, \sigma_2 \in \mathbb{R}^{d-2i-1}$  whenever  $i \neq \frac{d-1}{2}$ , and  $\sigma = (\sigma_1, \sigma_3) \in \mathbb{S}^d$  with  $\sigma_1, \sigma_3 \in \mathbb{R}^{\frac{d+1}{2}}$  whenever  $i = \frac{d-1}{2}$ . The  $G_{d,i}^r$ -invariant set  $D_i \subset \mathbb{S}^d$  can be defined as

$$D_i = \left\{ \sigma \in \mathbb{S}^d : \left| |\sigma_1| + |\sigma_3| - \frac{R+3r}{4} \right| \leq \frac{R-r}{8}, \left| \left| |\sigma_1| - |\sigma_3| \right| - \frac{R+3r}{4} \right| \leq \frac{R-r}{8} \right\}.$$

The geometrical image of the function  $w$  from (10.6) is shown by Figure 10.1.

### 10.3 Arbitrarily small solutions

Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be a continuous function and  $F(s) = \int_0^s f(t)dt$ . We assume that

$$(f_1^0). \quad -\infty < \liminf_{s \rightarrow 0^+} \frac{F(s)}{s^2} \leq \limsup_{s \rightarrow 0^+} \frac{F(s)}{s^2} = +\infty;$$

$$(f_2^0). \quad \liminf_{s \rightarrow 0^+} \frac{f(s)}{s} < 0.$$

**Theorem 10.2** *Let  $d \geq 5$  and  $f : \mathbb{R} \rightarrow \mathbb{R}$  be a continuous function which is odd in an arbitrarily small neighborhood of the origin, verifying  $(f_1^0)$  and  $(f_2^0)$ . Then there exist at least  $s_d = [d/2] + (-1)^{d+1} - 1$  sequences  $\{u_k^i\}_k \subset H_1^2(\mathbb{S}^d)$ ,  $i \in \{1, \dots, s_d\}$ , of sign-changing weak solutions of (P) distinguished by their symmetry properties. In addition,*

$$\lim_{k \rightarrow \infty} \|u_k^i\|_{L^\infty} = \lim_{k \rightarrow \infty} \|u_k^i\|_{H_1^2} = 0 \text{ for every } i \in \{1, \dots, s_d\}.$$

**Example 10.1** Let  $\alpha, \beta, \gamma \in \mathbb{R}$  such that  $\alpha + \beta > 1 > \alpha > 0$ , and  $\gamma \in (0, 1)$ . Then, the function  $f : \mathbb{R} \rightarrow \mathbb{R}$  defined by  $f(0) = 0$  and  $f(s) = |s|^{\alpha-1} s(\gamma + \sin |s|^{-\beta})$  near the origin (but  $s \neq 0$ ) and extended in an arbitrarily way to the whole  $\mathbb{R}$ , verifies both  $(f_1^0)$  and  $(f_2^0)$ .

In order to prove Theorem 10.2, we need some propositions. Throughout this section we assume the hypotheses of Theorem 10.2 are fulfilled.

Let  $\tilde{s} > 0$  be so small that  $f$  is odd on  $[-\tilde{s}, \tilde{s}]$ , and let us define  $\tilde{f}(s) = \text{sgn}(s)f(\min(|s|, \tilde{s}))$ . Clearly,  $\tilde{f}$  is continuous and odd on  $\mathbb{R}$ . Define also  $\tilde{F}(s) = \int_0^s \tilde{f}(t)dt$ ,  $s \in \mathbb{R}$ .

On account of  $(f_2^0)$ , one may fix  $c_0 > 0$  such that

$$\liminf_{s \rightarrow 0^+} \frac{f(s)}{s} < -c_0 < 0. \tag{10.7}$$

In particular, there is a sequence  $\{\bar{s}_k\}_k \subset (0, \tilde{s})$  converging (decreasingly) to 0, such that

$$\tilde{f}(\bar{s}_k) = f(\bar{s}_k) < -c_0 \bar{s}_k. \tag{10.8}$$

Let us define the functions

$$\psi(s) = \tilde{f}(s) + c_0 s \text{ and } \Psi(s) = \int_0^s \psi(t)dt = \tilde{F}(s) + \frac{c_0}{2} s^2, \quad s \in \mathbb{R}. \tag{10.9}$$

Due to (10.8),  $\psi(\bar{s}_k) < 0$ ; so, there are two sequences  $\{a_k\}_k, \{b_k\}_k \subset (0, \tilde{s})$ , both converging to 0, such that  $b_{k+1} < a_k < \bar{s}_k < b_k$  for every  $k \in \mathbb{N}$  and

$$\psi(s) \leq 0 \text{ for every } s \in [a_k, b_k]. \tag{10.10}$$

Since  $c_0 > 0$ , see (10.7), the norm

$$\|u\|_{c_0} = \left( \int_{\mathbb{S}^d} |\nabla u|_h^2 d\sigma_h + c_0 \int_{\mathbb{S}^d} u^2 d\sigma_h \right)^{1/2} \tag{10.11}$$

is equivalent to the standard norm  $\|\cdot\|_{H_1^2}$ . Now, we define  $\mathcal{E} : H_1^2(\mathbb{S}^d) \rightarrow \mathbb{R}$  by

$$\mathcal{E}(u) = \frac{1}{2}\|u\|_{c_0}^2 - \int_{\mathbb{S}^d} \Psi(u(\sigma))d\sigma_h,$$

which is well-defined since  $\psi$  has a subcritical growth, and  $H_1^2(\mathbb{S}^d)$  is compactly embedded into  $L^p(\mathbb{S}^d)$ ,  $p \in [1, 2^*)$ . Moreover,  $\mathcal{E}$  belongs to  $C^1(H_1^2(\mathbb{S}^d))$ , it is even, and it coincides with the energy functional associated to (P) on the set  $B^\infty(\tilde{s}) = \{u \in L^\infty(\mathbb{S}^d) : \|u\|_{L^\infty} \leq \tilde{s}\}$  because the functions  $f$  and  $\tilde{f}$  coincide on  $[-\tilde{s}, \tilde{s}]$ .

From now on, we fix  $i \in \{1, \dots, s_d\}$  and the corresponding subspace  $H_{G_{d,i}^{\tau_i}}(\mathbb{S}^d)$  of  $H_1^2(\mathbb{S}^d)$  introduced in the previous section. Let us denote by  $\mathcal{E}_i$  the restriction of the functional  $\mathcal{E}$  to  $H_{G_{d,i}^{\tau_i}}(\mathbb{S}^d)$  and for every  $k \in \mathbb{N}$ , consider the set

$$T_k^i = \{u \in H_{G_{d,i}^{\tau_i}}(\mathbb{S}^d) : \|u\|_{L^\infty} \leq b_k\}, \tag{10.12}$$

where  $b_k$  is from (10.10).

**Proposition 10.3** *The functional  $\mathcal{E}_i$  is bounded from below on  $T_k^i$  and its infimum  $m_k^i$  on  $T_k^i$  is attained at  $u_k^i \in T_k^i$ . Moreover,  $m_k^i = \mathcal{E}_i(u_k^i) < 0$  for every  $k \in \mathbb{N}$ .*

*Proof.* For every  $u \in T_k^i$  we have

$$\mathcal{E}_i(u) = \frac{1}{2}\|u\|_{c_0}^2 - \int_{\mathbb{S}^d} \Psi(u(\sigma))d\sigma_h \geq - \max_{[-b_k, b_k]} \Psi \cdot \text{Vol}_h(\mathbb{S}^d) > -\infty.$$

It is clear that  $T_k^i$  is convex and closed, thus weakly closed in  $H_{G_{d,i}^{\tau_i}}(\mathbb{S}^d)$ . Let  $m_k^i = \inf_{T_k^i} \mathcal{E}_i$ , and  $\{u_n\}_n \subset T_k^i$  be a minimizing sequence of  $\mathcal{E}_i$  for  $m_k^i$ . Then, for large  $n \in \mathbb{N}$ , we have

$$\frac{1}{2}\|u_n\|_{c_0}^2 \leq m_k^i + 1 + \max_{[-b_k, b_k]} \Psi \cdot \text{Vol}_h(\mathbb{S}^d),$$

thus  $\{u_n\}_n$  is bounded in  $H_{G_{d,i}^{\tau_i}}(\mathbb{S}^d)$ . Up to a subsequence,  $\{u_n\}_n$  weakly converges in  $H_{G_{d,i}^{\tau_i}}(\mathbb{S}^d)$  to some  $u_k^i \in T_k^i$ . Since  $\psi$  has a subcritical growth, by using the compactness of the embedding  $H_{G_{d,i}^{\tau_i}}(\mathbb{S}^d) \subset H_1^2(\mathbb{S}^d) \hookrightarrow L^p(\mathbb{S}^d)$ ,  $1 \leq p < 2^*$ , one can conclude the sequentially weak continuity of the function  $u \mapsto \int_{\mathbb{S}^d} \Psi(u(\sigma))d\sigma_h$ ,  $u \in H_{G_{d,i}^{\tau_i}}(\mathbb{S}^d)$ . Consequently,  $\mathcal{E}_i$  is sequentially weak lower semicontinuous. Combining this fact with the weak closedness of the set  $T_k^i$ , we obtain  $\mathcal{E}_i(u_k^i) = m_k^i = \inf_{T_k^i} \mathcal{E}_i$ .

The next task is to prove that  $m_k^i < 0$  for every  $k \in \mathbb{N}$ . First, due to (10.9) and  $(f_1^0)$ , we have

$$-\infty < \liminf_{s \rightarrow 0^+} \frac{\Psi(s)}{s^2} \leq \limsup_{s \rightarrow 0^+} \frac{\Psi(s)}{s^2} = +\infty. \tag{10.13}$$

Therefore, the left-hand side of (10.13) and the evenness of  $\Psi$  implies the existence of  $\underline{l} > 0$  and  $\varrho \in (0, \bar{s})$  such that

$$\Psi(s) \geq -\underline{l}s^2 \text{ for every } s \in (-\varrho, \varrho). \tag{10.14}$$

Let  $D_i \subset \mathbb{S}^d$  and  $C_i > 0$  be from Proposition 10.2 (which depend only on  $G_{d,i}$  and  $\tau_i$ ), and fix a number  $\bar{l} > 0$  large enough such that

$$\bar{l}\text{Vol}_h(D_i) > \left(\underline{l} + \frac{c_0}{2}\right) \text{Vol}_h(\mathbb{S}^d) + \frac{C_i^2}{2}, \tag{10.15}$$

$c_0 > 0$  being from (10.7). Taking into account the right-hand side of (10.13), there is a sequence  $\{s_k\}_k \subset (0, \varrho)$  such that  $s_k \leq b_k$  and  $\Psi(s_k) = \Psi(-s_k) > \bar{l}s_k^2$  for every  $k \in \mathbb{N}$ .

Let  $w_k := w_{s_k} \in H_{G_{d,i}^{\tau_i}}(\mathbb{S}^d)$  be the function from Proposition 10.2 corresponding to the value  $s_k > 0$ . Then  $w_k \in T_k^i$  and one has

$$\begin{aligned} \mathcal{E}_i(w_k) &= \frac{1}{2} \|w_k\|_{c_0}^2 - \int_{\mathbb{S}^d} \Psi(w_k(\sigma)) d\sigma_h \\ &\leq \frac{1}{2} (C_i^2 + c_0 \text{Vol}_h(\mathbb{S}^d)) s_k^2 - \int_{D_i} \Psi(w_k(\sigma)) d\sigma_h \\ &\quad - \int_{\mathbb{S}^d \setminus D_i} \Psi(w_k(\sigma)) d\sigma_h. \end{aligned}$$

On account of Proposition 10.2 (iii), we have

$$\int_{D_i} \Psi(w_k(\sigma)) d\sigma_h = \Psi(s_k) \text{Vol}_h(D_i) > \bar{l} \text{Vol}_h(D_i) s_k^2.$$

On the other hand, due to relation (10.14) and Proposition 10.2 (i), we have

$$\int_{\mathbb{S}^d \setminus D_i} \Psi(w_k(\sigma)) d\sigma_h \geq -\underline{l} \int_{\mathbb{S}^d \setminus D_i} w_k^2(\sigma) d\sigma_h > -\underline{l} \text{Vol}_h(\mathbb{S}^d) s_k^2.$$

Combining (10.15) with the above estimations, we obtain that  $m_k^i = \inf_{T_k^i} \mathcal{E}_i \leq \mathcal{E}_i(w_k) < 0$ , which proves our claim.  $\square$

**Proposition 10.4** *Let  $u_k^i \in T_k^i$  from Proposition 10.3. Then,  $\|u_k^i\|_{L^\infty} \leq a_k$ . (The number  $a_k$  is from (10.10).)*

*Proof.* Let  $A = \{\sigma \in \mathbb{S}^d : u_k^i(\sigma) \notin [-a_k, a_k]\}$  and suppose that  $\text{meas}(A) > 0$ . Define the function  $\gamma(s) = \text{sgn}(s) \min(|s|, a_k)$  and set  $w_k = \gamma \circ u_k^i$ . Since  $\gamma$  is Lipschitz continuous, then  $w_k \in H_1^2(\mathbb{S}^d)$ , see Hebey [138, Proposition 2.5, p. 24].

We first claim that  $w_k \in H_{G_{d,i}^{\tau_i}}(\mathbb{S}^d)$ . To see this, it suffices to prove that  $\tilde{g}w_k = w_k$  for every  $\tilde{g} \in G_{d,i}^{\tau_i}$ . First, let  $\tilde{g} = g \in G_{d,i}$ . Since  $gu_k^i = u_k^i$ , we have

$$gw_k(\sigma) = w_k(g^{-1}\sigma) = (\gamma \circ u_k^i)(g^{-1}\sigma) = \gamma(u_k^i(g^{-1}\sigma)) = \gamma(u_k^i(\sigma)) = w_k(\sigma)$$

for every  $\sigma \in \mathbb{S}^d$ . Now, let  $\tilde{g} = \tau_i g \in G_{d,i}^{\tau_i} \setminus G_{d,i}$  (with  $g \in G_{d,i}$ ). Since  $\gamma$  is an odd function and  $(\tau_i g)u_k^i = u_k^i$ , on account of (10.5) we have

$$\begin{aligned} (\tau_i g)w_k(\sigma) &= -w_k(g^{-1}\tau_i^{-1}\sigma) = -(\gamma \circ u_k^i)(g^{-1}\tau_i^{-1}\sigma) \\ &= \gamma(-u_k^i(g^{-1}\tau_i^{-1}\sigma)) = \gamma((\tau_i g)u_k^i(\sigma)) = \gamma(u_k^i(\sigma)) \\ &= w_k(\sigma) \end{aligned}$$

for every  $\sigma \in \mathbb{S}^d$ . In conclusion, the claim is true, and  $w_k \in H_{G_{d,i}^{\tau_i}}(\mathbb{S}^d)$ . Moreover,  $\|w_k\|_{L^\infty} \leq a_k$ . Consequently,  $w_k \in T_k^i$ .

We introduce the sets

$$A_1 = \{\sigma \in A : u_k^i(\sigma) < -a_k\} \quad \text{and} \quad A_2 = \{\sigma \in A : u_k^i(\sigma) > a_k\}.$$

Thus,  $A = A_1 \cup A_2$ , and we have that  $w_k(\sigma) = u_k^i(\sigma)$  for all  $\sigma \in \mathbb{S}^d \setminus A$ ,  $w_k(\sigma) = -a_k$  for all  $\sigma \in A_1$ , and  $w_k(\sigma) = a_k$  for all  $\sigma \in A_2$ . Moreover,

$$\begin{aligned} \mathcal{E}_i(w_k) - \mathcal{E}_i(u_k^i) &= \\ &= -\frac{1}{2} \int_A |\nabla u_k^i|_h^2 d\sigma_h + \frac{c_0}{2} \int_A [w_k^2 - (u_k^i)^2] d\sigma_h \\ &\quad - \int_A [\Psi(w_k) - \Psi(u_k^i)] d\sigma_h \\ &= -\frac{1}{2} \int_A |\nabla u_k^i(\sigma)|_h^2 d\sigma_h + \frac{c_0}{2} \int_A [a_k^2 - (u_k^i(\sigma))^2] d\sigma_h \\ &\quad - \int_{A_1} [\Psi(-a_k) - \Psi(u_k^i(\sigma))] d\sigma_h - \int_{A_2} [\Psi(a_k) - \Psi(u_k^i(\sigma))] d\sigma_h. \end{aligned}$$

Note that  $\int_A [w_k^2 - (u_k^i)^2] d\sigma_h \leq 0$ . Next, by the mean value theorem, for a.e.  $\sigma \in A_2$ , there exists  $\theta_k(\sigma) \in [a_k, b_k]$  such that  $\Psi(a_k) - \Psi(u_k^i(\sigma)) = \psi(\theta_k(\sigma))(a_k - u_k^i(\sigma))$ . Thus, on account of (10.10), one has

$$\int_{A_2} [\Psi(a_k) - \Psi(u_k^i(\sigma))] d\sigma_h \geq 0.$$

In the same way, using the oddness of  $\psi$ , we conclude that

$$\int_{A_1} [\Psi(-a_k) - \Psi(u_k^i(\sigma))] d\sigma_h \geq 0.$$

In conclusion, every term of the expression  $\mathcal{E}_i(w_k) - \mathcal{E}_i(u_k^i)$  is nonpositive. On the other hand, since  $w_k \in T_k^i$ , then  $\mathcal{E}_i(w_k) \geq \mathcal{E}_i(u_k^i) = \inf_{T_k^i} \mathcal{E}_i$ . So, every term in  $\mathcal{E}_i(w_k) - \mathcal{E}_i(u_k^i)$  should be zero. In particular,

$$\int_A |\nabla u_k^i(\sigma)|_h^2 d\sigma_h = \int_A [a_k^2 - (u_k^i(\sigma))^2] d\sigma_h = 0.$$

These equalities imply that  $\text{meas}(A)$  should be 0, contradicting our initial assumption.  $\square$

**Proposition 10.5**  $\lim_{k \rightarrow \infty} m_k^i = \lim_{k \rightarrow \infty} \|u_k^i\|_{L^\infty} = \lim_{k \rightarrow \infty} \|u_k^i\|_{H_1^2} = 0.$

*Proof.* Using Proposition 10.4, we have that  $\|u_k^i\|_{L^\infty} \leq a_k < \tilde{s}$  for a.e.  $\sigma \in \mathbb{S}^d$ . Therefore, we readily have that  $\lim_{k \rightarrow \infty} \|u_k^i\|_{L^\infty} = 0$ .

Moreover, the mean value theorem shows that

$$\begin{aligned} m_k^i = \mathcal{E}_i(u_k^i) &\geq - \int_{\mathbb{S}^d} \Psi(u_k^i(\sigma)) d\sigma_h \geq - \max_{[-\tilde{s}, \tilde{s}]} |\psi| \int_{\mathbb{S}^d} |u_k^i(\sigma)| d\sigma_h \\ &\geq - \max_{[-\tilde{s}, \tilde{s}]} |\psi| \text{Vol}_h(\mathbb{S}^d) a_k. \end{aligned}$$

Since  $\lim_{k \rightarrow \infty} a_k = 0$ , we have  $\lim_{k \rightarrow \infty} m_k^i \geq 0$ . On the other hand,  $m_k < 0$  for every  $k \in \mathbb{N}$ , see Proposition 10.3, which implies  $\lim_{k \rightarrow \infty} m_k^i = 0$ .

Note that

$$\frac{\|u_k^i\|_{c_0}^2}{2} = m_k^i + \int_{\mathbb{S}^d} \Psi(u_k^i(\sigma)) d\sigma_h \leq m_k^i + \max_{[-\tilde{s}, \tilde{s}]} |\psi| \text{Vol}_h(\mathbb{S}^d) a_k,$$

thus  $\lim_{k \rightarrow \infty} \|u_k^i\|_{c_0} = 0$ . But  $\|\cdot\|_{c_0}$  and  $\|\cdot\|_{H_1^2}$  are equivalent norms.  $\square$

Now, we prove the key result of this chapter where the non-smooth principle of symmetric criticality for Szulkin-type functions plays a crucial role.

**Proposition 10.6**  $u_k^i$  is a weak solution of (P) for every  $k \in \mathbb{N}$ .

*Proof.* We divide the proof into two parts. First, let

$$T_k = \{u \in H_1^2(\mathbb{S}^d) : \|u\|_{L^\infty} \leq b_k\}.$$

*Step 1.*  $\langle \mathcal{E}'(u_k^i), w - u_k^i \rangle_{H_1^2} \geq 0$  for every  $w \in T_k$ .

The set  $T_k$  is closed and convex in  $H_1^2(\mathbb{S}^d)$ . Let  $\zeta_{T_k}$  be the indicator

function of the set  $T_k$  (i.e.,  $\zeta_{T_k}(u) = 0$  if  $u \in T_k$ , and  $\zeta_{T_k}(u) = +\infty$ , otherwise). We define the Szulkin-type functional  $\mathcal{I}_k : H_1^2(\mathbb{S}^d) \rightarrow \mathbb{R} \cup \{+\infty\}$  by  $\mathcal{I}_k = \mathcal{E} + \zeta_{T_k}$ , see Section 1.6. We deduce that  $\mathcal{E}$  is of class  $C^1(H_1^2(\mathbb{S}^d))$ ,  $\zeta_{T_k}$  is convex, lower semicontinuous and proper. On account of (10.12), we have that  $T_k^i = T_k \cap H_{G_{d,i}^{\tau_i}}(\mathbb{S}^d)$ ; therefore, the restriction of  $\zeta_{T_k}$  to  $H_{G_{d,i}^{\tau_i}}(\mathbb{S}^d)$  is precisely the indicator function  $\zeta_{T_k^i}$  of the set  $T_k^i$ . Since  $u_k^i$  is a local minimum point of  $\mathcal{E}_i$  relative to  $T_k^i$  (see Proposition 10.3), then  $u_k^i$  is a critical point of the functional  $\mathcal{I}_k^i := \mathcal{E}_i + \zeta_{T_k^i}$  in the sense of Szulkin, see Proposition 1.3, i.e.,

$$0 \in \mathcal{E}'_i(u_k^i) + \partial\zeta_{T_k^i}(u_k^i) \quad \text{in} \quad (H_{G_{d,i}^{\tau_i}}(\mathbb{S}^d))^*, \quad (10.16)$$

where  $\partial\zeta_{T_k^i}$  stands for the subdifferential of the convex function  $\zeta_{T_k^i}$ .

Since  $\mathcal{E}$  is even, by means of (10.5) one can easily check that it is  $G_{d,i}^{\tau_i}$ -invariant. The function  $\zeta_{T_k}$  is also  $G_{d,i}^{\tau_i}$ -invariant since  $\tilde{g}T_k \subseteq T_k$  for every  $\tilde{g} \in G_{d,i}^{\tau_i}$  (we use again (10.5)). Finally, since  $G_{d,i}^{\tau_i} \subset O(d+1)$  is compact, and  $\mathcal{E}_i$  and  $\zeta_{T_k^i}$  are the restrictions of  $\mathcal{E}$  and  $\zeta_{T_k}$  to  $H_{G_{d,i}^{\tau_i}}(\mathbb{S}^d)$ , respectively, we may apply – via relation (10.16) – the principle of symmetric criticality, see Theorem 1.31. Thus, we obtain

$$0 \in \mathcal{E}'(u_k^i) + \partial\zeta_{T_k}(u_k^i) \quad \text{in} \quad (H_1^2(\mathbb{S}^d))^*.$$

Consequently, for every  $w \in H_1^2(\mathbb{S}^d)$ , we have

$$\langle \mathcal{E}'(u_k^i), w - u_k^i \rangle_{H_1^2} + \zeta_{T_k}(w) - \zeta_{T_k}(u_k^i) \geq 0,$$

which implies our claim.

*Step 2.* (Proof concluded)  $u_k^i$  is a weak solution of (P).

By Step 1, for all  $w \in T_k$ , we have

$$\int_{\mathbb{S}^d} \langle \nabla u_k^i, \nabla(w - u_k^i) \rangle_h d\sigma_h + c_0 \int_{\mathbb{S}^d} u_k^i(w - u_k^i) d\sigma_h - \int_{\mathbb{S}^d} \psi(u_k^i)(w - u_k^i) d\sigma_h \geq 0.$$

Recall from (10.9) that  $\psi(s) = \tilde{f}(s) + c_0 s$ ,  $s \in \mathbb{R}$ . Moreover,  $f$  and  $\tilde{f}$  coincide on  $[-\tilde{s}, \tilde{s}]$  and  $u_k^i(\sigma) \in [-a_k, a_k] \subset (-\tilde{s}, \tilde{s})$  for a.e.  $\sigma \in \mathbb{S}^d$  (see Proposition 10.4). Consequently, the above inequality reduces to

$$\int_{\mathbb{S}^d} \langle \nabla u_k^i, \nabla(w - u_k^i) \rangle_h d\sigma_h - \int_{\mathbb{S}^d} f(u_k^i)(w - u_k^i) d\sigma_h \geq 0, \quad \forall w \in T_k. \quad (10.17)$$

Let us define the function  $\gamma(s) = \text{sgn}(s) \min(|s|, b_k)$ , and fix  $\varepsilon > 0$  and  $v \in H_1^2(\mathbb{S}^d)$  arbitrarily. Since  $\gamma$  is Lipschitz continuous,  $w_k = \gamma \circ (u_k^i + \varepsilon v)$  belongs to  $H_1^2(\mathbb{S}^d)$ , see Hebey [138, Proposition 2.5, p. 24]. The explicit

expression of  $w_k$  is

$$w_k(\sigma) = \begin{cases} -b_k, & \text{if } \sigma \in \{u_k^i + \varepsilon v < -b_k\} \\ u_k^i(\sigma) + \varepsilon v(\sigma), & \text{if } \sigma \in \{-b_k \leq u_k^i + \varepsilon v < b_k\} \\ b_k, & \text{if } \sigma \in \{b_k \leq u_k^i + \varepsilon v\}. \end{cases}$$

Therefore,  $w_k \in T_k$ . Taking  $w = w_k$  as a test function in (10.17), we obtain

$$\begin{aligned} 0 &\leq - \int_{\{u_k^i + \varepsilon v < -b_k\}} |\nabla u_k^i|_h^2 + \int_{\{u_k^i + \varepsilon v < -b_k\}} f(u_k^i)(b_k + u_k^i) \\ &\quad + \varepsilon \int_{\{-b_k \leq u_k^i + \varepsilon v < b_k\}} \langle \nabla u_k^i, \nabla v \rangle_h - \varepsilon \int_{\{-b_k \leq u_k^i + \varepsilon v < b_k\}} f(u_k^i)v \\ &\quad - \int_{\{b_k \leq u_k^i + \varepsilon v\}} |\nabla u_k^i|_h^2 - \int_{\{b_k \leq u_k^i + \varepsilon v\}} f(u_k^i)(b_k - u_k^i). \end{aligned}$$

After a suitable rearrangement of the terms in this inequality, we obtain that

$$\begin{aligned} 0 &\leq \varepsilon \int_{\mathbb{S}^d} \langle \nabla u_k^i, \nabla v \rangle_h - \varepsilon \int_{\mathbb{S}^d} f(u_k^i)v \\ &\quad - \int_{\{u_k^i + \varepsilon v < -b_k\}} |\nabla u_k^i|_h^2 - \int_{\{b_k \leq u_k^i + \varepsilon v\}} |\nabla u_k^i|_h^2 \\ &\quad + \int_{\{u_k^i + \varepsilon v < -b_k\}} f(u_k^i)(b_k + u_k^i + \varepsilon v) + \int_{\{b_k \leq u_k^i + \varepsilon v\}} f(u_k^i)(-b_k + u_k^i + \varepsilon v) \\ &\quad - \varepsilon \int_{\{u_k^i + \varepsilon v < -b_k\}} \langle \nabla u_k^i, \nabla v \rangle_h - \varepsilon \int_{\{b_k \leq u_k^i + \varepsilon v\}} \langle \nabla u_k^i, \nabla v \rangle_h. \end{aligned}$$

Let  $M_k = \max_{[-a_k, a_k]} |f|$ . Since  $u_k^i(\sigma) \in [-a_k, a_k] \subset [-b_k, b_k]$  for a.e.  $\sigma \in \mathbb{S}^d$ , we have

$$\int_{\{u_k^i + \varepsilon v < -b_k\}} f(u_k^i)(b_k + u_k^i + \varepsilon v) \leq -\varepsilon M_k \int_{\{u_k^i + \varepsilon v < -b_k\}} v$$

and

$$\int_{\{b_k \leq u_k^i + \varepsilon v\}} f(u_k^i)(-b_k + u_k^i + \varepsilon v) \leq \varepsilon M_k \int_{\{b_k \leq u_k^i + \varepsilon v\}} v.$$

Using the above estimates and dividing by  $\varepsilon > 0$ , we obtain

$$\begin{aligned} 0 \leq & \int_{\mathbb{S}^d} \langle \nabla u_k^i, \nabla v \rangle_h d\sigma_h - \int_{\mathbb{S}^d} f(u_k^i) v d\sigma_h \\ & - M_k \int_{\{u_k^i + \varepsilon v < -b_k\}} v d\sigma_h + M_k \int_{\{b_k \leq u_k^i + \varepsilon v\}} v d\sigma_h \\ & - \int_{\{u_k^i + \varepsilon v < -b_k\}} \langle \nabla u_k^i, \nabla v \rangle_h d\sigma_h - \int_{\{b_k \leq u_k^i + \varepsilon v\}} \langle \nabla u_k^i, \nabla v \rangle_h d\sigma_h. \end{aligned}$$

Now, letting  $\varepsilon \rightarrow 0^+$ , and taking into account Proposition 10.4 (i.e.,  $-a_k \leq u_k^i(\sigma) \leq a_k$  for a.e.  $\sigma \in \mathbb{S}^d$ ), we have

$$\text{meas}(\{u_k^i + \varepsilon v < -b_k\}) \rightarrow 0 \quad \text{and} \quad \text{meas}(\{b_k \leq u_k^i + \varepsilon v\}) \rightarrow 0,$$

respectively. Consequently, the above inequality reduces to

$$0 \leq \int_{\mathbb{S}^d} \langle \nabla u_k^i, \nabla v \rangle_h d\sigma_h - \int_{\mathbb{S}^d} f(u_k^i) v d\sigma_h.$$

Putting  $(-v)$  instead of  $v$ , we see that  $u_k^i$  is a weak solution of (P), which completes the proof.  $\square$

*Proof of Theorem 10.2.* Fix  $i \in \{1, \dots, s_d\}$ . Combining Propositions 10.3 and 10.5, one can see that there are infinitely many distinct elements in the sequence  $\{u_k^i\}_k$ . These elements are weak solutions of (P) as Proposition 10.6 shows, and they change sign, see Remark 10.1. Moreover, due to Theorem 10.1 (ii), solutions in different spaces  $H_{G_{d,i}^{\tau_i}}(\mathbb{S}^d)$ ,  $i \in \{1, \dots, s_d\}$ , cannot be compared from symmetrical point of view. The  $L^\infty$ - and  $H_1^2$ -asymptotic behaviour of the sequences of solutions are described in Proposition 10.5.  $\square$

### 10.4 Arbitrarily large solutions

Instead of  $(f_1^0)$  and  $(f_2^0)$ , respectively, we assume

$$\begin{aligned} (f_1^\infty). \quad & -\infty < \liminf_{s \rightarrow \infty} \frac{F(s)}{s^2} \leq \limsup_{s \rightarrow \infty} \frac{F(s)}{s^2} = +\infty; \\ (f_2^\infty). \quad & \liminf_{s \rightarrow \infty} \frac{f(s)}{s} < 0. \end{aligned}$$

Unlike in Theorem 10.2 where no further assumption is needed at infinity, we have to control here the growth of  $f$ . We assume that  $f$  has an *asymptotically critical* growth at infinity, namely,

$$(f_3^\infty). \quad \sup_{s \in \mathbb{R}} \frac{|f(s)|}{1+|s|^{2^*-1}} < \infty, \quad \text{where } 2^* = \frac{2d}{d-2}.$$

**Theorem 10.3** *Let  $d \geq 5$  and  $f : \mathbb{R} \rightarrow \mathbb{R}$  be an odd, continuous function which verifies  $(f_1^\infty)$ ,  $(f_2^\infty)$  and  $(f_3^\infty)$ . Then there exist at least  $s_d = [d/2] + (-1)^{d+1} - 1$  sequences  $\{\tilde{u}_k^i\}_k \subset H_1^2(\mathbb{S}^d)$ ,  $i \in \{1, \dots, s_d\}$ , of sign-changing weak solutions of (P) distinguished by their symmetry properties. In addition,*

$$\lim_{k \rightarrow \infty} \|\tilde{u}_k^i\|_{L^\infty} = \lim_{k \rightarrow \infty} \|\tilde{u}_k^i\|_{H_1^2} = \infty \text{ for every } i \in \{1, \dots, s_d\}.$$

**Example 10.2** Let  $d \geq 5$  and  $\alpha, \beta, \gamma \in \mathbb{R}$  such that  $\frac{d+2}{d-2} \geq \alpha > 1$ ,  $|\alpha - \beta| < 1$ , and  $\gamma \in (0, 1)$ . Then, the function  $f : \mathbb{R} \rightarrow \mathbb{R}$  defined by  $f(s) = |s|^{\alpha-1}s(\gamma + \sin |s|^\beta)$  verifies the hypotheses  $(f_1^\infty)$ ,  $(f_2^\infty)$  and  $(f_3^\infty)$ , respectively.

Certain parts of the proof of Theorem 10.3 are similar to that of Theorem 10.2; so, we present only the differences. We assume throughout of this section that the hypotheses of Theorem 10.3 are fulfilled. Due to  $(f_2^\infty)$ , one can fix  $c_\infty > 0$  such that

$$\liminf_{s \rightarrow \infty} \frac{f(s)}{s} < -c_\infty < 0.$$

Let  $\{\bar{s}_k\} \subset (0, \infty)$  be a sequence converging (increasingly) to  $+\infty$ , such that  $f(\bar{s}_k) < -c_\infty \bar{s}_k$ . We define the functions

$$\psi(s) = f(s) + c_\infty s \text{ and } \Psi(s) = \int_0^s \psi(t)dt = F(s) + \frac{c_\infty}{2}s^2, \quad s \in \mathbb{R}. \tag{10.18}$$

By construction,  $\psi(\bar{s}_k) < 0$ ; consequently, there are two sequences  $\{a_k\}_k, \{b_k\}_k \subset (0, \infty)$ , both converging to  $\infty$ , such that  $a_k < \bar{s}_k < b_k < a_{k+1}$  for every  $k \in \mathbb{N}$  and

$$\psi(s) \leq 0 \text{ for every } s \in [a_k, b_k]. \tag{10.19}$$

Since  $c_\infty > 0$ , the norm  $\|\cdot\|_{c_\infty}$  defined in the same way as (10.11) with  $c_\infty$  instead of  $c_0$ , is equivalent to the standard norm  $\|\cdot\|_{H_1^2}$ . Now, we define the energy functional  $\mathcal{E} : H_1^2(\mathbb{S}^d) \rightarrow \mathbb{R}$  associated with (P) by

$$\mathcal{E}(u) = \frac{1}{2}\|u\|_{c_\infty}^2 - \int_{\mathbb{S}^d} \Psi(u(\sigma))d\sigma_h.$$

Since  $H_1^2(\mathbb{S}^d)$  is continuously embedded into  $L^p(\mathbb{S}^d)$ ,  $1 \leq p \leq 2^*$ , see [138, Corollary 2.1, p. 33], using hypothesis  $(f_3^\infty)$ , the functional  $\mathcal{E}$  is well-defined, and it belongs to  $C^1(H_1^2(\mathbb{S}^d))$ . Moreover, since  $f$  is odd on the whole  $\mathbb{R}$ , the functional  $\mathcal{E}$  is even.

We fix  $i \in \{1, \dots, s_d\}$  and the subspace  $H_{G_{d,i}^{\tau_i}}(\mathbb{S}^d)$  of  $H_1^2(\mathbb{S}^d)$ . Let  $\mathcal{E}_i$  be the restriction of the functional  $\mathcal{E}$  to  $H_{G_{d,i}^{\tau_i}}(\mathbb{S}^d)$  and for every  $k \in \mathbb{N}$ , define the set

$$Z_k^i = \{u \in H_{G_{d,i}^{\tau_i}}(\mathbb{S}^d) : \|u\|_{L^\infty} \leq b_k\},$$

where  $b_k$  is from (10.19).

**Proposition 10.7** *The functional  $\mathcal{E}_i$  is bounded from below on  $Z_k^i$  and its infimum  $\tilde{m}_k^i$  on  $Z_k^i$  is attained at  $\tilde{u}_k^i \in Z_k^i$ . Moreover,  $\lim_{k \rightarrow \infty} \tilde{m}_k^i = -\infty$ .*

*Proof.* It is easy to check that  $\mathcal{E}_i$  is bounded from below on  $Z_k^i$ . In order to see that it attains its infimum on  $Z_k^i$  we show that the function  $u \mapsto \int_{\mathbb{S}^d} \Psi(u(\sigma)) d\sigma_h$ ,  $u \in H_{G_{d,i}^{\tau_i}}(\mathbb{S}^d)$  is sequentially weak continuous; in such case,  $\mathcal{E}_i$  is sequentially weak lower semicontinuous and we may proceed in the standard way. On one hand, due to  $(f_3^\infty)$ , (10.18) and the oddness of  $\psi$ , one can find  $c_1 > 0$  such that

$$|\psi(s)| \leq c_1(1 + |s|^{2^*-1}), \quad s \in \mathbb{R}. \tag{10.20}$$

On the other hand, the definition of  $G_{d,i}$  shows that the  $G_{d,i}$ -orbit of every point  $\sigma \in \mathbb{S}^d$  has at least dimension 1, i.e.,  $\dim(G_{d,i}\sigma) \geq 1$  for every  $\sigma \in \mathbb{S}^d$ . Thus

$$d_G = \min\{\dim(G_{d,i}\sigma) : \sigma \in \mathbb{S}^d\} \geq 1.$$

Applying [31, Lemma 3.2], we conclude in particular that  $H_{G_{d,i}^{\tau_i}}(\mathbb{S}^d)$  is compactly embedded into  $L^q(\mathbb{S}^d)$ , whenever  $q \in \left[1, \frac{2d-2}{d-3}\right)$ . Since  $\frac{2d-2}{d-3} > 2^*$ , the embedding  $H_{G_{d,i}^{\tau_i}}(\mathbb{S}^d) \subset H_{G_{d,i}}(\mathbb{S}^d) \hookrightarrow L^{2^*}(\mathbb{S}^d)$  is compact. Combining (10.20) with the above compactness property, we conclude the sequentially weak continuity of the function  $u \mapsto \int_{\mathbb{S}^d} \Psi(u(\sigma)) d\sigma_h$ ,  $u \in H_{G_{d,i}^{\tau_i}}(\mathbb{S}^d)$ . Consequently, we may assert that the infimum  $\tilde{m}_k^i$  on  $Z_k^i$  is attained at the point  $\tilde{u}_k^i \in Z_k^i$ .

We will prove  $\lim_{k \rightarrow \infty} \tilde{m}_k^i = -\infty$ . First, due to (10.18) and  $(f_1^\infty)$ , we have

$$-\infty < \liminf_{s \rightarrow \infty} \frac{\Psi(s)}{s^2} \leq \limsup_{s \rightarrow \infty} \frac{\Psi(s)}{s^2} = +\infty. \tag{10.21}$$

The left inequality of (10.21) and the evenness of  $\Psi$  implies the existence of  $l, \varrho > 0$  such that

$$\Psi(s) \geq -ls^2 \text{ for every } |s| > \varrho. \tag{10.22}$$

Let  $D_i \subset \mathbb{S}^d$  and  $C_i > 0$  be from Proposition 10.2 (which depend only on  $G_{d,i}$  and  $\tau_i$ ), and fix a number  $\bar{l} > 0$  large enough such that

$$\bar{l}\text{Vol}_h(D_i) > \left(\bar{l} + \frac{c_\infty}{2}\right) \text{Vol}_h(\mathbb{S}^d) + \frac{C_i^2}{2}. \quad (10.23)$$

Taking into account the right-hand side of (10.21), there is a sequence  $\{\tilde{s}_k\}_k \subset (0, \infty)$  such that  $\lim_{k \rightarrow \infty} \tilde{s}_k = \infty$  and  $\Psi(\tilde{s}_k) = \Psi(-\tilde{s}_k) > \bar{l}\tilde{s}_k^2$  for every  $k \in \mathbb{N}$ .

Let  $\{b_{n_k}\}_k$  be an increasing subsequence of  $\{b_k\}_k$  such that  $\tilde{s}_k \leq b_{n_k}$  for every  $k \in \mathbb{N}$ . Let  $\tilde{w}_k := w_{\tilde{s}_k} \in H_{G_{d,i}^{\tau_i}}(\mathbb{S}^d)$  be the function from Proposition 10.2 corresponding to the value  $\tilde{s}_k > 0$ . Then  $\tilde{w}_k \in Z_{n_k}^i$  and one has

$$\begin{aligned} \mathcal{E}_i(\tilde{w}_k) &= \frac{1}{2} \|\tilde{w}_k\|_{c_\infty}^2 - \int_{\mathbb{S}^d} \Psi(\tilde{w}_k(\sigma)) d\sigma_h \\ &\leq \frac{1}{2} (C_i^2 + c_\infty \text{Vol}_h(\mathbb{S}^d)) \tilde{s}_k^2 - \int_{D_i} \Psi(\tilde{w}_k(\sigma)) d\sigma_h \\ &\quad - \int_{\mathbb{S}^d \setminus D_i} \Psi(\tilde{w}_k(\sigma)) d\sigma_h. \end{aligned}$$

On account of Proposition 10.2 iii), we have

$$\int_{D_i} \Psi(\tilde{w}_k(\sigma)) d\sigma_h = \Psi(\tilde{s}_k) \text{Vol}_h(D_i) > \bar{l} \text{Vol}_h(D_i) \tilde{s}_k^2.$$

Due to Proposition 10.2 i) and (10.22), we have

$$\begin{aligned} \int_{\mathbb{S}^d \setminus D_i} \Psi(\tilde{w}_k(\sigma)) d\sigma_h &= \int_{(\mathbb{S}^d \setminus D_i) \cap \{|\tilde{w}_k| \leq \varrho\}} \Psi(\tilde{w}_k(\sigma)) d\sigma_h \\ &\quad + \int_{(\mathbb{S}^d \setminus D_i) \cap \{|\tilde{w}_k| > \varrho\}} \Psi(\tilde{w}_k(\sigma)) d\sigma_h \\ &\geq - \left( \max_{[-\varrho, \varrho]} |\Psi| + \bar{l} \tilde{s}_k^2 \right) \text{Vol}_h(\mathbb{S}^d). \end{aligned}$$

Combining these estimates, we obtain that

$$\mathcal{E}_i(\tilde{w}_k) \leq \tilde{s}_k^2 \left( -\bar{l} \text{Vol}_h(D_i) + \left(\bar{l} + \frac{c_\infty}{2}\right) \text{Vol}_h(\mathbb{S}^d) + \frac{C_i^2}{2} \right) + \max_{[-\varrho, \varrho]} |\Psi| \text{Vol}_h(\mathbb{S}^d).$$

Taking into account (10.23) and  $\lim_{k \rightarrow \infty} \tilde{s}_k = \infty$ , we have  $\lim_{k \rightarrow \infty} \mathcal{E}_i(\tilde{w}_k) = -\infty$ . Since  $\tilde{m}_{n_k}^i = \mathcal{E}_i(\tilde{w}_{n_k}^i) = \inf_{Z_{n_k}^i} \mathcal{E}_i \leq \mathcal{E}_i(\tilde{w}_k)$ , then  $\lim_{k \rightarrow \infty} \tilde{m}_{n_k}^i = -\infty$ . Since the sequence  $\{\tilde{m}_k^i\}_k$  is non-increasing, the claim follows.  $\square$

**Proposition 10.8**  $\lim_{k \rightarrow \infty} \|\tilde{u}_k^i\|_{L^\infty} = \lim_{k \rightarrow \infty} \|\tilde{u}_k^i\|_{H_1^2} = \infty$ .

*Proof.* Assume first by contradiction that there exists a subsequence  $\{\tilde{u}_{n_k}^i\}_k$  of  $\{\tilde{u}_k^i\}_k$  such that  $\|\tilde{u}_{n_k}^i\|_{L^\infty} \leq M$  for some  $M > 0$ . In particular,  $\{\tilde{u}_{n_k}^i\} \subset Z_l^i$  for some  $l \in \mathbb{N}$ . Therefore, for every  $n_k \geq l$ , we have

$$\tilde{m}_l^i \geq \tilde{m}_{n_k}^i = \inf_{Z_{n_k}^i} \mathcal{E}_i = \mathcal{E}_i(\tilde{u}_{n_k}^i) \geq \inf_{Z_l^i} \mathcal{E}_i = \tilde{m}_l^i.$$

Consequently,  $\tilde{m}_{n_k} = \tilde{m}_l$  for every  $n_k \geq l$ , and since the sequence  $\{\tilde{m}_k^i\}_k$  is non-increasing, there exists  $k_0 \in \mathbb{N}$  such that for every  $k \geq k_0$  we have  $\tilde{m}_k^i = \tilde{m}_l^i$ , contradicting Proposition 10.7.

It remains to prove that  $\lim_{k \rightarrow \infty} \|\tilde{u}_k^i\|_{H_1^2} = \infty$ . Note that (10.20) and the continuity of the embedding  $H_1^2(\mathbb{S}^d)$  into  $L^{2^*}(\mathbb{S}^d)$  implies that for some  $C > 0$  we have

$$\left| \int_{\mathbb{S}^d} \Psi(u(\sigma)) d\sigma_h \right| \leq C(\|u\|_{H_1^2} + \|u\|_{H_1^{2^*}}), \quad \forall u \in H_1^2(\mathbb{S}^d).$$

Similarly as above, we assume that there exists a subsequence  $\{\tilde{u}_{n_k}^i\}_k$  of  $\{\tilde{u}_k^i\}_k$  such that for some  $M > 0$ , we have  $\|\tilde{u}_{n_k}^i\|_{H_1^2} \leq M$ . Since  $\|\cdot\|_{c_\infty}$  is equivalent with  $\|\cdot\|_{H_1^2}$ , due to the above inequality, the sequence  $\{\mathcal{E}_i(\tilde{u}_{n_k}^i)\}_k$  is bounded. But  $\tilde{m}_{n_k}^i = \mathcal{E}_i(\tilde{u}_{n_k}^i)$ , thus, the sequence  $\{\tilde{m}_{n_k}^i\}_k$  is also bounded. This fact contradicts Proposition 10.7.  $\square$

*Proof of Theorem 10.3.* Due to Proposition 10.7, we can find infinitely many distinct elements  $\tilde{u}_k^i$ ; similar reasoning as in Propositions 10.4 and 10.6 show that  $\tilde{u}_k^i$  are weak solutions of (P) for every  $k \in \mathbb{N}$ . The  $L^\infty$ - and  $H_1^2$ -asymptotic behaviour of the sequences of solutions are described in Proposition 10.8. The rest is similar as in Theorem 10.2.  $\square$

**Remark 10.2** Theorems 10.2 and 10.3 can be successfully applied to treat Emden-Fowler equations of the form

$$-\Delta v = |x|^{\alpha-2} f(|x|^{-\alpha} v), \quad x \in \mathbb{R}^{d+1} \setminus \{0\} \quad (\alpha < 0) \quad (10.24)$$

whenever  $f : \mathbb{R} \rightarrow \mathbb{R}$  is enough smooth and oscillates either at zero or at infinity having an asymptotically critical growth. Finding solutions of (10.24) in the form  $v(x) = v(r, \sigma) = r^\alpha u(\sigma)$ ,  $(r, \sigma) = (|x|, x/|x|) \in (0, \infty) \times \mathbb{S}^d$  being the spherical coordinates, we obtain

$$-\Delta_h u + \alpha(1 - d - \alpha)u = f(u) \quad \text{on } \mathbb{S}^d, \quad (10.25)$$

see also (9.1) $_\lambda$ . Assuming  $(f_1^L)$  and  $\liminf_{s \rightarrow L} \frac{f(s)}{s} < \alpha(1 - d - \alpha)$  with

$L \in \{0^+, +\infty\}$ , and  $(f_3^\infty)$  whenever  $L = +\infty$ , we may formulate multiplicity results for (10.25), so for (10.24). Note that the obtained solutions of (10.24) are sign-changing and *non*-radial.

### 10.5 Historical notes, comments and perspectives

*A. Historical notes.* Elliptic problems involving oscillatory nonlinearities have been studied in Omari and Zanolin [232], Ricceri [262], Saint Raymond [269], subjected to standard Neumann or Dirichlet boundary value conditions on bounded open domains of  $\mathbb{R}^n$ , or even on unbounded domains, see Faraci and Kristály [113], Kristály [168]. Results in finding sign-changing solutions for semilinear problems can be found in Li and Wang [190], Zou [306] and references therein. The strategy in these last papers is to construct suitable closed convex sets which contain all the positive and negative solutions in the interior, and are invariant with respect to some vector fields. Our approach is rather different than those of [190], [306] and is related to the works of Bartsch, Schneider and Weth [31], Bartsch and Willem [33], where the existence of non-radial and sign-changing solutions are studied for Schrödinger and polyharmonic equations defined on  $\mathbb{R}^n$ .

*B. Comments.* The minimal number of those sequences of solutions for (P) which contain mutually symmetrically distinct elements is  $s_d = [d/2] + (-1)^{d+1} - 1$ . Note that  $s_d \sim d/2$  as  $d \rightarrow \infty$ . However, in lower dimensions, Theorems 10.2 and 10.3 are not spectacular. For instance,  $s_4 = 0$ ; therefore, on  $\mathbb{S}^4$  we have no analogous results as Theorems 10.2 and 10.3. Note that  $s_3 = 1$ ; in fact, for  $G_{3,1} = O(2) \times O(2)$  we may apply our arguments. Hence, on  $\mathbb{S}^3$  one can find a sequence of solutions of (P) with the described properties in our theorems. We may compare these results with that of Bartsch and Willem [33]; they studied the lower bound of those sequences of solutions for a Schrödinger equation on  $\mathbb{R}^{d+1}$  which contain elements in different  $O(d+1)$ -orbits. Due to [33, Proposition 4.1, p. 457], we deduce that their lower bound is  $s'_d = [\log_2 \frac{d+3}{3}]$  whenever  $d \geq 3$  and  $d \neq 4$ .

Let  $\alpha, \beta \in L^\infty(\mathbb{S}^d)$  be two  $G_{d,i}$ -invariant functions such that  $\text{essinf}_{\mathbb{S}^d} \beta > 0$  and consider the problem

$$-\Delta_h u + \alpha(\sigma)u = \beta(\sigma)f(u) \quad \text{on } \mathbb{S}^d. \quad (10.26)$$

If  $f : \mathbb{R} \rightarrow \mathbb{R}$  has an asymptotically critical growth fulfilling  $(f_1^L)$  and

$$\liminf_{s \rightarrow L} \frac{f(s)}{s} < \operatorname{ess\,inf}_{\mathbb{S}^d} \frac{\alpha}{\beta},$$

problem (10.26) admits a sequence of  $G_{d,i}$ -invariant (perhaps not sign-changing) weak solutions in both cases, i.e.  $L \in \{0^+, \infty\}$ . The proofs can be carried out following Theorems 10.2 and 10.3, respectively, considering instead of  $H_{G_{d,i}^{\tau_i}}(\mathbb{S}^d)$  the space  $H_{G_{d,i}}(\mathbb{S}^d)$ . Note that  $\alpha : \mathbb{S}^d \rightarrow \mathbb{R}$  may change its sign. In particular, this type of result complements the paper of Cotsoolis and Iliopoulos [79].

The main results of this chapter are mainly contained in the papers of Kristály [171], and Kristály and Marzantowicz [174].

*C. Further perspectives.* The symmetry and compactness of the sphere  $\mathbb{S}^d$  have been deeply exploited in our framework. A challenging problem related to (P) may be formulated on *non-compact Riemannian symmetric spaces* (for instance, on the hyperbolic space  $\mathbb{H}^d = SO_0(d, 1)/SO(d)$  which is the dual companion of  $\mathbb{S}^d = SO(d + 1)/SO(d)$ ). In order to handle this kind of problem, the action of the isometry group of the symmetric space seems to be essential, as shown by Hebey [138, Chapter 9], Hebey and Vaugon [140].

# 11

## Equations with Critical Exponent

Numbers are the highest degree of knowledge. It is knowledge itself.

---

Plato (429–347 B.C.)

In this chapter we treat elliptic equations with critical exponent defined on compact Riemannian manifolds. This problem is closely related to the so-called *Yamabe problem*, i.e., for any smooth compact Riemannian manifold  $(M, g)$  of dimension  $d \geq 3$ , there exists a conformal metric  $\tilde{g}$  to  $g$  with constant scalar curvature.

### 11.1 Introduction

The starting point of this chapter is the *Yamabe problem*, whose geometric formulation is as follows:

*For any smooth compact Riemannian manifold  $(M, g)$  of dimension  $d \geq 3$ , there exists a conformal metric to  $g$  with constant scalar curvature.*

This problem can be transposed into an elliptic equation which involves the critical exponent. In the sequel, we focus our attention to this transformation as well as to establish some existence results.

Let  $(M, g)$  be a  $d$ -dimensional Riemannian manifold and  $\nabla$  its Levi-Civita (or, Riemannian) connection. Let  $(p_i)$  be a local system of coordinates on  $M$ , then

$$\nabla_{\frac{\partial}{\partial p_k}} \left( \frac{\partial}{\partial p_j} \right) = \Gamma_{ij}^k \frac{\partial}{\partial p_k}.$$

Here and in the sequel, we use the Einstein convention for the summa-

tion. The functions  $\Gamma_{ij}^k$  are called the Christoffel symbols of the Levi-Civita connection  $\nabla$ . In the local system of coordinates  $(p_i)$  the components of metric tensor  $g$  we denote by  $(g_{ij})$  with the inverse matrix  $(g^{ij})$ , and let  $|g| = \det(g_{ij})$ . The divergence operator  $\operatorname{div}_g$  on the  $C^1$  vector field  $X = (X^i)$  is defined by

$$\operatorname{div}_g X = \frac{1}{\sqrt{|g|}} \frac{\partial}{\partial p_i} (\sqrt{|g|} X^i),$$

and the Laplace–Beltrami operator by  $\Delta_g u = \operatorname{div}_g(\nabla u)$ . Here

$$\nabla u = g^{ij} \frac{\partial u}{\partial p_i}$$

In the local system of coordinates  $(p_i)$  the Laplace–Beltrami operator  $\Delta_g$ , has the following expression:

$$\Delta_g u = g^{ij} \left( \frac{\partial^2 u}{\partial p_k \partial p_j} - \Gamma_{ij}^k \frac{\partial u}{\partial p_k} \right). \quad (11.1)$$

The *curvature* associated to the connection  $\nabla$  is defined by:

$$R(X, Y) = \nabla_X \nabla_Y - \nabla_Y \nabla_X - \nabla_{[X, Y]},$$

where  $[\cdot, \cdot]$  denote the Poisson bracket of the vector fields  $X$  and  $Y$ . Consider the 4-covariant tensor  $R(X, Y, Z, T) = g(X, R(Z, T)Y)$  with components  $R_{lkij} = g_{lm} R_{kij}^m$ . The Ricci tensor is obtained from curvature tensor by contraction and its components are  $R_{ij} = R_{ikj}^k$ . The Ricci tensor is symmetric and its contraction  $S_g = R_{ij} g^{ij}$  is called the *scalar curvature*. A metric of the form  $\tilde{g} = e^u g$  is said to be *conformal metric* to  $g$ . We denote by  $[g]$  the conformal class of the metric  $g$ . By definition

$$[g] = \{e^u g; u \in C^\infty(M)\}. \quad (11.2)$$

If  $S_g$  and  $S_{\tilde{g}}$  are the scalar curvatures of  $g$  and  $\tilde{g}$ , and  $\tilde{g} \in [g]$ , one gets that

$$e^{2u} S_{\tilde{g}} = S_g - 2(d-1)\Delta_g u - (d-1)(d-2)|\nabla u|_g^2. \quad (11.3)$$

Let us now write  $\tilde{g}$  under the form  $\tilde{g} = u^{\frac{4}{d-2}} g$  where  $u : M \rightarrow \mathbb{R}$  is a smooth positive function. The above relation becomes

$$-\Delta_g u + \frac{d-2}{4(d-1)} S_g u = \frac{d-2}{4(d-1)} S_{\tilde{g}} u^{\frac{d+2}{d-2}}. \quad (11.4)$$

The geometric version of Yamabe's problem can be formulated in language of PDE's as follows:

For any smooth compact Riemannian manifold  $(M, g)$  of dimension  $d \geq 3$ , there exists  $u \in C^\infty(M)$ ,  $u > 0$  and there exists  $\lambda \in \mathbb{R}$  such that

$$-\Delta_g u + \frac{d-2}{4(d-1)} S_g u = \lambda u^{\frac{d+2}{d-2}}. \quad (11.5)$$

If  $u$  and  $\lambda$  satisfies equation (11.5), and if  $\tilde{g} = u^{\frac{4}{d-2}} g$ , then we obtain

$$S_{\tilde{g}} = \frac{4(d-1)}{d-2} \lambda.$$

In particular, this gives a conformal metric to  $g$  of constant scalar curvature. The left hand side in this equation (11.5) is referred to as the *conformal Laplacian*. We denoted by

$$L_g u = -\Delta_g u + \frac{d-2}{4(d-1)} S_g u.$$

Note that  $L_g$  is conformally invariant in the following sense: If  $\tilde{g} = \varphi^{\frac{4}{d-2}} g$  is a conformal metric to  $g$ , then, for all  $u \in C^\infty(M)$ ,

$$L_{\tilde{g}}(u) = \varphi^{-\frac{d+2}{d-2}} L_g(u\varphi).$$

Taking into account (11.4) and (11.5), in this chapter we study the problem

$$(CE) \quad \begin{cases} -\Delta_g u + \alpha u = \lambda u^{2^*-1}, & \text{in } M \\ u > 0, & \text{in } M, \end{cases}$$

where  $(M, g)$  be a smooth compact Riemannian manifold of dimension  $d \geq 3$ ,  $\lambda \in \mathbb{R}$ ,  $\alpha : M \rightarrow \mathbb{R}$  is a smooth function, and  $2^* = \frac{2d}{d-2}$  is the critical exponent of the embedding  $H_1^2(M) \hookrightarrow L^{2^*}(M)$ . To complete this, we first study a subcritical problem in Section 11.2, while in Section 11.3 we deal with problem (CE).

## 11.2 Subcritical case

Let  $(M, g)$  be a smooth compact manifold of dimension  $d \geq 3$ ,  $\alpha : M \rightarrow \mathbb{R}$  a smooth function and  $q \in (2, 2^*)$  a fixed number. In this section we study the subcritical equation

$$\begin{cases} -\Delta_g u + \alpha u = \lambda u^{q-1}, & \text{in } M \\ u > 0, & \text{in } M. \end{cases} \quad (11.6)$$

**Definition 11.1** We say that  $u \in H_1^2(M)$  is a weak solution of the equation (11.6) if for every  $\varphi \in H_1^2(M)$  we have

$$\int_M \langle \nabla u, \nabla \varphi \rangle_g d\sigma_g + \int_M \alpha u \varphi d\sigma_g = \lambda \int_M u^{q-1} \varphi d\sigma_g.$$

We have the following regularity result.

**Theorem 11.1** *If  $u \in H_1^2(M)$  is a weak solution of (11.6), then  $u \in C^\infty(M)$ .*

*Proof* First we prove that the weak solution  $u$  is smooth, i.e.  $u \in C^\infty(M)$ . For this let  $f = \mu_q u^{q-1}$ , and  $p_1 = 2^*$ . Since  $u \in H_1^2(M)$ , from Theorem B.9 follows that  $u \in L^{p_1}(M)$ . Hence  $f \in L^{\frac{p_1}{q-1}}(M)$ , and it follows from Theorem B.12 that  $u \in H_2^{\frac{p_1}{q-1}}(M)$ . Using again Theorem B.9 embedding theorem, we obtain

- 1)  $u \in L^{p_2}(M)$ , where  $p_2 = \frac{dp_1}{d(q-1)-2p_1}$  if  $d(q-1) > 2p_1$   
or
- 2)  $u \in L^s(M)$  for all  $s$  if  $d(q-1) \leq 2p_1$ .

If we repeat this process, we get by finite induction that  $u \in L^s(M)$  for all  $s$ . Indeed, let  $p_0 = \frac{d(q-2)}{2}$ . Then  $p_1 > p_0$ . We define  $p_k$  by induction letting

$$\begin{cases} p_{k+1} = \frac{dp_k}{d(q-1)-2p_k}, & \text{if } d(q-1) > 2p_k \\ p_{k+1} = +\infty, & \text{if } d(q-1) \leq 2p_k. \end{cases} \quad (11.7)$$

For every  $k \in \mathbb{N}^*$  we have  $p_k > p_0$ . It follows that  $p_{k+1} > p_k$ . Moreover,  $u \in L^{p_{k+1}}(M)$  if  $d(q-1) > 2p_k$ , and  $u \in L^s(M)$  for all  $s$  if  $d(q-1) \leq 2p_k$ . Now, either exists  $k \in \mathbb{N}^*$  such that  $p_k > \frac{d(q-1)}{2}$ , or  $p_k \leq \frac{d(q-1)}{2}$  for all  $k$ . In the first case,  $p_{k+1} = +\infty$  and we get that  $u \in L^s(M)$  for all  $s$ . In the second case,  $(p_k)$  is an increasing sequence bounded from above. Thus  $(p_k)$  converges, and if  $p$  is the limit of the  $p_k$ 's, then

$$p = \frac{dp}{d(q-1) - 2p}$$

so that  $p = \frac{d(q-2)}{2}$ , which is impossible. This prove that  $u \in L^s(M)$  for all  $s$ . By Theorem B.12 we get  $u \in H_2^s(M)$  for all  $s$ . From c) Theorem B.9 follows that  $u \in C^1(M)$ . Then, since  $q > 2$ ,  $u^{q-1} \in C^1(M)$ , and, in particular  $u^{q-1} \in H_1^s(M)$  for all  $s$ . Using again Theorem B.12, it follows that  $u \in H_3^s(M)$  for all  $s$ , and c) Theorem B.9 implies  $u \in C^2(M)$ . Since

$u \neq 0$ , applying the maximum principle, i.e. Theorem B.1 we get than  $u > 0$ . From Theorem B.12 follows that  $u \in C^\infty(M)$ . Thus the assertion of theorem is proved.  $\square$

We define

$$\mu_q = \inf_{u \in \mathcal{H}_q} \int_M (|\nabla u|_g^2 + \alpha u^2) d\sigma_g, \quad (11.8)$$

where

$$\mathcal{H}_q = \{u \in H_1^2 \mid \text{such that } \int_M |u|^q d\sigma_g = 1\}. \quad (11.9)$$

We have the following result

**Theorem 11.2** *Let  $(M, g)$  be a smooth compact Riemannian manifold of dimension  $d \geq 3$ . Given  $q \in (2, 2^*)$ , there exists  $u \in C^\infty(M)$ ,  $u > 0$  in  $M$  such that*

$$-\Delta_g u + \alpha u = \mu_q u^{q-1},$$

where  $\int_M u^q d\sigma_g = 1$ , where  $\mu_q$  is as above.

*Proof* Let  $(u_k) \subset \mathcal{H}_q$  a minimizing sequence for  $\mu_q$ . Taking into account Proposition 3.49 [19] we have  $|\nabla|u|| = |\nabla u|$  a.e., up to replacing  $u_k$  by  $|u_k|$ , we can assume that  $u_k \geq 0$  for all  $k$ . Since  $(u_k) \subset \mathcal{H}_q$  and  $q > 2$ ,  $(u_k)$  using Hölder inequality we get

$$\int_M u_k^2 d\sigma_g \leq \text{Vol}_g(M)^{\frac{q-2}{q}} \left( \int_M |u_k|^q d\sigma_g \right)^{\frac{2}{q}}.$$

From this inequality follows that  $(u_k)$  is bounded in  $L^2(M)$ . In particular,  $\mu_q$  is finite, and  $(u_k)$  is bounded in  $H_1^2(M)$ . Since  $H_1^2(M)$  is a Hilbert space, follows that is reflexive. Taking into account that the embedding  $H_1^2(M) \hookrightarrow L^q(M)$  is compact, there exists  $u \in H_1^2(M)$  and a subsequence of  $(u_k)$  which will denote in the sameway, such that:

- a)  $u_k \rightharpoonup u$  weakly in  $H_1^2(M)$
- b)  $u_k \rightarrow u$  strongly in  $L^q(M)$
- c)  $u_k(\sigma) \rightarrow u(\sigma)$  a.e.  $\sigma \in M$ .

From c) follows that  $u \geq 0$ , and by b),  $u \in \mathcal{H}_q$ . Because the norm is weakly lower semicontinuous and  $u_k \rightharpoonup u$  weakly in  $H_1^2(M)$  follows that

$$\|u\|_{H_1^2} \leq \liminf_{i \rightarrow +\infty} \|u_k\|_{H_1^2}.$$

Because  $u_k \rightarrow u$  strongly in  $L^q(M)$  and since  $L^q(M) \subset L^2(M)$ , then we obtain

$$\mu_q = \int_M (|\nabla u|_g^2 + \alpha u^2) d\sigma_g.$$

In particular,  $u$  is minimizer for  $\mu_q$ . By Theorem 1.13 follows the existence of  $\nu \in \mathbb{R}$ , such that for any  $\varphi \in H_1^2(M)$ ,

$$\int_M \langle \nabla u, \nabla \varphi \rangle_g d\sigma_g + \int_M \alpha u \varphi d\sigma_g = \nu \int_M u^{q-1} \varphi d\sigma_g.$$

Taking  $\varphi = u$ , we get that  $\nu = \mu_q$ . Thus  $u \in \mathcal{H}_q, u \geq 0$ , a weak solutions of (11.6). □

### 11.3 Critical case

In this section we study the critical case, i.e. the problem

$$(CE) \quad \begin{cases} -\Delta_g u + \alpha u = \lambda u^{2^*-1}, & \text{in } M \\ u > 0, & \text{in } M, \end{cases}$$

where,  $(M, g)$  be a smooth compact manifold of dimension  $d \geq 3$ ,  $\alpha : M \rightarrow \mathbb{R}$  a smooth function, where  $\lambda \in \mathbb{R}$  and  $2^* = \frac{2d}{d-2}$  is the critical exponent.

We define

$$\mu = \inf_{u \in \mathcal{H}} \int_M (|\nabla u|_g^2 + \alpha u^2) d\sigma_g,$$

where

$$\mathcal{H} = \{u \in H_1^2 \mid \text{such that } \int_M |u|^{2^*} d\sigma_g = 1\}.$$

Because  $L^{2^*}(M) \subset L^2(M)$ , follows that  $\mu$  is finite. First we state the following regularity result, which essentially is due to Trudinger [289]. Here we follows the Hebey work [139].

We have the following regularity result.

**Theorem 11.3** *Let  $(M, g)$  be a smooth compact Riemannian manifold of dimension  $d \geq 3$ , and let  $\alpha : M \rightarrow \mathbb{R}$  be a smooth function. If  $u \in H_1^2(M), u \geq 0$  is a weak solution of the equation*

$$-\Delta_g u + \alpha u = \lambda u^{2^*-1}$$

where  $\lambda \in \mathbb{R}$ , then  $u \in C^\infty(M)$  and either  $u \equiv 0$ , or  $u > 0$  everywhere.

*Proof* It is enough to prove that  $u \in L^s(M)$  for some  $s > 2^*$ , because using the argument from Theorem 11.2 follows that  $u \in C^\infty(M)$ . For a fixed number  $L > 0$  we consider the following functions  $F_L, G_L : \mathbb{R} \rightarrow \mathbb{R}$  given by

$$F_L(t) = \begin{cases} |t|^{2^*/2}, & \text{if } |t| \leq L \\ \frac{2^*}{2} L^{(2^*-2)/2} |t| - \frac{2^*-2}{2} L^{2^*/2}, & \text{if } |t| > L, \end{cases}$$

and

$$G_L(t) = \begin{cases} |t|^{2^*-1}, & \text{if } |t| \leq L \\ \frac{2^*}{2} L^{(2^*-2)} |t| - \frac{2^*-2}{2} L^{2^*-1}, & \text{if } |t| > L. \end{cases}$$

From the definitions of the functions  $F_L$  and  $G_L$  follows

$$F_L \leq t^{2^*/2}, \quad G_L \leq t^{2^*-1} \quad \text{and} \quad (F_L(t))^2 \geq t G_L(t).$$

From these, follows easily  $(F'_L(t))^2 \leq \frac{2^*}{2} G'_L(t)$  for  $t \neq L$ . Taking into account that  $F_L$  and  $G_L$  are locally Lipschitz and if we denote  $\tilde{F}_L = F_L(u)$  and  $\tilde{G}_L = G_L(u)$  follows that  $\tilde{F}_L, \tilde{G}_L \in H^2_1(M)$ . Since  $u \in H^2_1(M)$  is a weak solutions the equation (CE), follows that

$$\int_M \langle \nabla u, \nabla \tilde{G}_L \rangle_g d\sigma_g + \int_M \alpha u \tilde{G}_L d\sigma_g = \lambda \int_M u^{2^*-1} \tilde{G}_L d\sigma_g.$$

Since  $\tilde{G}_L(u) \leq u^{2^*-1}$ , and  $u \in L^{2^*}(M)$ , it follows that there exists  $C_1, C_2 > 0$ , independent of  $L$ , such that

$$\int_M G'_L(u) |\nabla u|_g^2 d\sigma_g \leq C_1 + C_2 \int_M u^{2^*-1} \tilde{G}_L d\sigma_g$$

and since  $(F'_L(t))^2 \leq \frac{2^*}{2} G'_L(t)$  and  $t G_L(t) \leq (F_L(t))^2$ , follows that

$$\frac{2^*}{2} \int_M |\nabla \tilde{F}_L|_g d\sigma_g \leq C_1 + C_2 \int_M u^{2^*-2} \tilde{F}_L^2 d\sigma_g.$$

Given  $K > 0$ , let

$$K^+ = \{x \in M : u(x) \leq K\},$$

and

$$K^- = \{x \in M : u(x) \geq K\}.$$

Because  $H_1^2(M) \hookrightarrow L^{2^*}(M)$  the embedding is continuous, from the Hölder's inequalities we get

$$\begin{aligned} \int_M u^{2^*-2} \tilde{F}_L^2 d\sigma_g &= \int_{K^-} u^{2^*-2} \tilde{F}_L^2 d\sigma_g + \int_{K^+} u^{2^*-2} \tilde{F}_L^2 d\sigma_g \\ &\leq \int_{K^-} u^{2^*-2} \tilde{F}_L^2 d\sigma_g + \left( \int_{K^+} u^{2^*} \right)^{2/n} \left( \int_{K^+} \tilde{F}_L^{2^*} \right)^{2/2^*} \\ &\leq \int_{K^-} u^{2^*-2} \tilde{F}_L^2 d\sigma_g + \varepsilon(K) \left( \int_M \tilde{F}_L^{2^*} \right)^{2/2^*} \\ &\leq \int_{K^-} u^{2^*-2} \tilde{F}_L^2 d\sigma_g + C_3 \varepsilon(K) \int_M (|\nabla \tilde{F}_L|^2 + \tilde{F}_L^2) d\sigma_g, \end{aligned}$$

where  $\varepsilon(K) = (\int_{K^+} u^{2^*} d\sigma_g)^{2/d}$ , and  $C_3$  does not depend on  $K$  and  $L$ . Since  $u \in L^{2^*}(M)$ , we have  $\lim_{K \rightarrow +\infty} \varepsilon(K) = 0$ . We fix  $K > 0$  such that  $C_2 C_3 \varepsilon(K) < \frac{2}{2^*}$ . We suppose that  $L > K$ , then we have

$$\int_{K^-} u^{2^*-2} \tilde{F}_L^2 d\sigma_g \leq K^{2(2^*-1)} \text{Vol}_g(M),$$

where  $\text{Vol}_g(M)$  denotes the volume of  $M$  with respect the Riemannian metric  $g$ . Since  $F_L(t) \leq t^{2^*/2}$  and  $u \in L^{2^*}(M)$  follows the existence of a real number  $C_4 > 0$ , which does not depend on  $L$  such that

$$\int_M \tilde{F}_L^2 d\sigma_g \leq C_4.$$

In consequence, then there exist  $C_5, C_6 > 0$ , independent of  $L$ , with  $C_6 < 1$ , such that

$$\int_M |\nabla \tilde{F}_L|^2 d\sigma_g \leq C_5 + C_6 \int_M |\nabla \tilde{F}_L|^2 d\sigma_g.$$

From this inequality follows that

$$\int_M |\nabla \tilde{F}_L|^2 d\sigma_g \leq \frac{C_5}{1 - C_6}.$$

Taking into account the Sobolev inequality for the embedding  $H_1^2(M) \hookrightarrow L^{2^*}(M)$  and the above inequality we get

$$\int_M \tilde{F}_L^{2^*} d\sigma_g \leq C_7,$$

where  $C_7$  does not depend on  $L$ . If  $L \rightarrow +\infty$ , follows that  $u \in L^{(2^*)^2/2}(M)$ . But  $(2^*)^2/2 > 2$  and we get the existence of some  $s > 2^*$  such that  $u \in L^s(M)$ . Now we can use the procedure from Theorem 11.2 to prove

that  $u \in C^\infty(M)$ . Now from Theorem B.1 it follows that either  $u \equiv 0$  or  $u > 0$ . □

**Theorem 11.4** *Let  $(M, g)$  be a smooth compact manifold of dimension  $d \geq 3$ , and let  $\alpha : M \rightarrow \mathbb{R}$  be a smooth function. If  $\mu \leq 0$ , then the problem (CE) has a smooth positive solution.*

*Proof* In the following we distinguish two cases.

### 11.3.1 The negative case: $\mu < 0$

Fix  $q \in (2, 2^*)$  and we consider the equation

$$-\Delta_g u + \alpha u = \mu_q u^{q-1},$$

where  $\mu_q$  is defined in (11.2).

From Theorem 11.2, there exists  $u_q \in C^\infty(M)$ ,  $u_q > 0$ , such that

$$-\Delta_g u_q + \alpha u_q = \mu_q u_q^{q-1},$$

and  $\int_M u_q^q d\sigma_g = 1$ . Hence there exists  $u \in \mathcal{H}$  such that  $I(u) < 0$ , where

$$I(u) = \int_M (|\nabla u|_g^2 + \alpha u^2) d\sigma_g.$$

We have

$$\mu_q \leq I\left(\frac{u}{\left(\int_M |u|^q\right)^{1/q}}\right)$$

and that  $\int_M |u|^q d\sigma_g \leq \text{Vol}_g(M)^{1-\frac{q}{2^*}}$ , where  $\text{Vol}_g(M)$  is the volume of  $M$  with respect to  $g$ , we easily get that there exists  $\varepsilon_0 > 0$  such that  $\mu_q \leq \varepsilon_0$  for every  $q \in (2, 2^*)$ . In similar way, we easily get that there exists  $K > 0$  such that  $\mu_q \geq -K$  for all  $q \in (2, 2^*)$ . Hence, there exists  $\varepsilon_0 > 0$  such that

$$-\frac{1}{\varepsilon_0} \leq \mu_q \leq -\varepsilon_0$$

for all  $q \in (2, 2^*)$ . Let  $\sigma_q$  be a point where  $u_q$  is maximum. Then  $-\Delta_g u_q(\sigma_q) \geq 0$ . It follows from the equation satisfied by  $u_q$  that

$$h(\sigma_q)u_q \leq \mu_q u_q^{q-1}(\sigma_q).$$

In particular,  $h(\sigma_q) < 0$ , and

$$u_q^{q-2}(\sigma_q) \leq \frac{1}{\varepsilon_0} \max_{x \in M} |h(x)|,$$

therefore the  $u'_q$ s are uniformly bounded. From Theorem B.12 follows that the sequence  $u'_q$ s are bounded in  $H_2^p$  for all  $p$ . In particular, a subsequence of the  $u'_q$ s converge to some  $u$  in  $C^1(M)$  as  $q \rightarrow 2^*$ . Assuming that the  $\mu'_q$ s converge to some  $\lambda$  as  $q \rightarrow 2^*$ , we get that  $u$  is a weak solution of

$$-\Delta_g u + \alpha u = \lambda u^{2^*-1}.$$

Because  $\int_M u_q^q d\sigma_g = 1$ , follows that  $u$  is nonzero and  $u_q \rightarrow u$  uniformly as  $q \rightarrow 2^*$ .

From Theorem 11.3 follows that  $u$  is smooth and Proposition B.1 implies that  $u$  is everywhere positive. In particular,  $u$  is a strong solution of the above equation. With similar arguments as above, we get that  $\limsup_{q \rightarrow 2^*} \mu_q \leq \mu$ . Independently, it is straightforward that  $\mu \leq$

$I\left(\left(\int_M u_q^{2^*} d\sigma_g\right)^{-\frac{1}{2^*}} u_q\right)$ , therefore

$$\left(\int_M u_q^{2^*} d\sigma_g\right)^{\frac{2}{2^*}} \mu \leq \mu_q$$

for all  $q$ . Since  $u_q \rightarrow u$  uniformly, we have that  $\int_M u_q^{2^*} d\sigma_g \rightarrow \int_M u^{2^*} d\sigma_g$ . Hence, we also have that  $\liminf_{q \rightarrow 2^*} \mu_q \geq \mu$ . It follows that  $\mu_q \rightarrow \mu$  as  $q \rightarrow 2^*$ , so that  $\lambda = \mu$ . Summarizing, we proved that if  $\mu < 0$ , then there exists  $u \in C^\infty(M), u > 0$ , such that

$$-\Delta_g u + \alpha u = \mu u^{2^*-1}$$

and  $\int_M u^{2^*} d\sigma_g = 1$ . In particular,  $u$  is a minimizing solution of the equation. Moreover,  $u$  is obtained as the uniform limit of a subsequence of the  $u'_q$ s.

### 11.3.2 The null case: $\mu = 0$

For the fixed number  $q \in (2, 2^*)$  we consider the equation

$$-\Delta_g u + \alpha u = \mu_q u^{q-1},$$

where  $\mu_q$  is defined in (11.8).

From Theorem 11.2 follows the existence of an element  $u_q \in C^\infty(M), u_q >$

0 be such that

$$-\Delta_g u_q + \alpha u_q = \mu_q u_q^{q-1}$$

and  $\int_M u_q^q d\sigma_g = 1$ . First we claim that if  $\mu = 0$ , then  $\mu_q = 0$  for all  $q$ . Given  $\varepsilon > 0$ , we let  $u_\varepsilon \in \mathcal{H}$  be such that  $I(u_\varepsilon) \leq \varepsilon$ . Thanks to Sobolev inequality, there exists  $A > 0$  such that for any  $u \in H_1^2(M)$ ,

$$\|u\|_{2^*}^2 \leq A(\|u\|_2^2 + \|u\|_2^2).$$

Taking  $u = u_\varepsilon$  in the above inequality, we get that for any  $\varepsilon > 0$ ,

$$1 \leq A(\varepsilon + B\|u_\varepsilon\|_2^2),$$

where  $B = 1 + \max_{x \in M} |h(x)|$ . Hence, there exists  $C > 0$  such that  $\|u_\varepsilon\|_2 \geq C$  for all  $\varepsilon > 0$  sufficiently small. In particular, for  $q > 2$ , there exists  $C_q > 0$  such that

$$\int_M |u_\varepsilon|^q d\sigma_g \geq C_q.$$

Independently, it is clear that  $\mu_q \leq I(\|u_\varepsilon\|_q^{-1} u_\varepsilon)$ , so that  $\|u_\varepsilon\|_q^2 \mu_q \leq \varepsilon$ .

Fixing  $q > 2$ , and letting  $\varepsilon \rightarrow 0$ , it follows that  $\mu_q \leq 0$ . On the other hand,

$$\mu_q = I(u_q) = \|u_q\|_{2^*}^2 I(\|u_q\|_{2^*}^{-1} u_q) \geq \|u_q\|_{2^*}^2 \mu,$$

so that  $\mu_q \geq 0$ . This prove the above claim if  $\mu = 0$ , then  $\mu_q = 0$  for all  $q$ . Letting  $u = \|u_q\|_{2^*}^{-1} u_q$  for some  $q$ , we get that  $u$  is a positive smooth solution of the equation

$$-\Delta_g u + \alpha u = \mu u^{2^*-1}$$

such that  $\int_M u^{2^*} d\sigma_g = 1$ . □

### 11.3.3 The positive case: $\mu > 0$

In this case first we prove that the operator  $\Delta_g + h$  is coercive in the sense that, there exists  $\lambda > 0$  such that for any  $u \in H_1^2(M)$ ,

$$\int_M (|\nabla u|_g^2 + \alpha u^2) d\sigma_g \geq \lambda \|u\|_{H_1^2}^2.$$

From condition  $\mu > 0$  and from Hölder's inequalities follows the existence of  $\tilde{\mu} > 0$  such that for any  $u \in H_1^2(M)$ ,

$$\int_M (|\nabla u|_g^2 + \alpha u^2) d\sigma_g \geq \tilde{\mu} \int_M u^2 d\sigma_g.$$

We let  $\varepsilon \in (0, \frac{\tilde{\mu}}{2})$  be such that  $(1 - \varepsilon)\tilde{\mu} + \varepsilon h \geq \frac{\tilde{\mu}}{2}$ . Then

$$\begin{aligned} \int_M (|\nabla u|_g^2 + \alpha u^2) d\sigma_g &\geq \varepsilon \int_M (|\nabla u|_g^2 + \alpha u^2) d\sigma_g + (1 - \varepsilon) \int_M u^2 d\sigma_g \\ &\geq \int_M |\nabla u|_g^2 d\sigma_g + \frac{\tilde{\mu}}{2} \int_M u^2 d\sigma_g \\ &\geq \varepsilon \int_M (|\nabla u|_g^2 + u^2) d\sigma_g. \end{aligned}$$

In consequence, the operator  $\Delta_g + h$  is coercive.

The main result in the case  $\mu > 0$  is the following:

**Theorem 11.5** *Let  $(M, g)$  be a smooth compact Riemannian manifold of dimension  $d \geq 3$ , and  $\alpha : M \rightarrow \mathbb{R}$  be a smooth function. If*

$$\inf_{u \in \mathcal{H}} \int_M (|\nabla u|_g^2 + \alpha u^2) d\sigma_g < \frac{1}{K_d^2}.$$

*Then there exists  $u \in C^\infty(M), u > 0$  such that*

$$-\Delta_g u + \alpha u = \mu u^{2^* - 1}$$

*and  $\int_M u^{2^*} d\sigma_g = 1$ . In particular,  $u$  is a minimizing solution of the equation.*

*Proof* Let  $(u_k) \subset \mathcal{H}$  be a minimizing sequence for  $\mu$ . Using again Proposition 3.49 [19], we can replace  $u_k$  by  $|u_k|$ , so we can assume that the  $u_k$ 's are nonnegative. We have that  $(u_k)$  is bounded in  $H_1^2(M)$ . Taking into account that  $H_1^2(M)$  is a Hilbert space, we may thus assume that there exists  $u \in H_1^2(M)$ , such that

- 1)  $u_k \rightharpoonup u$  weakly in  $H_1^2(M)$ ,
- 2)  $u_k \rightarrow u$  strongly in  $L^2(M)$ ,
- 3)  $u_k \rightarrow u$  almost everywhere as  $k \rightarrow +\infty$ .

In particular,  $u$  is nonnegative. From the weak convergence  $u_k \rightharpoonup u$  follows that

$$\|u_k\|_2^2 = \|\nabla(u_k - u)\|_2^2 + \|u\|_2^2 + o(1),$$

for all  $k$ , where  $o(1) \rightarrow 0$  as  $k \rightarrow +\infty$ . From Brézis-Lieb Lemma (see Theorem B.2), it follows that

$$\|u_k\|_{2^*}^{2^*} = \|u_k - u\|_{2^*}^{2^*} + \|u\|_{2^*}^{2^*} + o(1),$$

for all  $k$ , where, as above,  $o(1) \rightarrow 0$  as  $k \rightarrow +\infty$ . By Sobolev embeddings, there exists  $B > 0$  such that for any  $k$ ,

$$\|u_k - u\|_{2^*}^2 \leq K_d^2 \|\nabla(u_k - u)\|_2^2 + B\|u_k - u\|_2^2.$$

Since  $u_k \in \mathcal{H}$ , it follows that

$$(1 - \|u\|_{2^*}^{2^*})^{2/2^*} \leq K_d^2 (\|\nabla u_k\|_2^2 - \|\nabla u\|_2^2) + o(1).$$

Since  $I(u_k) \rightarrow \mu$  and  $u_k \rightarrow u$  strongly in  $L^2(M)$ , we also have that

$$\begin{aligned} K_d^2 (\|\nabla u_k\|_2^2 - \|\nabla u\|_2^2) &= K_d^2 \mu - K_d^2 \left( \int_M |\nabla u|_g^2 d\sigma_g + \int_M \alpha u^2 d\sigma_g \right) + o(1) \\ &\leq K_d^2 \mu - K_d^2 \mu \|u\|_{2^*}^2 + o(1). \end{aligned}$$

Hence,

$$(1 - \|u\|_{2^*}^{2^*})^{2/2^*} \leq K_d^2 (1 - \|u\|_{2^*}^2).$$

From the assumption  $\mu K_d^2 < 1$  follows that

$$1 - \|u\|_{2^*}^2 \leq (1 - \|u\|_{2^*}^{2^*})^{2/2^*}.$$

From this we have  $\|u\|_{2^*} = 1$ . Then,  $\|u_k\|_2 \rightarrow \|u\|_2$  as  $k \rightarrow +\infty$ , and since

$$\|\nabla u_k\|_2^2 = \|\nabla(u_k - u)\|_2^2 + \|\nabla u\|_2^2 + o(1),$$

we get that  $u_k \rightarrow u$  strongly in  $H_1^2(M)$  as  $k \rightarrow +\infty$ . In particular,  $u$  is a minimizer for  $\mu$ , and  $u$  is a weak nonnegative solution of the equation

$$-\Delta_g u + \alpha u = \mu u^{2^*-1}.$$

From Theorem 11.3 and Proposition B.1 follows that  $u$  is smooth and positive.  $\square$

## 11.4 Comments and historical notes

In 1960, Yamabe [299] attempted to solve the problem: *Given a compact Riemannian manifold  $(M, g)$  of dimension  $d \geq 3$ , then there exists a conformal metric with constant scalar curvature.* Unfortunately, his proof contained an error, discovered by Trudinger [289] who repaired the proof, but only with a rather restrictive assumption on the manifold  $M$ . In 1976, Aubin [18] and Schoen [271] trait the remaining case. In 1984, Cherrier [69] extend the Yamabe problem to the compact Riemann

manifold with boundary. In 1987 appear a very nice paper due to Lee and Parker [188] about the Yamabe problem, where here gave a proof of the Yamabe problem unifying Aubin's and Schoen's arguments. The Yamabe problem on compact Riemannian manifold with boundary was studied in many papers by Escobar [106], [107], [108]. In 1993, Hebey and Vaugon [140] studied the equivariant Yamabe problem. In the last years many papers are dedicated to study different aspects of Yamabe or Yamabe like problems. For this see the books of Ambrosetti and Malchiodi [7], Druet, Hebey, and Robert [98], and Hebey [137], [138] and also the papers of Li and Li [191] and Il'yasov and Runst [144] and there references. Also many papers is dedicated to study the Yamabe invariant, see for example a very nice paper of Ammann, Dahl, and Humbert [11]. These problems suggest to study elliptic or semilinear elliptic problems on a compact Riemannian manifold with or without boundary. In the present exists a very reach literature which are dedicated to study these problem.

The results presented in the present chapter are contained in the paper of Hebey [139].

# 12

## Problems to Part II

In mathematics the art of proposing a question must be held of higher value than solving it.

---

Georg Cantor (1845–1918)

**Problem 12.1** (Hopf's lemma) Let  $(M, g)$  be a connected, compact Riemannian manifold and  $f : M \rightarrow \mathbb{R}$  a smooth function such that

$$\Delta_g f \geq 0.$$

Then  $f$  is constant.

**Problem 12.2** Consider  $\mathbb{S}^n$  the  $n$ -dimensional unit sphere endowed with the Riemannian metric induced by the inclusion  $\iota : \mathbb{S}^n \hookrightarrow \mathbb{R}^{n+1}$ . Show that for any smooth function  $f : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$  we have

$$(\Delta_{\mathbb{R}^{n+1}} f)|_{\mathbb{S}^n} = \Delta_{\mathbb{S}^n}(f|_{\mathbb{S}^n}) - \frac{\partial^2 f}{\partial r^2}|_{\mathbb{S}^n} - n \frac{\partial f}{\partial r}|_{\mathbb{S}^n},$$

where  $\Delta_{\mathbb{R}^{n+1}}$ ,  $\Delta_{\mathbb{S}^n}$  and  $\frac{\partial}{\partial r}$  are the Laplace operator on  $\mathbb{R}^{n+1}$ ,  $\mathbb{S}^n$ , and the radial derivative, respectively.

**Problem 12.3** Let  $\mathbb{S}^n$  be the unit sphere endowed with the usual Riemannian structure from  $\mathbb{R}^{n+1}$ . Denote by  $\mathcal{H}_k$  the vector space of the harmonic polynomial of degree  $k \geq 0$  defined on  $\mathbb{R}^{n+1}$ . Let  $\tilde{\mathcal{H}}_k = \{f|_{\mathbb{S}^n} : f \in \mathcal{H}_k\}$ .

- Show that  $\Delta_{\mathbb{S}^n} f = k(n+k-1)f$ , for all  $f \in \tilde{\mathcal{H}}_k$  and hence  $k(n+k-1)$  is an eigenvalue of the Laplace operator  $\Delta_{\mathbb{S}^n}$ .
- $\tilde{\mathcal{H}}_k$  is the eigenspace corresponding to the eigenvalue  $\lambda_k = k(n+k-1)$ .
- The set  $\{k(n+k-1) : k \in \mathbb{N}\}$  is the set of eigenvalues of  $\Delta_{\mathbb{S}^n}$ .

**Problem 12.4** (M. Obata [230] ) If  $g$  is a metric on  $\mathbb{S}^n$  that is conformal to the standard metric  $\bar{g}$  and has constant scalar curvature, then up to a constant scale factor,  $g$  is obtained from  $\bar{g}$  by a conformal diffeomorphism of the sphere.

**Problem 12.5** Let  $g_0$  be the standard Riemannian metric on  $\mathbb{R}^n$ . Let  $h = (h_{ij})$  be a symmetric bilinear form with compact support and consider the metric  $g_\varepsilon = g_0 + \varepsilon h$  on  $\mathbb{R}^n$ . If  $R_{g_\varepsilon}$  denotes the scalar curvature of  $g_\varepsilon$  prove that

$$R_{g_\varepsilon}(x) = \varepsilon R_1(x) + \varepsilon^2 R_2(x) + o(\varepsilon^2),$$

where

$$R_1(x) = \sum_{i,j} \frac{\partial^2 h_{ij}}{\partial x_i \partial x_j} - \Delta \operatorname{tr} h,$$

and

$$\begin{aligned} R_2 = & -2 \sum_{k,j,l} h_{kj} \frac{\partial^2 h_{ij}}{\partial^2 x_l} + \sum_{k,j,l} h_{kj} \frac{\partial^2 h_{ll}}{\partial x_j \partial x_k} + \frac{3}{4} \sum_{k,j,l} \frac{\partial h_{jl}}{\partial x_k} \frac{\partial h_{jl}}{\partial x_k} \\ & - \sum_{k,j,l} \frac{\partial h_{jl}}{\partial x_l} \frac{\partial h_{jk}}{\partial x_k} + \sum_{k,j,l} \frac{\partial h_{jl}}{\partial x_l} \frac{\partial h_{kk}}{\partial x_j} - \frac{1}{4} \sum_{k,j,l} \frac{\partial h_{ll}}{\partial x_j} \frac{\partial h_{kk}}{\partial x_j} - \frac{1}{2} \sum_{k,j,l} \frac{\partial h_{kl}}{\partial x_j} \frac{\partial h_{jk}}{\partial x_l}. \end{aligned}$$

Also the following formula holds.

$$|g_\varepsilon|^{\frac{1}{2}} = 1 + \frac{\varepsilon}{2} \operatorname{tr} h + \varepsilon^2 \left( \frac{1}{8} (\operatorname{tr} h)^2 - \frac{1}{4} \operatorname{tr}(h^2) \right) + o(\varepsilon^2).$$

**Problem 12.6** Let  $(M, \overset{\circ}{g})$  be a compact Riemannian manifold with boundary  $n \geq 3$ . Let  $0 \in \partial M$  be a point. We assume that  $1 \leq i, j, k, l \leq n-1$ . Let  $(x_1, x_2, \dots, x_{n-1})$  be normal coordinate on  $\partial M$  at the point 0. Let  $\gamma(t)$  be the geodesics leaving from  $(x_1, x_2, \dots, x_{n-1})$  in the orthogonal direction to  $\partial M$  and parameterized by the arc length. In this case  $(x_1, x_2, \dots, x_{n-1})$  will be called Fermi coordinates at  $0 \in \partial M$ . In these coordinates the arc length is written as

$$ds^2 = dt^2 + g_{ij}(x, t) dx_i dx_j.$$

If  $g = \det(\overset{\circ}{g})$  then  $g = \det(g_{ij})$ . If  $x_i$  and  $t$  are sufficiently small, then the following formula holds

$$\sqrt{g} = 1 - Ht + \frac{1}{2} (H^2 - \|\pi\|^2 - \operatorname{Ric}(\eta))t^2 - H_i t x_i - \frac{1}{6} \bar{R}_{ij} x_i x_j + O(|(x, t)|^3),$$

where  $\pi$  is the second fundamental form,  $H$  is its trace and  $\bar{R}_{ij}$  are the coefficients of the Ricci tensor of  $\partial M$ .

**Problem 12.7** Let  $(M, g)$  be a connected compact  $n$ -dimensional Riemannian manifold without boundary. An eigenvalue of the  $p$ -Laplace operator  $\Delta_p u := \operatorname{div}_g(|\nabla u|^{p-2} \nabla u)$  is a real number  $\lambda$  such that there exists  $u \in W_1^p(M) \setminus \{0\}$  such that

$$-\Delta_p u = \lambda |u|^{p-2} u.$$

We denote by  $\sigma_p(M, g)$  the set of all nonzero eigenvalues. The set  $\sigma_p(M, g)$  is not empty, unbounded and included in  $(0, +\infty)$ . If  $\lambda_1 = \inf \sigma_p(M, g)$  prove that  $\lambda_1 \in \sigma_p(M, g)$  and

$$\lambda_1 = \min \left\{ \int_M |\nabla \varphi|^p dv_g : \varphi \in \Sigma_0 \right\}$$

where

$$\Sigma_0 = \left\{ \varphi \in W_1^p(M) : \int_M |\varphi|^p dv_g = 1, \int_M |\varphi|^{p-2} \varphi dv_g = 0 \right\}.$$

If  $\omega \subset M$  is a non-empty open subset, define

$$\mu(\omega) = \min \left\{ \int_\omega |\nabla \varphi|^p : \varphi \in W_0^{1,p}(\omega), \int_\omega |\varphi|^p dv_g = 1 \right\}.$$

Prove that

$$\lambda_1 = \min_{(\omega, \tilde{\omega})} \max\{\mu(\omega), \mu(\tilde{\omega})\},$$

where  $(\omega, \tilde{\omega})$  runs over the set of couples of non-empty disjoint open subset of  $M$  and if  $u$  is any nonzero eigenfunction associated to  $\lambda_1$ , then  $M \setminus u^{-1}(0)$  has two connected open components.

**Problem 12.8** Let  $(M, g)$  be an  $n$ -dimensional compact Riemannian manifold without boundary,  $K : M \rightarrow \mathbb{R}$  a continuous function ( $K \neq 0$ ) and  $\lambda \in \mathbb{R}$ . We consider the following equation

$$\Delta_g u + \lambda u + K(x)u^q = 0.$$

- a) If  $\lambda = 0$  and  $1 < q < \frac{n+2}{n-2}$  then the equation admits a positive solutions if and only if the following conditions hold:
- i).  $\int_M K dv_g < 0$ ;
  - ii). there exists  $x_0 \in M$  such that  $K(x_0) > 0$ .
- b) If  $\lambda < 0$  and  $1 < q < \frac{n+2}{n-2}$  then the equation admits positive solutions if and only if the condition ii) holds.

**Problem 12.9** Let  $\Omega \subset \mathbb{R}^N$  be a bounded domain with smooth boundary and let  $\lambda_1$  denote the first eigenvalue of  $(-\Delta)$  in  $H_0^1(\Omega)$ . Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be a differentiable function such that  $f(0) = 0$ ,  $f'(0) < \lambda_1$ , and

$$\lambda_1 < \lim_{u \rightarrow +\infty} \frac{f(u)}{u} < \infty.$$

Prove that the problem

$$\begin{cases} -\Delta u = f(u) & \text{in } \Omega \\ u > 0 & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega \end{cases}$$

has a solution.

**Problem 12.10** Let  $\Omega \subset \mathbb{R}^N$  be a bounded domain with smooth boundary and assume that  $p$  is a real number such that  $1 < p < (N + 2)/(N - 2)$  if  $N \geq 3$  and  $1 < p < \infty$  if  $N \in \{1, 2\}$ .

Prove that there exists  $\lambda^* > 0$  such that the problem

$$\begin{cases} -\Delta u = \lambda(1 + u)^p & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega \end{cases}$$

has at least two solutions for any  $\lambda \in (0, \lambda^*)$ .

**Problem 12.11** Let  $\Omega \subset \mathbb{R}^N$  be a bounded domain with smooth boundary and assume that  $p$  is a real number such that  $1 < p < (N + 2)/(N - 2)$  if  $N \geq 3$  and  $1 < p < \infty$  if  $N \in \{1, 2\}$ . Consider the problem

$$\begin{cases} -\Delta u = u^p + f(x) & \text{in } \Omega \\ u > 0 & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (12.1)$$

where  $f$  is a smooth function.

Prove that there exists  $\delta > 0$  such that problem (12.1) has a solution, provided that  $\|f\|_{L^\infty} \leq \delta$ .

**Problem 12.12** Let  $\Omega \subset \mathbb{R}^N$  be a bounded domain with smooth boundary and assume that  $p$  is a real number such that  $1 < p < (N + 2)/(N - 2)$  if  $N \geq 3$  and  $1 < p < \infty$  if  $N \in \{1, 2\}$ . Consider the problem

$$\begin{cases} -\Delta u = |u|^p + f(x) & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (12.2)$$

where  $f$  is a smooth function.

Denote

$$K = \{f \in C^{0,\alpha}(\bar{\Omega}); \text{ problem (12.2) has a solution}\}$$

and

$$P = \{h \in C^{0,\alpha}(\bar{\Omega}); h \geq 0 \text{ in } \Omega\}.$$

- a) Prove that  $K$  is a convex set and  $K \setminus P \subset K$ .
- b) Prove that for every  $f \in \text{Int } K$ , problem (12.2) has at least two solutions. In particular, if  $f \leq 0$  in  $\Omega$ , then problem (12.2) has at least two solutions.

# 13

## Mathematical Preliminaries

Knowledge is power. (Ipsa  
Scientia Potestas Est)

---

Sir Francis Bacon  
(1561–1626), *Meditationes  
Sacrae. De Hæresibus*

### 13.1 Metrics, geodesics, flag curvature

Let  $M$  be a connected  $m$ -dimensional  $C^\infty$  manifold and let  $TM = \bigcup_{p \in M} T_p M$  be its tangent bundle. If the continuous function  $F : TM \rightarrow [0, \infty)$  satisfies the conditions that it is  $C^\infty$  on  $TM \setminus \{0\}$ ;  $F(tu) = tF(u)$  for all  $t \geq 0$  and  $u \in TM$ , i.e.,  $F$  is positively homogeneous of degree one; and the matrix  $g_{ij}(u) := [\frac{1}{2}F^2]_{y^i y^j}(u)$  is positive definite for all  $u \in TM \setminus \{0\}$ , then we say that  $(M, F)$  is a *Finsler manifold*. If  $F$  is absolutely homogeneous, then  $(M, F)$  is said to be *reversible*. Let  $\pi^*TM$  be the pull-back of the tangent bundle  $TM$  by  $\pi : TM \setminus \{0\} \rightarrow M$ ; then

$$g_{(p,y)} := g_{ij(p,y)} dp^i \otimes dp^j := [\frac{1}{2}F^2]_{y^i y^j} dp^i \otimes dp^j, \quad p \in M, \quad y \in T_p M, \quad (13.1)$$

is the natural Riemannian metric on the pulled-back bundle  $\pi^*TM$ .

Unlike the Levi-Civita connection in Riemann geometry, there is no unique natural connection in the Finsler case. Among these connections on  $\pi^*TM$ , we choose the *Chern connection* whose coefficients are denoted by  $\Gamma_{jk}^i$ ; see Bao-Chern-Shen [23, p.38]. This connection induces the *curvature tensor*, denoted by  $R$ ; see Bao-Chern-Shen [23, Chapter 3]. The Chern connection defines the *covariant derivative*  $D_V U$  of a vector field  $U$  in the direction  $V \in T_p M$ . Since, in general, the Chern connection coefficients  $\Gamma_{jk}^i$  in natural coordinates have a directional dependence, we must say explicitly that  $D_V U$  is defined with a fixed reference vector.

In particular, let  $\sigma : [0, r] \rightarrow M$  be a smooth curve with velocity field  $T = \dot{\sigma}(t)$ . Suppose that  $U$  and  $W$  are vector fields defined along  $\sigma$ . We define  $D_T U$  with reference vector  $W$  as

$$D_T U = \left[ \frac{dU^i}{dt} + U^j T^k (\Gamma_{jk}^i)_{(\sigma, W)} \right] \frac{\partial}{\partial p^i} \Big|_{\sigma(t)},$$

where  $\left\{ \frac{\partial}{\partial p^i} \Big|_{\sigma(t)} \right\}_{i=1, m}$  is a basis of  $T_{\sigma(t)} M$ . A  $C^\infty$  curve  $\sigma : [0, r] \rightarrow M$ , with velocity  $T = \dot{\sigma}$  is a (Finslerian) *geodesic* if

$$D_T \left[ \frac{T}{F(T)} \right] = 0 \quad \text{with reference vector } T. \quad (13.2)$$

If the Finslerian velocity of the geodesic  $\sigma$  is constant, then (13.2) becomes

$$\frac{d^2 \sigma^i}{dt^2} + \frac{d\sigma^j}{dt} \frac{d\sigma^k}{dt} (\Gamma_{jk}^i)_{(\sigma, T)} = 0, \quad i = 1, \dots, m = \dim M. \quad (13.3)$$

For any  $p \in M$  and  $y \in T_p M$  we may define the *exponential map*  $\exp_p : T_p M \rightarrow M$ ,  $\exp_p(y) = \sigma(1, p, y)$ , where  $\sigma(t, p, y)$  is the unique solution (geodesic) of the second order differential equation (13.2) (or, (13.3)) which passes through  $p$  at  $t = 0$  with velocity  $y$ .

If  $U$ ,  $V$  and  $W$  are vector fields along a curve  $\sigma$ , which has velocity  $T = \dot{\sigma}$ , we have the *derivative rule*

$$\frac{d}{dt} g_{(\sigma, W)}(U, V) = g_{(\sigma, W)}(D_T U, V) + g_{(\sigma, W)}(U, D_T V) \quad (13.4)$$

whenever  $D_T U$  and  $D_T V$  are with reference vector  $W$  and *one* of the following conditions holds:

- $U$  or  $V$  is proportional to  $W$ , or
- $W = T$  and  $\sigma$  is a geodesic.

Let  $\gamma : [0, 1] \rightarrow M$  be a smooth regular curve and  $\Sigma : [0, 1] \times [-\varepsilon, \varepsilon] \rightarrow M$  be a smooth regular variation of  $\gamma$  (i.e.  $\Sigma(t, 0) = \gamma(t)$  for all  $t \in [0, 1]$ ) with variation vector field  $U = U(t, u) = \frac{\partial \Sigma}{\partial u}$ . Then

$$\frac{\partial}{\partial u} g_{(\sigma, T)}(T, T) = 2g_{(\sigma, T)}(T, D_U T), \quad (13.5)$$

where  $T = \frac{\partial \Sigma}{\partial t}$  and the covariant derivative  $D_U T$  is with reference vector  $T$ .

A vector field  $J$  along a geodesic  $\sigma : [0, r] \rightarrow M$  (with velocity field  $T$ ) is said to be a *Jacobi field* if it satisfies the equation

$$D_T D_T J + R(J, T)T = 0, \quad (13.6)$$

where  $R$  is the curvature tensor. Here, the covariant derivative  $D_T$  is defined with reference vector  $T$ .

We say that  $q$  is conjugate to  $p$  along the geodesic  $\sigma$  if there exists a nonzero Jacobi field  $J$  along  $\sigma$  which vanishes at  $p$  and  $q$ .

Let  $\gamma : [0, r] \rightarrow M$  be a piecewise  $C^\infty$  curve. Its integral length is defined as

$$L_F(\gamma) = \int_0^r F(\gamma(t), \dot{\gamma}(t)) dt.$$

Let  $\Sigma : [0, r] \times [-\varepsilon, \varepsilon] \rightarrow M$  ( $\varepsilon > 0$ ) be a piecewise  $C^\infty$  variation of a geodesic  $\gamma : [0, r] \rightarrow M$  with  $\Sigma(\cdot, 0) = \gamma$ . Let  $T = T(t, u) = \frac{\partial \Sigma}{\partial t}$ ,  $U = U(t, u) = \frac{\partial \Sigma}{\partial u}$  the velocities of the  $t$ -curves and  $u$ -curves, respectively. The formula for the first variation of arc length, see Bao-Chern-Shen [23, Exercise 5.1.4], gives

$$L'_F(\Sigma(\cdot, 0)) := \frac{d}{du} L_F(\Sigma(\cdot, u))|_{u=0} = \left[ g_T \left( U, \frac{T}{F(T)} \right) \Big|_{u=0} \right] \Big|_{t=0}^{t=r}. \quad (13.7)$$

For  $p, q \in M$ , denote by  $\Gamma(p, q)$  the set of all piecewise  $C^\infty$  curves  $\gamma : [0, r] \rightarrow M$  such that  $\gamma(0) = p$  and  $\gamma(r) = q$ . Define the map  $d_F : M \times M \rightarrow [0, \infty)$  by

$$d_F(p, q) = \inf_{\gamma \in \Gamma(p, q)} L_F(\gamma). \quad (13.8)$$

Of course, we have  $d_F(p, q) \geq 0$ , where equality holds if and only if  $p = q$ , and the triangle inequality holds, i.e.,

$$d_F(p_0, p_2) \leq d_F(p_0, p_1) + d_F(p_1, p_2)$$

for every  $p_0, p_1, p_2 \in M$ . In general, since  $F$  is only a positive homogeneous function,  $d_F(p, q) \neq d_F(q, p)$ ; thus,  $(M, d_F)$  is only a quasi-metric space. If  $(M, g)$  is a Riemannian manifold, we use the notation  $d_g$  instead of  $d_F$  which becomes a usual metric function.

A geodesic  $\sigma : [0, 1] \rightarrow M$  from  $p = \sigma(0)$  to  $q = \sigma(1)$  is said to be minimal if  $L_F(\sigma) = d_F(p, q)$ .

For  $p \in M$ ,  $r > 0$ , we define the forward and backward Finsler-metric balls, respectively, with center  $p \in M$  and radius  $r > 0$ , by

$$\mathcal{B}_p^+(r) = \{q \in M : d_F(p, q) < r\} \quad \text{and} \quad \mathcal{B}_p^-(r) = \{q \in M : d_F(q, p) < r\}.$$

We denote by  $B_p(r) := \{y \in T_p M : F(p, y) < r\}$  the open tangent ball at  $p \in M$  with radius  $r > 0$ . It is well-known that the topology generated by the forward (resp. backward) metric balls coincides with the underlying manifold topology, respectively.

By Whitehead's theorem (see [297] or [23, Exercise 6.4.3, p. 164]) and [23, Lemma 6.2.1, p. 146] we can conclude the following useful local result (see also [182]).

**Proposition 13.1** *Let  $(M, F)$  be a Finsler manifold, where  $F$  is positively (but perhaps not absolutely) homogeneous of degree one. For every point  $p \in M$  there exist a small  $\rho_p > 0$  and  $c_p > 1$  (depending only on  $p$ ) such that for every pair of points  $q_0, q_1$  in  $\mathcal{B}_p^+(\rho_p)$  we have*

$$\frac{1}{c_p} d_F(q_1, q_0) \leq d_F(q_0, q_1) \leq c_p d_F(q_1, q_0). \quad (13.9)$$

Moreover, for every real number  $k \geq 1$  and  $q \in \mathcal{B}_p^+(\rho_p/k)$  the mapping  $\exp_q$  is  $C^1$ -diffeomorphism from  $B_q(2\rho_p/k)$  onto  $\mathcal{B}_q^+(2\rho_p/k)$  and every pair of points  $q_0, q_1$  in  $\mathcal{B}_p^+(\rho_p/k)$  can be joined by a unique minimal geodesic from  $q_0$  to  $q_1$  lying entirely in  $\mathcal{B}_p^+(\rho_p/k)$ .

A set  $M_0 \subseteq M$  is *forward bounded* if there exist  $p \in M$  and  $r > 0$  such that  $M_0 \subseteq \mathcal{B}_p^+(r)$ . Similarly,  $M_0 \subseteq M$  is *backward bounded* if there exist  $p \in M$  and  $r > 0$  such that  $M_0 \subseteq \mathcal{B}_p^-(r)$ .

A set  $M_0 \subseteq M$  is *geodesic convex* if for any two points of  $M_0$  there exists a unique geodesic joining them which belongs entirely to  $M_0$ .

Let  $(p, y) \in TM \setminus 0$  and let  $V$  be a section of the pulled-back bundle  $\pi^*TM$ . Then,

$$K(y, V) = \frac{g_{(p,y)}(R(V, y)y, V)}{g_{(p,y)}(y, y)g_{(p,y)}(V, V) - [g_{(p,y)}(y, V)]^2} \quad (13.10)$$

is the *flag curvature* with flag  $y$  and transverse edge  $V$ . In particular, when the Finsler structure  $F$  arises from a Riemannian metric  $g$  (i.e., the fundamental tensor  $g_{ij} = [\frac{1}{2}F^2]_{y_i y_j}$  does not depend on the direction  $y$ ), the flag curvature coincides with the usual *sectional curvature*.

If  $K(V, W) \leq 0$  for every  $0 \neq V, W \in T_p M$ , and  $p \in M$ , with  $V$  and  $W$  not collinear, we say that the flag curvature of  $(M, F)$  is *non-positive*.

A Finsler manifold  $(M, F)$  is said to be *forward (resp. backward) geodesically complete* if every geodesic  $\sigma : [0, 1] \rightarrow M$  parameterized to have constant Finslerian speed, can be extended to a geodesic defined on  $[0, \infty)$  (resp.  $(-\infty, 1]$ ).  $(M, F)$  is *geodesically complete* if every geodesic  $\sigma : [0, 1] \rightarrow M$  can be extended to a geodesic defined on  $(-\infty, \infty)$ . A geodesically complete Riemannian manifold is called *complete Riemannian manifold*.

**Theorem 13.1** [Theorem of Hopf-Rinow, [23, p.168]] *Let  $(M, F)$  be*

a connected Finsler manifold, where  $F$  is positively (but perhaps not absolutely) homogeneous of degree one. The following two criteria are equivalent:

- (a).  $(M, F)$  is forward (backward) geodesically complete;
- (b). Every closed and forward (backward) bounded subset of  $(M, d_F)$  is compact.

Moreover, if any of the above holds, then every pair of points in  $M$  can be joined by a minimizing geodesic.

**Theorem 13.2** [Theorem of Cartan-Hadamard, [23, p.238]] *Let  $(M, F)$  be a forward/backward geodesically complete, simply connected Finsler manifold of non-positive flag curvature. Then:*

- (a). Geodesics in  $(M, F)$  do not contain conjugate points.
- (b). The exponential map  $\exp_p : T_pM \rightarrow M$  is a  $C^1$  diffeomorphism from the tangent space  $T_pM$  onto the manifold  $M$ .

A Finsler manifold  $(M, F)$  is a *Minkowski space* if  $M$  is a vector space and  $F$  is a Minkowski norm inducing a Finsler structure on  $M$  by translation; its flag curvature is identically zero, the geodesics are straight lines, and for any two points  $p, q \in M$ , we have  $F(q - p) = d_F(p, q)$ , see Bao-Chern-Shen [23, Chapter 14]. In particular,  $(M, F)$  is both forward and backward geodesically complete. The *fundamental inequality* for Minkowski norms implies

$$|g_y(y, w)| \leq \sqrt{g_y(y, y)} \cdot \sqrt{g_w(w, w)} = F(y) \cdot F(w) \text{ for all } y \neq 0 \neq w, \quad (13.11)$$

which is the *generalized Cauchy-Schwarz inequality*, see Bao-Chern-Shen [23, p. 6-10].

A Finsler manifold is of *Berwald type* if the Chern connection coefficients  $\Gamma_{ij}^k$  in natural coordinates depend only on the base point. Special Berwald spaces are the (locally) *Minkowski spaces* and the *Riemannian manifolds*. In the latter case, the Chern connection coefficients  $\Gamma_{ij}^k$  coincide with the usual Christoffel symbols

$$\bar{\Gamma}_{ij}^k(p) = \frac{1}{2} \left[ \left( \frac{\partial g_{mj}}{\partial p_i} \right)_p + \left( \frac{\partial g_{mi}}{\partial p_j} \right)_p - \left( \frac{\partial g_{ij}}{\partial p_m} \right)_p \right] g^{mk}(p)$$

where the  $g^{ij}$ 's are such that  $g_{im}g^{mj} = \delta_{ij}$ .

A Riemannian manifold is said to be of *Hadamard-type*, if it is simply connected, complete, and has non-positive sectional curvature.

**Theorem 13.3** [Cosine inequality, see [93, Lemma 3.1]] *Let  $(M, g)$  be a Hadamard-type Riemannian manifold. Consider the geodesic triangle*

determined by vertices  $a, b, c \in M$ . If  $\hat{c}$  is the angle belonging to vertex  $c$  and if  $A = d_g(b, c)$ ,  $B = d_g(a, c)$ ,  $C = d_g(a, b)$ , then

$$A^2 + B^2 - 2AB \cos \hat{c} \leq C^2.$$

The following result is probably known, but since we have not found an explicit reference, we give its proof.

**Proposition 13.2** *Let  $(M, g)$  be a complete, finite-dimensional Riemannian manifold. Then any geodesic convex set  $K \subset M$  is contractible.*

*Proof* Let us fix  $p \in K$  arbitrarily. Since  $K$  is geodesic convex, every point  $q \in K$  can be connected to  $p$  uniquely by the geodesic segment  $\gamma_q : [0, 1] \rightarrow K$ , i.e.,  $\gamma_q(0) = p$ ,  $\gamma_q(1) = q$ . Moreover, the map  $K \ni q \mapsto \exp_p^{-1}(q) \in T_p M$  is well-defined and continuous. Note actually that  $\gamma_q(t) = \exp_p(t \exp_p^{-1}(q))$ . We define the map  $F : [0, 1] \times K \rightarrow K$  by  $F(t, q) = \gamma_q(t)$ . It is clear that  $F$  is continuous,  $F(1, q) = q$  and  $F(0, q) = p$  for all  $q \in K$ , i.e., the identity map  $\text{id}_K$  is homotopic to the constant map  $p$ .  $\square$

### 13.2 Busemann-type inequalities on Finsler manifolds

In the forties, Busemann developed a synthetic geometry on metric spaces. In particular, he axiomatically elaborated a whole theory of non-positively curved metric spaces which have no differential structure a priori and they possess the essential qualitative geometric properties of Finsler manifolds. These spaces are the so-called *G-spaces*, see Busemann [55, p. 37]. This notion of non-positive curvature requires that in small geodesic triangles the length of a side is at least the twice of the geodesic distance of the mid-points of the other two sides, see Busemann [55, p. 237].

To formulate in a precise way this notion, let  $(M, d)$  be a quasi-metric space and for every  $p \in M$  and radius  $r > 0$ , we introduce the *forward* and *backward metric balls*

$$B_p^+(r) = \{q \in M : d(p, q) < r\} \quad \text{and} \quad B_p^-(r) = \{q \in M : d(q, p) < r\}.$$

A continuous curve  $\gamma : [a, b] \rightarrow M$  with  $\gamma(a) = x$ ,  $\gamma(b) = y$  is a *shortest geodesic*, if  $l(\gamma) = d(x, y)$ , where  $l(\gamma)$  denotes the *generalized length* of  $\gamma$

and it is defined by

$$l(\gamma) = \sup\left\{\sum_{i=1}^n d(\gamma(t_{i-1}), \gamma(t_i)) : a = t_0 < t_1 < \dots < t_n = b, n \in \mathbb{N}\right\}.$$

In the sequel, we always assume that the shortest geodesics are parametrized proportionally to arclength, i.e.,  $l(\gamma|_{[0,t]}) = tl(\gamma)$ .

**Remark 13.1** A famous result of Busemann-Meyer (see [56, Theorem 2, p. 186]) from Calculus of Variations shows that the generalized length  $l(\gamma)$  and the integral length  $L_F(\gamma)$  of any (piecewise)  $C^\infty$  curves coincide for Finsler manifolds. Therefore, the minimal Finsler geodesic and shortest geodesic notions coincide.

We say that  $(M, d)$  is a *locally geodesic (length) space* if for every point  $p \in M$  there is a  $\rho_p > 0$  such that for every two points  $x, y \in B_p^+(\rho_p)$  there exists a shortest geodesic joining them.

**Definition 13.1** A *locally geodesic space*  $(M, d)$  is said to be a Busemann non-positive curvature space (*shortly*, Busemann NPC space), if for every  $p \in M$  there exists  $\rho_p > 0$  such that for any two shortest geodesics  $\gamma_1, \gamma_2 : [0, 1] \rightarrow M$  with  $\gamma_1(0) = \gamma_2(0) = x \in B_p^+(\rho_p)$  and with endpoints  $\gamma_1(1), \gamma_2(1) \in B_p^+(\rho_p)$  we have

$$2d(\gamma_1(\frac{1}{2}), \gamma_2(\frac{1}{2})) \leq d(\gamma_1(1), \gamma_2(1)).$$

(We shall say that  $\gamma_1$  and  $\gamma_2$  satisfy the Busemann NPC inequality).

Let  $(M, g)$  be a Riemannian manifold and  $(M, d_g)$  the metric space induced by itself. In this context, the Busemann NPC inequality is well-known. Namely, we have

**Proposition 13.3** [55, Theorem (41.6)]  $(M, d_g)$  is a Busemann NPC space if and only if the sectional curvature of  $(M, g)$  is non-positive.

However, the picture for Finsler spaces is quite different. To see this, we consider the Hilbert metric of the interior of a simple, closed and convex curve  $C$  in the Euclidean plane. In order to describe this metric, let  $M_C \subset \mathbb{R}^2$  be the region defined by the interior of the curve  $C$  and fix  $x_1, x_2 \in \text{Int}(M_C)$ . Assume first that  $x_1 \neq x_2$ . Since  $C$  is a convex curve, the straight line passing to the points  $x_1, x_2$  intersects the curve  $C$  in two points; denote them by  $u_1, u_2 \in C$ . Then, there are  $\tau_1, \tau_2 \in (0, 1)$  such that  $x_i = \tau_i u_1 + (1 - \tau_i) u_2$  ( $i = 1, 2$ ).

The Hilbert distance between  $x_1$  and  $x_2$  is

$$d_H(x_1, x_2) = \left| \log \left( \frac{1 - \tau_1}{1 - \tau_2} \cdot \frac{\tau_2}{\tau_1} \right) \right|.$$

We complete this definition by  $d_H(x, x) = 0$  for every  $x \in \text{Int}(M_C)$ . One can easily prove that  $(\text{Int}(M_C), d_H)$  is a metric space and it is a projective Finsler metric with constant flag curvature  $-1$ . However, one has

**Proposition 13.4** [Kelly-Straus [158]] *The metric space  $(\text{Int}(M_C), d_H)$  is a Busemann NPC space if and only if the curve  $C \subset \mathbb{R}^2$  is an ellipse.*

This means that although for Riemannian spaces the non-positivity of the sectional curvature and Busemann's curvature conditions are mutually equivalent, the non-positivity of the flag curvature of a *generic* Finsler manifold is not enough to guarantee Busemann's property.

Therefore, in order to obtain a characterization of Busemann's curvature condition for Finsler spaces, we have two possibilities:

- (I). To find a *new* notion of curvature in Finsler geometry such that for an arbitrary Finsler manifold the non-positivity of this curvature is equivalent with the Busemann non-positive curvature condition, as it was proposed by Shen [280, Open Problem 41]; or,
- (II). To keep the flag curvature, but put some restrictive condition on the Finsler metric.

In spite of the fact that (reversible) Finsler manifolds are included in  $G$ -spaces, only few results are known which establish a link between the *differential invariants* of a Finsler manifold and the *metric properties* of the induced metric space. The main result of this section is due Kristály-Kozma [173] (see also Kristály-Varga-Kozma [182]), which makes a strong connection between an analytical property and a synthetic concept of non-positively curved metric spaces. Namely, we have

**Theorem 13.4** *Let  $(M, F)$  be a Berwald space with non-positive flag curvature, where  $F$  is positively (but perhaps not absolutely) homogeneous of degree one. Then  $(M, d_F)$  is a Busemann NPC space.*

*Proof* Let us fix  $p \in M$  and consider  $\rho_p > 0$ ,  $c_p > 1$  from Proposition 13.1. We will prove that  $\rho'_p = \frac{\rho_p}{c_p}$  is an appropriate choice in Definition 13.1. To do this, let  $\gamma_1, \gamma_2 : [0, 1] \rightarrow M$  be two (minimal) geodesics with  $\gamma_1(0) = \gamma_2(0) = x \in \mathcal{B}_p^+(\rho'_p)$  and  $\gamma_1(1), \gamma_2(1) \in \mathcal{B}_p^+(\rho'_p)$ . By Proposition 13.1, we can construct a unique geodesic  $\gamma : [0, 1] \rightarrow M$  joining  $\gamma_1(1)$

with  $\gamma_2(1)$  and  $d_F(\gamma_1(1), \gamma_2(1)) = L_F(\gamma)$ . Clearly,  $\gamma(s) \in \mathcal{B}_p^+(\rho'_p)$  for all  $s \in [0, 1]$  (we applied Proposition 13.1 for  $k = c_p$ ). Moreover,  $x \in \mathcal{B}_{\gamma(s)}^+(2\rho_p)$ . Indeed, by (13.9), we obtain

$$\begin{aligned} d_F(\gamma(s), x) &\leq d_F(\gamma(s), p) + d_F(p, x) \\ &\leq c_p d_F(p, \gamma(s)) + \rho'_p \leq (c_p + 1)\rho'_p \\ &< 2\rho_p. \end{aligned}$$

Therefore, we can define  $\Sigma : [0, 1] \times [0, 1] \rightarrow M$  by

$$\Sigma(t, s) = \exp_{\gamma(s)}((1 - t) \cdot \exp_{\gamma(s)}^{-1}(x)).$$

The curve  $t \mapsto \Sigma(1 - t, s)$  is a radial geodesic which joins  $\gamma(s)$  with  $x$ . Taking into account that  $(M, F)$  is of Berwald type, the reverse of  $t \mapsto \Sigma(1 - t, s)$ , i.e.,  $t \mapsto \Sigma(t, s)$  is also a geodesic for all  $s \in [0, 1]$ , see Bao-Chern-Shen [23, Exercise 5.3.3, p. 128]. Moreover,  $\Sigma(0, 0) = x = \gamma_1(0)$ ,  $\Sigma(1, 0) = \gamma(0) = \gamma_1(1)$ . From the uniqueness of the geodesic between  $x$  and  $\gamma_1(1)$ , we have  $\Sigma(\cdot, 0) = \gamma_1$ . Analogously, we have  $\Sigma(\cdot, 1) = \gamma_2$ . Since  $\Sigma$  is a geodesic variation (of the curves  $\gamma_1$  and  $\gamma_2$ ), the vector field  $J_s$ , defined by

$$J_s(t) = \frac{\partial}{\partial s} \Sigma(t, s) \in T_{\Sigma(t, s)} M$$

is a Jacobi field along  $\Sigma(\cdot, s)$ ,  $s \in [0, 1]$ , see Bao-Chern-Shen [23, p. 130]. In particular, we have  $\Sigma(1, s) = \gamma(s)$ ,  $J_s(0) = 0$ ,  $J_s(1) = \frac{\partial}{\partial s} \Sigma(1, s) = \frac{d\gamma}{ds}$  and  $J_s(\frac{1}{2}) = \frac{\partial}{\partial s} \Sigma(\frac{1}{2}, s)$ .

Now, we fix  $s \in [0, 1]$ . Since  $J_s(0) = 0$  and the flag curvature in non-positive, then the geodesic  $\Sigma(\cdot, s)$  has no conjugated points, see Theorem 13.2(a). Therefore,

$$J_s(t) \neq 0 \quad \text{for all } t \in (0, 1].$$

Hence  $g_{J_s}(J_s, J_s)(t)$  is well defined for every  $t \in (0, 1]$ . Moreover,

$$F(J_s)(t) := F(\Sigma(t, s), J_s(t)) = [g_{J_s}(J_s, J_s)]^{\frac{1}{2}}(t) \neq 0, \quad \forall t \in (0, 1]. \quad (13.12)$$

Let  $T_s$  the velocity field of  $\Sigma(\cdot, s)$ . Applying twice formula (13.4), we obtain

$$\begin{aligned} \frac{d^2}{dt^2} [g_{J_s}(J_s, J_s)]^{\frac{1}{2}}(t) &= \frac{d^2}{dt^2} F(J_s)(t) = \frac{d}{dt} \left[ \frac{g_{J_s}(D_{T_s} J_s, J_s)}{F(J_s)} \right] (t) = \\ &= \frac{[g_{J_s}(D_{T_s} D_{T_s} J_s, J_s) + g_{J_s}(D_{T_s} J_s, D_{T_s} J_s)] \cdot F(J_s) - g_{J_s}^2(D_{T_s} J_s, J_s) \cdot F(J_s)^{-1}}{F^2(J_s)}(t) = \\ &= \frac{g_{J_s}(D_{T_s} D_{T_s} J_s, J_s) \cdot F^2(J_s) + g_{J_s}(D_{T_s} J_s, D_{T_s} J_s) \cdot F^2(J_s) - g_{J_s}^2(D_{T_s} J_s, J_s)}{F^3(J_s)}(t), \end{aligned}$$

where the covariant derivatives (for generic Finsler manifolds) are with reference vector  $J_s$ . Since  $(M, F)$  is a Berwald space, the Chern connection coefficients do not depend on the direction, i.e., the notion of reference vector becomes irrelevant. Therefore, we can use the Jacobi equation (13.6), concluding that

$$g_{J_s}(D_{T_s} D_{T_s} J_s, J_s) = -g_{J_s}(R(J_s, T_s)T_s, J_s).$$

Using the symmetry property of the curvature tensor, the formula of the flag curvature, and the Schwarz inequality we have

$$\begin{aligned} -g_{J_s}(R(J_s, T_s)T_s, J_s) &= -g_{J_s}(R(T_s, J_s)J_s, T_s) \\ &= -K(J_s, T_s) \cdot [g_{J_s}(J_s, J_s)g_{J_s}(T_s, T_s) - g_{J_s}^2(J_s, T_s)] \\ &\geq 0. \end{aligned}$$

For the last two terms of the numerator we apply again the Schwarz inequality and we conclude that

$$\frac{d^2}{dt^2} F(J_s)(t) \geq 0 \quad \text{for all } t \in (0, 1].$$

Since  $J_s(t) \neq 0$  for  $t \in (0, 1]$ , the mapping  $t \mapsto F(J_s)(t)$  is  $C^\infty$  on  $(0, 1]$ . From the above inequality and the second order Taylor expansion about  $v \in (0, 1]$ , we obtain

$$F(J_s)(v) + (t - v) \frac{d}{dt} F(J_s)(v) \leq F(J_s)(t) \quad \text{for all } t \in (0, 1]. \quad (13.13)$$

Letting  $t \rightarrow 0$  and  $v = 1/2$  in (13.13), by the continuity of  $F$ , we obtain

$$F(J_s)\left(\frac{1}{2}\right) - \frac{1}{2} \frac{d}{dt} F(J_s)\left(\frac{1}{2}\right) \leq 0.$$

Let  $v = 1/2$  and  $t = 1$  in (13.13), and adding the obtained inequality with the above one, we conclude that

$$2F\left(\Sigma\left(\frac{1}{2}, s\right), \frac{\partial}{\partial s} \Sigma\left(\frac{1}{2}, s\right)\right) = 2F(J_s)\left(\frac{1}{2}\right) \leq F(J_s)(1) = F\left(\gamma(s), \frac{d\gamma}{ds}\right).$$

Integrating the last inequality with respect to  $s$  from 0 to 1, we obtain

$$\begin{aligned} 2L_F\left(\Sigma\left(\frac{1}{2}, \cdot\right)\right) &= 2 \int_0^1 F\left(\Sigma\left(\frac{1}{2}, s\right), \frac{\partial}{\partial s} \Sigma\left(\frac{1}{2}, s\right)\right) ds \\ &\leq \int_0^1 F\left(\gamma(s), \frac{d\gamma}{ds}\right) ds \\ &= L_F(\gamma) \\ &= d_F(\gamma_1(1), \gamma_2(1)). \end{aligned}$$

Since  $\Sigma(\frac{1}{2}, 0) = \gamma_1(\frac{1}{2})$ ,  $\Sigma(\frac{1}{2}, 1) = \gamma_2(\frac{1}{2})$  and  $\Sigma(\frac{1}{2}, \cdot)$  is a  $C^\infty$  curve, by the

definition of the metric function  $d_F$ , we conclude that  $\gamma_1$  and  $\gamma_2$  satisfy the Busemann NPC inequality. This concludes the proof.  $\square$

**Remark 13.2** In view of Theorem 13.4, Berwald spaces seem to be the first class of Finsler metrics that are non-positively curved in the sense of Busemann and which are neither flat nor Riemannian. Moreover, the above result suggests a full characterization of the Busemann curvature notion for Berwald spaces. Indeed, we refer the reader to Kristály-Kozma [173] where the converse of Theorem 13.4 is also proved; here we omit this technical part since only the above result is applied for Economical problems.

**Remark 13.3** Note that Theorem 13.4 includes a partial answer to the question of Busemann [55] (see also [238, p. 87]), i.e., every reversible Berwald space of non-positive flag curvature has convex capsules (i.e., the loci equidistant to geodesic segments).

**Remark 13.4** In the fifties, Aleksandrov introduced independently an other notion of curvature in metric spaces, based on the convexity of the distance function. It is well-known that the condition of Busemann curvature is weaker than the Aleksandrov one, see Jost [152, Corollary 2.3.1]. Nevertheless, in Riemannian spaces the Aleksandrov curvature condition holds if and only if the sectional curvature is non-positive, see Bridson-Haefliger [53, Theorem 1A.6], but in the Finsler case the picture is quite rigid. Namely, if on a reversible Finsler manifold  $(M, F)$  the Aleksandrov curvature condition holds (on the induced metric space by  $(M, F)$ ) then  $(M, F)$  it must be Riemannian, see Bridson-Haefliger [53, Proposition 1.14].

A direct consequence of Theorem 13.4 is

**Corollary 13.1** *Let  $(M, F)$  be a forward/backward geodesically complete, simply connected Berwald space with non-positive flag curvature, where  $F$  is positively (but perhaps not absolutely) homogeneous of degree one. Then  $(M, d_F)$  is a global Busemann NPC space, i.e., the Busemann NPC inequality holds for any pair of geodesics.*

The following result is crucial in Chapters 14 and 15.

**Proposition 13.5** *Let  $(M, F)$  be a forward/backward geodesically complete, simply connected Berwald space with non-positive flag curvature, where  $F$  is positively (but perhaps not absolutely) homogeneous of degree one. Fix two geodesics  $\gamma_1, \gamma_2 : [0, 1] \rightarrow M$ . Then, the function  $t \mapsto d_F(\gamma_1(t), \gamma_2(t))$  is convex.*

*Proof* Due to Hopf-Rinow theorem (see Theorem 13.1), there exists a geodesic  $\gamma_3 : [0, 1] \rightarrow M$  joining  $\gamma_1(0)$  and  $\gamma_2(1)$ . Moreover, due to Cartan-Hadamard theorem (see Theorem 13.2),  $\gamma_3$  is unique. Applying Corollary 13.1 first to the pair  $\gamma_1, \gamma_3$  and then to the pair  $\gamma_3, \gamma_2$  (with opposite orientation), we obtain

$$d_F(\gamma_1(\frac{1}{2}), \gamma_3(\frac{1}{2})) \leq \frac{1}{2}d_F(\gamma_1(1), \gamma_3(1));$$

$$d_F(\gamma_3(\frac{1}{2}), \gamma_2(\frac{1}{2})) \leq \frac{1}{2}d_F(\gamma_3(0), \gamma_2(0)).$$

Note that the opposite of  $\gamma_3$  and  $\gamma_2$  are also geodesics, since  $(M, F)$  is of Berwald type, see [23, Example 5.3.3]. Now, using the triangle inequality, we obtain

$$d_F(\gamma_1(\frac{1}{2}), \gamma_2(\frac{1}{2})) \leq \frac{1}{2}d_F(\gamma_1(1), \gamma_2(1)) + \frac{1}{2}d_F(\gamma_1(0), \gamma_2(0)),$$

which means actually the  $\frac{1}{2}$ -convexity of the function  $t \mapsto d_F(\gamma_1(t), \gamma_2(t))$ . The continuity of the function  $t \mapsto d_F(\gamma_1(t), \gamma_2(t))$  together with the  $\frac{1}{2}$ -convexity lead to its convexity.  $\square$

### 13.3 Variational inequalities

Existence results for Nash equilibria are often derived from intersection theorems, KKM theorems, or fixed point theorems. For instance, the original proof of Nash concerning equilibrium points is based on the Brouwer fixed point theorem. These theorems are actually equivalent to minimax theorems or variational inequalities, as Fan minimax theorem, Sion-type result, etc. In this section we recall a few results which will be used in Chapter 16.

A nonempty set  $X$  is *acyclic* if it is connected and its Čech homology (coefficients in a fixed field) is zero in dimensions greater than zero. Note that every contractible set is acyclic (but the converse need not hold in general). The following result is a Fan-type minimax theorem.

**Theorem 13.5** [McClendon [210, Theorem 3.1]] *Suppose that  $X$  is a compact acyclic finite-dimensional ANR. Suppose  $h : X \times X \rightarrow \mathbb{R}$  is a function such that  $\{(x, y) : h(y, y) > h(x, y)\}$  is open and  $\{x : h(y, y) > h(x, y)\}$  is contractible or empty for all  $y \in X$ . Then there is a  $y_0 \in X$  with  $h(y_0, y_0) \leq h(x, y_0)$  for all  $x \in X$ .*

In the sequel, we state some well-known results from the theory of variational inequalities. To do this, we consider a Gâteaux differentiable function  $f : K \rightarrow \mathbb{R}$  where  $K$  is a closed convex subset of the topological vector space  $X$ .

**Lemma 13.1** *Let  $x_0 \in K$  be a relative minimum point of  $f$  to  $K$ , i.e.,  $f(x) \geq f(x_0)$  for every  $x \in K$ . Then,*

$$f'(x_0)(x - x_0) \geq 0, \quad \forall x \in K. \quad (13.14)$$

*Furthermore, if  $f$  is convex, then the converse also holds.*

**Lemma 13.2** *Let  $x_0 \in K$  such that*

$$f'(x)(x - x_0) \geq 0, \quad \forall x \in K. \quad (13.15)$$

*Then,  $x_0 \in K$  is a relative minimum point of  $f$  to  $K$ . Furthermore, if  $f$  is convex, then the converse also holds.*

Note that if we replace  $f'$  by an operator  $A : X \rightarrow X^*$ , then (13.14) appearing in Lemma 13.1 is called a *Stampacchia-type variational inequality*, while (13.15) from Lemma 13.2 is a *Minty-type variational inequality*.

# 14

## Minimization of Cost-functions on Manifolds

Geography has made us  
neighbors. History has made  
us friends. Economics has  
made us partners, and  
necessity has made us allies.  
Those whom God has so  
joined together, let no man  
put asunder.

---

John F. Kennedy (1917–1963)

### 14.1 Introduction

Let us consider three markets  $P_1, P_2, P_3$  placed on an inclined plane (slope) with an angle  $\alpha$  to the horizontal plane, denoted by  $(S_\alpha)$ . Assume that three cars transport products from (resp. to) a deposit  $P \in (S_\alpha)$  to (resp. from) markets  $P_1, P_2, P_3 \in (S_\alpha)$  such that

- they move always in  $(S_\alpha)$  along straight roads;
- the Earth gravity acts on them (we omit other physical perturbations such as friction, air resistance, etc.);
- the transport costs coincide with the *distance* (measuring actually the *time* elapsed to arrive) from (resp. to) deposit  $P$  to (resp. from) markets  $P_i$  ( $i = 1, 2, 3$ ).

We emphasize that usually the two distances, i.e., from the deposit to the markets and conversely, are *not* the same. The point here is that the travel speed depends heavily on both the slope of the terrain and the direction of travel. More precisely, if a car moves with a constant speed

$v$  [ $m/s$ ] on a horizontal plane, it goes  $l_t = vt + \frac{g}{2}t^2 \sin \alpha \cos \theta$  meters in  $t$  seconds on  $(S_\alpha)$ , where  $\theta$  is the angle between the straight road and the direct downhill road ( $\theta$  is measured in clockwise direction). The law of the above phenomenon can be described relatively to the horizontal plane by means of the parametrized function

$$F_\alpha(y_1, y_2) = \frac{y_1^2 + y_2^2}{v\sqrt{y_1^2 + y_2^2} + \frac{g}{2}y_1 \sin \alpha}, \quad (y_1, y_2) \in \mathbb{R}^2 \setminus \{(0, 0)\}. \quad (14.1)$$

Here,  $g \approx 9.81[m/s^2]$ . The *distance* (measuring the *time* to arrive) from  $P = (P^1, P^2)$  to  $P_i = (P_i^1, P_i^2)$  is

$$d_\alpha(P, P_i) = F_\alpha(P_i^1 - P^1, P_i^2 - P^2),$$

and for the converse it is

$$d_\alpha(P_i, P) = F_\alpha(P^1 - P_i^1, P^2 - P_i^2).$$

Consequently, we have to minimize the functions

$$C_f(P) = \sum_{i=1}^3 d_\alpha(P, P_i) \quad \text{and} \quad C_b(P) = \sum_{i=1}^3 d_\alpha(P_i, P), \quad (14.2)$$

when  $P$  moves on  $(S_\alpha)$ . The function  $C_f$  (resp.  $C_b$ ) denotes the *total forward* (resp. *backward*) *cost* between the deposit  $P \in (S_\alpha)$  and markets  $P_1, P_2, P_3 \in (S_\alpha)$ . The minimum points of  $C_f$  and  $C_b$ , respectively, may be far from each other (see Figure 14.1), due to the fact that  $F_\alpha$  (and  $d_\alpha$ ) is not symmetric unless  $\alpha = 0$ , i.e.,  $F_\alpha(-y_1, -y_2) \neq F_\alpha(y_1, y_2)$  for each  $(y_1, y_2) \in \mathbb{R}^2 \setminus \{(0, 0)\}$ .

We will use in general  $T_f$  (resp.  $T_b$ ) to denote a minimum point of  $C_f$  (resp.  $C_b$ ), which corresponds to the position of a deposit when we measure costs in forward (resp. backward) manner, see (14.2).

In the case  $\alpha = 0$  (when  $(S_\alpha)$  is a horizontal plane), the functions  $C_f$  and  $C_b$  coincide (the same is true for  $T_f$  and  $T_b$ ). The minimum point  $T = T_f = T_b$  is the well-known *Torricelli point* corresponding to the triangle  $P_1P_2P_3$ . Note that  $F_0(y_1, y_2) = \sqrt{y_1^2 + y_2^2}/v$  corresponds to the standard Euclidean metric; indeed,

$$d_0(P, P_i) = d_0(P_i, P) = \sqrt{(P_i^1 - P^1)^2 + (P_i^2 - P^2)^2}/v$$

measures the time, which is needed to arrive from  $P$  to  $P_i$  (and vice-versa) with constant velocity  $v$ .

Unfortunately, finding critical points as possible minima does not yield any result: either the minimization function is not smooth enough (usually, it is only a locally Lipschitz function) or the system, which would




Figure 14.1 We fix  $P_1 = (-250, -50)$ ,  $P_2 = (0, -100)$  and  $P_3 = (-50, 100)$  on the slope  $(S_\alpha)$  with angle  $\alpha = 35^\circ$ . If  $v = 10$ , the minimum of the total forward cost on the slope is  $C_f \approx 40.3265$ ; the corresponding deposit is located at  $T_f \approx (-226.11, -39.4995) \in (S_\alpha)$ . However, the minimum of the total backward cost on the slope is  $C_b \approx 38.4143$ ; the corresponding deposit has the coordinates  $T_b \approx (-25.1332, -35.097) \in (S_\alpha)$ .

give the critical points, becomes very complicated even in quite simple cases (see (14.5) below). Consequently, the main purpose of the present chapter is to study the set of these minima (existence, location) in various geometrical settings.

Note, that the function appearing in (14.1) is a typically Finsler metric on  $\mathbb{R}^2$ , introduced and studied first by Matsumoto [207]. In this way, elements from Riemann-Finsler geometry are needed in order to handle the question formulated above. In the next sections we prove some necessarily, existence, uniqueness and multiplicity results for the economical problem on non-positively curved Berwald space which model various real life phenomena. Simultaneously, relevant numerical examples and counterexamples are constructed by means of evolutionary methods and computational geometry tools, emphasizing the applicability and sharpness of our results.

## 14.2 A necessary condition

Let  $(M, F)$  be an  $m$ -dimensional connected Finsler manifold, where  $F$  is positively (but perhaps not absolutely) homogeneous of degree one.

In this section we prove some results concerning the set of minima for functions

$$C_f(P_i, n, s)(P) = \sum_{i=1}^n d_F^s(P, P_i) \quad \text{and} \quad C_b(P_i, n, s)(P) = \sum_{i=1}^n d_F^s(P_i, P),$$

where  $s \geq 1$  and the points  $P_i \in M$ ,  $i = 1, \dots, n$ , correspond to  $n \in \mathbb{N}$  markets. The value  $C_f(P_i, n, s)(P)$  (resp.  $C_b(P_i, n, s)(P)$ ) denotes the *total  $s$ -forward* (resp.  *$s$ -backward*) cost between the deposit  $P \in M$  and the markets  $P_i \in M$ ,  $i = 1, \dots, n$ . When  $s = 1$ , we simply say *total forward* (resp. *backward*) cost.

By using the triangle inequality, for every  $x_0, x_1, x_2 \in M$  we have

$$|d_F(x_1, x_0) - d_F(x_2, x_0)| \leq \max\{d_F(x_1, x_2), d_F(x_2, x_1)\}. \quad (14.3)$$

Given any point  $P \in M$ , there exists a coordinate map  $\varphi_P$  defined on the closure of some precompact open subset  $U$  containing  $P$  such that  $\varphi_P$  maps the set  $U$  diffeomorphically onto the open Euclidean ball  $B^m(r)$ ,  $r > 0$ , with  $\varphi_P(P) = 0_{\mathbb{R}^m}$ . Moreover, there is a constant  $c > 1$ , depending only on  $P$  and  $U$  such that

$$c^{-1} \|\varphi_P(x_1) - \varphi_P(x_2)\| \leq d_F(x_1, x_2) \leq c \|\varphi_P(x_1) - \varphi_P(x_2)\| \quad (14.4)$$

for every  $x_1, x_2 \in U$ ; see [23, p. 149]. Here,  $\|\cdot\|$  denotes the Euclidean norm on  $\mathbb{R}^m$ . We claim that for every  $Q \in M$ , the function  $d_F(\varphi_P^{-1}(\cdot), Q)$  is a Lipschitz function on  $\varphi_P(U) = B^m(r)$ . Indeed, for every  $y_i = \varphi_P(x_i) \in \varphi_P(U)$ ,  $i = 1, 2$ , due to (14.3) and (14.4), one has

$$\begin{aligned} |d_F(\varphi_P^{-1}(y_1), Q) - d_F(\varphi_P^{-1}(y_2), Q)| &= |d_F(x_1, Q) - d_F(x_2, Q)| \leq \\ &\leq \max\{d_F(x_1, x_2), d_F(x_2, x_1)\} \leq c \|y_1 - y_2\|. \end{aligned}$$

Consequently, for every  $Q \in M$ , there exists the *generalized gradient* of the locally Lipschitz function  $d_F(\varphi_P^{-1}(\cdot), Q)$  on  $\varphi_P(U) = B^m(r)$ , see Clarke [72, p. 27], i.e., for every  $y \in \varphi_P(U) = B^m(r)$  we have

$$\partial d_F(\varphi_P^{-1}(\cdot), Q)(y) = \{\xi \in \mathbb{R}^m : d_F^0(\varphi_P^{-1}(\cdot), Q)(y; h) \geq \langle \xi, h \rangle, \forall h \in \mathbb{R}^m\},$$

where  $\langle \cdot, \cdot \rangle$  denotes the standard inner product on  $\mathbb{R}^m$  and

$$d_F^0(\varphi_P^{-1}(\cdot), Q)(y; h) = \limsup_{z \rightarrow y, t \rightarrow 0^+} \frac{d_F(\varphi_P^{-1}(z + th), Q) - d_F(\varphi_P^{-1}(z), Q)}{t}$$

is the *generalized directional derivative*.

**Theorem 14.1** (Necessary Condition) *Assume that  $T_f \in M$  is a minimum point for  $C_f(P_i, n, s)$  and  $\varphi_{T_f}$  is a map as above. Then*

$$0_{\mathbb{R}^m} \in \sum_{i=1}^n d_F^{s-1}(T_f, P_i) \partial d_F(\varphi_{T_f}^{-1}(\cdot), P_i)(\varphi_{T_f}(T_f)). \quad (14.5)$$

*Proof* Since  $T_f \in M$  is a minimum point of the locally Lipschitz function  $C_f(P_i, n, s)$ , then

$$0_{\mathbb{R}^m} \in \partial \left( \sum_{i=1}^n d_F^s(\varphi_{T_f}^{-1}(\cdot), P_i) \right) (\varphi_{T_f}(T_f)).$$

Now, using the basic properties of the generalized gradient, see Appendix D, we conclude the proof.  $\square$

**Remark 14.1** A result similar to Theorem 14.1 can also be obtained for  $C_b(P_i, n, s)$ .

**Example 14.1** Let  $M = \mathbb{R}^m$ ,  $m \geq 2$ , be endowed with the natural Euclidean metric. Taking into account (14.5), a simple computation shows that the unique minimum point  $T_f = T_b$  (i.e., the place of the deposit) for  $C_f(P_i, n, 2) = C_b(P_i, n, 2)$  is the centre of gravity of markets  $\{P_1, \dots, P_n\}$ , i.e.,  $\frac{1}{n} \sum_{i=1}^n P_i$ . In this case,  $\varphi_{T_f}$  can be the identity map on  $\mathbb{R}^m$ .

**Remark 14.2** The system (14.5) may become very complicated even for simple cases; it is enough to consider the Matsumoto metric given by (14.1). In such cases, we are not able to give an explicit formula for minimal points.

### 14.3 Existence and uniqueness results

The next result gives an alternative concerning the number of minimum points of the function  $C_f(P_i, n, s)$  in a general geometrical framework. Similar result can be obtained for  $C_b(P_i, n, s)$ . We have the following theorem.

**Theorem 14.2** *Let  $(M, F)$  be a simply connected, geodesically complete Berwald manifold of nonpositive flag curvature, where  $F$  is positively (but perhaps not absolutely) homogeneous of degree one. Then*

- (a). *there exists either a unique or infinitely many minimum points for  $C_f(P_i, n, 1)$ ;*
- (b). *there exists a unique minimum point for  $C_f(P_i, n, s)$  whenever  $s > 1$ .*

*Proof* First of all, we observe that  $M$  is not a backward bounded set. Indeed, if we assume the contrary, then  $M$  is compact due to Hopf-Rinow theorem, see Theorem 13.1. On the other hand, due Cartan-Hadamard theorem, see Theorem 13.2, the exponential map  $\exp_p : T_pM \rightarrow M$  is a diffeomorphism for every  $p \in M$ . Thus, the tangent space  $T_pM = \exp_p^{-1}(M)$  is compact, a contradiction. Since  $M$  is not backward bounded, in particular, for every  $i = 1, \dots, n$ , we have that

$$\sup_{P \in M} d_F(P, P_i) = \infty.$$

Consequently, outside of a large backward bounded subset of  $M$ , denoted by  $M_0$ , the value of  $C_f(P_i, n, s)$  is large. But,  $M_0$  being compact, the continuous function  $C_f(P_i, n, s)$  attains its infimum, i.e., the set of the minima for  $C_f(P_i, n, s)$  is always nonempty.

On the other hand, due to Proposition 13.5, for every nonconstant geodesic  $\sigma : [0, 1] \rightarrow M$  and  $p \in M$ , the function  $t \mapsto d_F(\sigma(t), p)$  is convex and  $t \mapsto d_F^s(\sigma(t), p)$  is strictly convex, whenever  $s > 1$ , see also Jost [152, Corollary 2.2.6].

(a) Let us assume that there are at least two minimum points for  $C_f(P_i, n, 1)$ , denoting them by  $T_f^0$  and  $T_f^1$ . Let  $\sigma : [0, 1] \rightarrow M$  be a geodesic with constant Finslerian speed such that  $\sigma(0) = T_f^0$  and  $\sigma(1) = T_f^1$ . Then, for every  $t \in (0, 1)$  we have

$$\begin{aligned} C_f(P_i, n, 1)(\sigma(t)) &= \sum_{i=1}^n d_F(\sigma(t), P_i) \\ &\leq (1-t) \sum_{i=1}^n d_F(\sigma(0), P_i) + t \sum_{i=1}^n d_F(\sigma(1), P_i) \\ &= (1-t) \min C_f(P_i, n, 1) + t \min C_f(P_i, n, 1) \\ &= \min C_f(P_i, n, 1). \end{aligned} \tag{14.6}$$

Consequently, for every  $t \in [0, 1]$ ,  $\sigma(t) \in M$  is a minimum point for  $C_f(P_i, n, 1)$ .

(b) It follows directly from the strict convexity of the function  $t \mapsto d_F^s(\sigma(t), p)$ , whenever  $s > 1$ ; indeed, in (14.6) we have strict inequality instead of " $\leq$ " which shows we cannot have more than one minimum point of the function  $C_f(P_i, n, s)$ .  $\square$

**Example 14.2** Let  $F$  be the Finsler metric introduced in (14.1). One




Figure 14.2 A hexagon with vertices  $P_1, P_2, \dots, P_6$  in the Matsumoto space. Increasing the slope's angle  $\alpha$  from  $0$  to  $\pi/2$ , points  $T_f^\alpha$  and  $T_b^\alpha$  are wandering in the presented directions. Orbits of points  $T_f^\alpha$  and  $T_b^\alpha$  were generated by natural cubic spline curve interpolation.

can see that  $(\mathbb{R}^2, F)$  is a typically nonsymmetric Finsler manifold. Actually, it is a (locally) Minkowski space, so a Berwald space as well; its Chern connection vanishes, see [23, p.384]. According to (13.3) and (13.10), the geodesics are straight lines (hence  $(\mathbb{R}^2, F)$  is geodesically complete in both sense) and the flag curvature is identically 0. Thus, we can apply Theorem 14.2. For instance, if we consider the points  $P_1 = (a, -b) \in \mathbb{R}^2$  and  $P_2 = (a, b) \in \mathbb{R}^2$  with  $b \neq 0$ , the minimum points of the function  $C_f(P_i, 2, 1)$  form the segment  $[P_1, P_2]$ , *independently* of the value of  $\alpha$ . The same is true for  $C_b(P_i, 2, 1)$ . However, considering more complicated constellations, the situation changes dramatically, see Figure 14.2.

It would be interesting to study in similar cases the precise orbit of

the (Torricelli) points  $T_f^\alpha$  and  $T_b^\alpha$  when  $\alpha$  varies from 0 to  $\pi/2$ . Several numerical experiments show that  $T_f^\alpha$  tends to a top point of the convex polygon (as in the Figure 14.2).

In the sequel, we want to study our problem in a special constellation: we assume the markets are situated on a common "straight line", i.e., on a geodesic which is in a Riemannian manifold. Note that, in the Riemannian context, the forward and backward costs coincide, i.e.,

$$C_f(P_i, n, 1) = C_b(P_i, n, 1).$$

We denote this common value by  $C(P_i, n, 1)$ . We have

**Theorem 14.3** *Let  $(M, g)$  be a Hadamard-type Riemannian manifold. Assume the points  $P_i \in M$ ,  $i = 1, \dots, n$ , ( $n \geq 2$ ), belong to a geodesic  $\sigma : [0, 1] \rightarrow M$  such that  $P_i = \sigma(t_i)$  with  $0 \leq t_1 < \dots < t_n \leq 1$ . Then*

- (a). *the unique minimum point for  $C(P_i, n, 1)$  is  $P_{[n/2]}$  whenever  $n$  is odd;*
- (b). *the minimum points for  $C(P_i, n, 1)$  is the whole geodesic segment situated on  $\sigma$  between  $P_{n/2}$  and  $P_{n/2+1}$  whenever  $n$  is even.*

*Proof* Since  $(M, g)$  is complete, we extend  $\sigma$  to  $(-\infty, \infty)$ , keeping the same notation. First, we prove that the minimum point(s) for  $C(P_i, n, 1)$  belong to the geodesic  $\sigma$ . We assume the contrary, i.e., let  $T \in M \setminus \text{Im}(\sigma)$  be a minimum point of  $C(P_i, n, 1)$ . Let  $T_\perp \in \text{Im}(\sigma)$  be the projection of  $T$  on the geodesic  $\sigma$ , i.e.

$$d_g(T, T_\perp) = \min_{t \in \mathbb{R}} d_g(T, \sigma(t)).$$

It is clear that the (unique) geodesic lying between  $T$  and  $T_\perp$  is perpendicular to  $\sigma$  with respect to the Riemannian metric  $g$ .

Let  $i_0 \in \{1, \dots, n\}$  such that  $P_{i_0} \neq T_\perp$ . Applying the cosine inequality, see Theorem 13.3, for the triangle with vertices  $P_{i_0}$ ,  $T$  and  $T_\perp$  (so,  $\widehat{T_\perp} = \pi/2$ ), we have

$$d_g^2(T_\perp, T) + d_g^2(T_\perp, P_{i_0}) \leq d_g^2(T, P_{i_0}).$$

Since

$$d_g(T_\perp, T) > 0,$$

we have

$$d_g(T_\perp, P_{i_0}) < d_g(T, P_{i_0}).$$

Consequently,

$$C(P_i, n, 1)(T_\perp) = \sum_{i=1}^n d_g(T_\perp, P_i) < \sum_{i=1}^n d_g(P, P_i) = \min C(P_i, n, 1),$$

a contradiction. Now, conclusions (a) and (b) follow easily by using simple arithmetical reasons.  $\square$

#### 14.4 Examples on the Finslerian-Poincaré disc

We emphasize that Theorem 14.3 is sharp in the following sense: neither the nonpositivity of the sectional curvature (see Example 14.3) nor the Riemannian structure (see Example 14.4) can be omitted.

**Example 14.3** (Sphere) Let us consider the 2-dimensional unit sphere  $\mathbb{S}^2 \subset \mathbb{R}^3$  endowed with its natural Riemannian metric  $h$  inherited by  $\mathbb{R}^3$ . We know that it has constant curvature 1. Let us fix  $P_1, P_2 \in \mathbb{S}^2$ ,  $P_1 \neq \pm P_2$  and their antipodals  $P_3 = -P_1$ ,  $P_4 = -P_2$ . There exists a unique great circle (geodesic) connecting  $P_i$ ,  $i = 1, \dots, 4$ . However, we observe that the function  $C(P_i, 4, 1)$  is *constant* on  $\mathbb{S}^2$ ; its value is  $2\pi$ . Consequently, *every* point on  $\mathbb{S}^2$  is a minimum for the function  $C(P_i, 4, 1)$ .

**Example 14.4** (Finslerian-Poincaré disc) Let us consider the disc

$$M = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 < 4\}.$$

Introducing the polar coordinates  $(r, \theta)$  on  $M$ , i.e.,  $x = r \cos \theta$ ,  $y = r \sin \theta$ , we define the non-reversible Finsler metric on  $M$  by

$$F((r, \theta), V) = \frac{1}{1 - \frac{r^2}{4}} \sqrt{p^2 + r^2 q^2} + \frac{pr}{1 - \frac{r^4}{16}},$$

where

$$V = p \frac{\partial}{\partial r} + q \frac{\partial}{\partial \theta} \in T_{(r, \theta)} M.$$

The pair  $(M, F)$  is the so-called *Finslerian-Poincaré disc*. Within the classification of Finsler manifolds,  $(M, F)$  is a *Randers space*, see Bao-Chern-Shen [23, Section 12.6], which has the following properties:

- (p1). it has constant negative flag curvature  $-1/4$ ;
- (p2). the geodesics have the following trajectories: Euclidean circular arcs that intersect the boundary  $\partial M$  of  $M$  at Euclidean right angles; Euclidean straight rays that emanate from the origin; and Euclidean straight rays that aim to the origin;




Figure 14.3

*Step 1:* The minimum of the total backward (resp. forward) cost function  $C_b(P_i, 2, 1)$  (resp.  $C_f(P_i, 2, 1)$ ) is restricted to the geodesic determined by  $P_1(1.6, 170^\circ)$  and  $P_2(1.3, 250^\circ)$ . The point which minimizes  $C_b(P_i, 2, 1)$  (resp.  $C_f(P_i, 2, 1)$ ) is approximated by  $P_b(0.8541, 212.2545^\circ)$  (resp.  $P_f = P_1$ ); in this case  $C_b(P_i, 2, 1)(P_b) \approx 1.26$  (resp.  $C_f(P_i, 2, 1)(P_f) \approx 2.32507$ ).  
*Step 2:* The minimum of the total backward (resp. forward) cost function  $C_b(P_i, 2, 1)$  (resp.  $C_f(P_i, 2, 1)$ ) is on the whole Randers space  $M$ . The minimum point of total backward (resp. forward) cost function is approximated by  $T_b(0.4472, 212.5589^\circ)$  (resp.  $T_f(1.9999, 171.5237^\circ)$ ), which gives  $C_b(P_i, 2, 1)(T_b) \approx 0.950825 < C_b(P_i, 2, 1)(P_b)$  (resp.  $C_f(P_i, 2, 1)(T_f) \approx 2.32079 < C_f(P_i, 2, 1)(P_f)$ ).

(p3).  $\text{dist}_F((0,0), \partial M) = \infty$ , while  $\text{dist}_F(\partial M, (0,0)) = \log 2$ .

Although  $(M, F)$  is forward geodesically complete (but *not* backward geodesically complete), it has constant negative flag curvature  $-\frac{1}{4}$  and it is contractible (thus, simply connected), the conclusion of Theorem 14.3 may be false. Indeed, one can find points in  $M$  (belonging to the same geodesic) such that the minimum point for the total forward (resp. backward) cost function is *not* situated on the geodesic, see Figure 14.3.

**Remark 14.3** Note that Example 14.4 (Finslerian-Poincaré disc) may give a model of a *gravitational field* whose centre of gravity is located at the origin  $O = (0, 0)$ , while the boundary  $\partial M$  means the "infinity". Suppose that in this gravitational field, we have several spaceships, which are delivering some cargo to certain bases or to another spacecraft. Also, assume that these spaceships are of the same type and they consume  $k$  liter/second fuel ( $k > 0$ ). Note that the expression  $F(d\sigma)$  denotes the

physical *time* elapsed to traverse a short portion  $d\sigma$  of the spaceship orbit. Consequently, traversing a short path  $d\sigma$ , a spaceship consumes  $kF(d\sigma)$  liter of fuel. In this way, the number  $k \int_0^1 F(\sigma(t), d\sigma(t)) dt$  expresses the quantity of fuel used up by a spaceship traversing an orbit  $\sigma : [0, 1] \rightarrow M$ .

Suppose that two spaceships have to meet each other (for logistical reasons) starting their trip from bases  $P_1$  and  $P_2$ , respectively. Consuming as low total quantity of fuel as possible, they will choose  $T_b$  as a meeting point and *not*  $P_b$  on the geodesic determined by  $P_1$  and  $P_2$ . Thus, the point  $T_b$  could be a position for an optimal deposit-base.

Now, suppose that we have two damaged spacecraft (e.g., without fuel) at positions  $P_1$  and  $P_2$ . Two rescue spaceships consuming as low total quantity of fuel as possible, will blastoff from base  $T_f$  and not from  $P_f = P_1$  on the geodesic determined by  $P_1$  and  $P_2$ . In this case, the point  $T_f$  is the position for an optimal rescue-base. If the spaceships in trouble are close to the center of the gravitational field  $M$ , then any rescue-base located closely also to the center  $O$ , implies the consumption of a great amount of energy (fuel) by the rescue spaceships in order to reach their destinations (namely,  $P_1$  and  $P_2$ ). Indeed, they have to overcome the strong gravitational force near the center  $O$ . Consequently, this is the reason why the point  $T_f$  is so far from  $O$ , as Figure 14.3 shows. Note that further numerical experiments support this observation. However, there are certain special cases when the position of the optimal rescue-base is either  $P_1$  or  $P_2$ : from these two points, the farthest one from the gravitational center  $O$  will be the position of the rescue-base. In such case, the orbit of the (single) rescue spaceship is exactly the geodesic determined by points  $P_1$  and  $P_2$ .

## 14.5 Comments and further perspectives

*A. Comments.* The results of this chapter are based on the paper of Kristály-Kozma [173] and Kristály-Moroşanu-Róth [175]. In this chapter we studied variational problems arising from Economical contexts via Riemann-Finsler geometry. Real life phenomena may be well modeled involving Finsler metrics, which represent external force as current or gravitation. A special class of Finsler manifolds, called Berwald space, played a central role in our investigation. Indeed, a recent result of Kristály-Kozma [173] concerning metric relations on non-positively curved Berwald spaces has been exploited which includes several real




Figure 14.4 The force of the current flow towards the waterfall may be described by means of the Finslerian-Poincaré disc model. The points  $x_1(0)$  and  $x_2(0)$  correspond to the alerting moment when the lifeboats from the mother-ship  $M_S$  start their trips, while  $x_1(T)$  and  $x_2(T)$  are the last possible points where the lifeboats and the corresponding pleasure boats may meet each other.

life applications (as the slope metric of a hillside, described by Matsumoto [207]).

*B. Further perspectives.* We propose an optimization problems which arises in a real life situation.

**Problem 14.1** *There are given  $n$  ships moving on different paths (we know all the details on their speed, direction, etc.). Determine the position of the optimum point(s) of the aircraft-carrier (mother ship) from where other  $n$  ships can reach the first  $n$  ships within a given time interval using the lowest amount of (total) combustible.*

We give some hints concerning Problem 14.1. We consider it in a particular case as follows. Two pleasure boats are moving on given paths within an estuary ending in a waterfall (see Figure 14.4), and a mother ship  $M_S$  is positioned in the same area for safety reasons transporting two lifeboats. The problem is to determine the optimal position of the mother ship  $M_S$  at every moment such that the two lifeboats reach the two pleasure boats within a  $T$  period consuming the minimal total combustible (for a lifeboat it is allowed to wait the pleasure boat but not

conversely). On Figure 14.4, the points  $x_1(0)$  and  $x_2(0)$  correspond to the alerting moment (when the lifeboats start their trips) while  $x_1(T)$  and  $x_2(T)$  are the last possible points where the lifeboats and the corresponding pleasure boats may meet each other. In this case, the external force is the *current flow* towards the waterfall; the law describing this force – up to some constants – can be given as a *submanifold* of the Finslerian-Poincaré disc, see Example 14.4.

We now formulate in mathematical terminology a slightly general problem than Problem 14.1. We consider the quasi-metric space  $(M, d_F)$  associated with a Finsler manifold  $(M, F)$  which is not necessarily symmetric, the paths  $x_i : [0, T] \rightarrow M$  ( $i = 1, \dots, n$ ) and some numbers  $v_i > 0$  ( $i = 1, \dots, n$ ) representing the speeds of the last  $n$  ships moving towards the paths  $x_i$  in order to meet the  $i^{\text{th}}$  ship from the first group. Let us consider the function

$$f_i(P) = \min_{t \in [0, T]} \{d_F(P, x_i(t)) : d_F(P, x_i(t)) \leq v_i t\}$$

and the convex functions  $\psi_i : \mathbb{R} \rightarrow \mathbb{R}$  ( $i = 1, \dots, n$ ). The variational problem we are dealing with is

$$\min\{f(P) : P \in M\} \quad (P_{\text{optim}})$$

where

$$f(P) = \sum_{i=1}^n \psi_i(f_i(P)).$$

We assume that

$$S = \bigcap_{i=1}^n \{P \in M : d_F(P, x_i(T)) \leq v_i T\} \neq \emptyset,$$

which is a sufficient condition for the existence of a solution to the variational problem  $(P_{\text{optim}})$ .

# 15

## Best Approximation Problems on Manifolds

If that enabled us to predict the succeeding situation with the same approximation, that is all we require, and we should say that the phenomenon had been predicted, that it is governed by the laws.

---

Henri Poincaré (1854–1912)

### 15.1 Introduction

One of the most famous questions of functional and numerical analysis is the best approximation problem: find the nearest point (called also projection) from a given point to a nonempty set, both situated in an ambient space endowed with a certain metric structure by means of which one can measure metric distances. In the classical theory of best approximations the ambient space has a vector space structure, the distance function is symmetric which comes from a norm or, specially, from an inner product. However, *non*-symmetry is abundant in important real life situations; indeed, it is enough to consider the swimming (with/against a current) in a river, or the walking (up/down) on a mountain slope. These kinds of non-symmetric phenomena lead us to spaces with possible no vector space structure while the distance function is not necessarily symmetric, see also Chapter 14. Thus, the most appropriate framework is to consider not necessarily reversible Finsler

manifolds. This nonlinear context throws completely new light upon the problem of best approximations where well-tried methods usually fail.

The purpose of this chapter is to initiate a systematic study of best approximation problems on Finsler manifolds by exploiting notions from Finsler geometry as geodesics, forward/backward geodesically completeness, flag curvature, fundamental inequality of Finsler geometry, and first variation of arc length.

Throughout this chapter,  $(M, F)$  is a connected finite-dimensional Finsler manifold,  $F$  is positively (but perhaps not absolutely) homogeneous of degree one. Let  $d_F : M \times M \rightarrow [0, \infty)$  be the quasi-metric associated with  $F$ , see (13.8). Let  $q \in M$  be a point and  $S \subset M$  a nonempty set. Since  $d_F$  is not necessarily symmetric, let

$$\text{dist}_F(q, S) = \inf\{d_F(q, s) : s \in S\}$$

and

$$\text{dist}_F(S, q) = \inf\{d_F(s, q) : s \in S\}$$

the *forward* and *backward distances* between the point  $q$  and the set  $S$ , respectively. The *forward* (resp. *backward*) *best approximation problem* can be formulated as follows: find  $s_f \in S$  (resp.  $s_b \in S$ ) such that

$$d_F(q, s_f) = \text{dist}_F(q, S) \quad (\text{resp.} \quad d_F(s_b, q) = \text{dist}_F(S, q)).$$

We define the set of *forward* (resp. *backward*) *projections* of  $q$  to  $S$  by

$$P_S^+(q) = \{s_f \in S : d_F(q, s_f) = \text{dist}_F(q, S)\}$$

$$(\text{resp.} \quad P_S^-(q) = \{s_b \in S : d_F(s_b, q) = \text{dist}_F(S, q)\}).$$

If  $P_S^+(q)$  and  $P_S^-(q)$  coincide (for instance, when the Finsler metric  $F$  is absolutely homogeneous), we simply write  $P_S(q)$  instead of the above sets. If any of the above sets is a singleton, we do not make any difference between the set and its unique point.

In the following sections we are dealing in detail with the existence, characterization and non-expansiveness of projections, geodesic convexity and Chebyshevity of closed sets as well as with nearest points between two sets.

## 15.2 Existence of projections

Let  $(M, F)$  be a Finsler manifold. A set  $S \subset M$  is *forward* (resp. *backward*) *proximal* if  $P_S^+(q) \neq \emptyset$  (resp.  $P_S^-(q) \neq \emptyset$ ) for every  $q \in M$ .

**Theorem 15.1** *Let  $(M, F)$  be a forward (resp. backward) geodesically complete Finsler manifold,  $S \subset M$  be a nonempty closed set. Then,  $S$  is a forward (resp. backward) proximal set.*

*Proof* We consider the statement only for the "forward" case, the "backward" case works similarly. Let us fix  $q \in M$  arbitrarily and let  $\{s_n\}_{n \in \mathbb{N}} \subset S$  be a forward minimizing sequence, i.e.,  $\lim_{n \rightarrow \infty} d_F(q, s_n) = \text{dist}_F(q, S) =: d_0$ . In order to conclude the proof, it is enough to show that the sequence  $\{s_n\}_{n \in \mathbb{N}}$  contains a convergent subsequence. Indeed, let  $\{s_{k_n}\}_n \subset S$  be a convergent subsequence of  $\{s_n\}_n$ . Since  $S$  is closed, then  $\lim_{n \rightarrow \infty} s_{k_n} = s_f \in S$ . Moreover, since  $d_F(q, \cdot)$  is continuous, we have  $d_F(q, s_f) = \lim_{n \rightarrow \infty} d_F(q, s_{k_n}) = \text{dist}_F(q, S)$ , which proves that  $s_f \in P_S^+(q)$ , i.e.,  $S$  is a forward proximal set.

Let  $A = \{s_n : n \in \mathbb{N}\}$  and assume that  $A$  has no any convergent subsequence. Then, for every  $a \in \bar{A}$ , there exists  $r_a > 0$  such that

$$\mathcal{B}_a^+(r_a) \cap A \subseteq \{a\}. \quad (15.1)$$

On the other hand, for  $n$  large enough, we have  $d_F(q, s_n) < d_0 + 1$ ; thus, we may assume that  $A \subset \mathcal{B}_q^+(d_0 + 1)$ . Since the set  $\bar{A}$  is forward bounded and closed, due to Hopf-Rinow's theorem, it is also compact. Since  $\bar{A} \subseteq \cup_{a \in \bar{A}} \mathcal{B}_a^+(r_a)$  and  $\mathcal{B}_a^+(r_a)$  are open balls in the manifold topology, there exist  $a_1, \dots, a_l \in \bar{A}$  such that  $\bar{A} \subseteq \cup_{i=1}^l \mathcal{B}_{a_i}^+(r_{a_i})$ . In particular, for every  $n \in \mathbb{N}$ , there exists  $i_0 \in \{1, \dots, l\}$  such that  $s_n \in \mathcal{B}_{a_{i_0}}^+(r_{a_{i_0}})$ . Due to (15.1),

$$s_n \in \mathcal{B}_{a_{i_0}}^+(r_{a_{i_0}}) \cap A \subseteq \{a_{i_0}\},$$

i.e.,  $s_n = a_{i_0}$ . Consequently, the set  $A$  contains finitely many elements, which contradicts our assumption.  $\square$

The geodesically completeness cannot be dropped in Theorem 15.1. We describe here a concrete example on the Finslerian-Poincaré disc, see Example 14.4. The idea to construct a closed, not backward proximal set in  $(M, F)$  comes from the property (p3) and from the fact that  $(M, F)$  is *not* geodesically backward complete, see Bao-Chern-Shen [23, p.342]. We consider the point  $q = (1, 0) \in M$  and the set  $S = \{(t, -\sqrt{2}) : t \in [1, \sqrt{2}]\} \subset M$ , see Figure 15.1. Clearly,  $S$  is closed in the topology of  $(M, F)$ . However, a numerical calculation based on property (p2) and on Maple codes shows that the function  $t \mapsto d_F((t, -\sqrt{2}), q)$ ,  $t \in [1, \sqrt{2}]$ , is strictly *decreasing*, thus the value

$$\text{dist}_F(S, q) = \lim_{t \rightarrow \sqrt{2}} d_F((t, -\sqrt{2}), q) \approx 0.8808$$




Figure 15.1  $S = \{(t, -\sqrt{2}) : t \in [1, \sqrt{2}]\}$  is a closed, but not backward proximal set on the Finslerian-Poincaré disc.

is *not* achieved by any point of  $S$ , i.e.,  $P_S^-(q) = \emptyset$ .

### 15.3 Geometric properties of projections

Let  $(X, \langle \cdot, \cdot \rangle)$  be a real prehilbert space,  $S \subset X$  be a nonempty convex set,  $x \in X$ ,  $y \in S$ . Due to Moskovitz-Dines [222] one has:

$$y \in P_S(x) \Leftrightarrow \langle x - y, z - y \rangle \leq 0 \quad \text{for all } z \in S. \quad (15.2)$$

The geometrical meaning of this characterization is that the vectors  $x - y$  and  $z - y$  (for every  $z \in S$ ) form an obtuse-angle with respect to the inner product  $\langle \cdot, \cdot \rangle$ .

For a Finsler manifold  $(M, F)$ , the function  $F$  is not necessarily induced by an inner product, thus angle-measuring loses its original meaning. The 'inner product' on a Finsler manifold  $(M, F)$  is the Riemannian metric on the pulled-back bundle  $\pi^*TM$  depending not only on the points of  $M$  but also on the *directions* in  $TM$ , defined by  $g_{(p,y)}$  from (13.1),  $y \in T_pM$ .

Let  $S \subset M$  be a nonempty set and  $q \in M$ . We consider the following statements:

- $(MD_1^+)$  :  $s \in P_S^+(q)$ ;
- $(MD_2^+)$  : If  $\gamma : [0, 1] \rightarrow M$  is the unique minimal geodesic from  $\gamma(0) = q$  to  $\gamma(1) = s \in S$ , then every geodesic  $\sigma : [0, \delta] \rightarrow S$  ( $\delta \geq 0$ ) emanating from the point  $s$  fulfills  $g_{(s, \dot{\gamma}(1))}(\dot{\gamma}(1), \dot{\sigma}(0)) \geq 0$ .

**Theorem 15.2** *Let  $(M, F)$  be a forward geodesically complete Finsler manifold,  $S \subset M$  be a nonempty closed set. Then,  $(MD_1^+) \Rightarrow (MD_2^+)$ .*

*Proof* One may assume that  $\delta > 0$ ; otherwise, the statement is trivial since the curve  $\sigma$  shrinks to a point. Standard ODE theory shows that the geodesic segment  $\sigma : [0, \delta] \rightarrow M$  has a  $C^\infty$  extension  $\tilde{\sigma} : (-\varepsilon, \delta] \rightarrow M$ , where  $\varepsilon > 0$ . Since  $\gamma$  is the unique minimal geodesic lying the points  $q$  and  $s$ , there exists  $v_0 \in T_q M$  such that  $\gamma(t) = \exp_q(tv_0)$  and  $\exp_{q^*}$  is nonsingular at the point  $v_0$ . One may fix  $0 < \varepsilon_0 < \min\{\varepsilon, \delta\}$  and a  $C^\infty$  vector field  $V : [-\varepsilon_0, \varepsilon_0] \rightarrow T_q M$  such that  $V(0) = v_0$ ,  $\exp_q(V(u)) = \tilde{\sigma}(u)$  and  $\exp_{q^*}$  is not singular at  $V(u)$  for every  $u \in [-\varepsilon_0, \varepsilon_0]$ . Let us introduce the map  $\Sigma : [0, 1] \times [-\varepsilon_0, \varepsilon_0] \rightarrow M$  by  $\Sigma(t, u) = \exp_q(tV(u))$ . Note that  $\Sigma$  is well-defined and  $\Sigma$  is a variation of the geodesic  $\gamma$  with  $\Sigma(\cdot, 0) = \gamma$ . By definition, we have

$$\begin{aligned} L'_F(\Sigma(\cdot, 0)) &= \frac{d}{du} L_F(\Sigma(\cdot, u))|_{u=0} = \lim_{u \rightarrow 0^+} \frac{L_F(\Sigma(\cdot, u)) - L_F(\Sigma(\cdot, 0))}{u} \\ &\geq \lim_{u \rightarrow 0^+} \frac{d_F(q, \tilde{\sigma}(u)) - L_F(\gamma)}{u} \\ &= \lim_{u \rightarrow 0^+} \frac{d_F(q, \sigma(u)) - d_F(q, s)}{u}. \end{aligned}$$

Since  $s \in P_S^+(q)$  and  $\sigma(u) \in S$  for small values of  $u \geq 0$ , the latter limit is non-negative; thus,  $L'_F(\Sigma(\cdot, 0)) \geq 0$ . On the other hand, combining this relation with the first variation formula, see (13.7), we obtain

$$0 \leq L'_F(\Sigma(\cdot, 0)) = \frac{g_{(s, \dot{\gamma}(1))}(\dot{\gamma}(1), \dot{\sigma}(0))}{F(s, \dot{\gamma}(1))},$$

which concludes the proof. □

We now consider the above statements for the "backward" case:

$(MD_1^-) : s \in P_S^-(q)$ ;

$(MD_2^-) :$  If  $\gamma : [0, 1] \rightarrow M$  is the unique minimal geodesic from  $\gamma(0) = s \in S$  to  $\gamma(1) = q$ , then every geodesic  $\sigma : [0, \delta] \rightarrow S$  ( $\delta \geq 0$ ) emanating from the point  $s$  fulfills  $g_{(s, \dot{\gamma}(0))}(\dot{\gamma}(0), \dot{\sigma}(0)) \leq 0$ .

The proof of the following result works similarly to that of Theorem 15.2.

**Theorem 15.3** *Let  $(M, F)$  be a backward geodesically complete Finsler manifold,  $S \subset M$  be a nonempty, closed set. Then,  $(MD_1^-) \Rightarrow (MD_2^-)$ .*

**Remark 15.1** Let  $(M, g)$  be a Riemannian manifold. In this special case, since the fundamental tensor  $g$  is independent on the directions, the statements  $(MD_1^+)$  and  $(MD_2^+)$  (as well as  $(MD_1^-)$  and  $(MD_2^-)$ ) reduce to:

$(MD_1)$  :  $s \in P_S(q)$ ;

$(MD_2)$  : If  $\gamma : [0, 1] \rightarrow M$  is the unique minimal geodesic from  $\gamma(0) = s \in S$  to  $\gamma(1) = q$ , then every geodesic  $\sigma : [0, \delta] \rightarrow S$  ( $\delta \geq 0$ ) emanating from the point  $s$  fulfills  $g(\dot{\gamma}(0), \dot{\sigma}(0)) \leq 0$ .

**Remark 15.2** The implications in Theorems 15.2 and 15.3 *cannot* be reversed in general, not even for Riemannian manifolds. Indeed, we consider the  $m$ -dimensional unit sphere  $(\mathbb{S}^m, g_0)$  ( $m \geq 2$ ), and we fix the set  $S$  as the equator of  $\mathbb{S}^m$ . Let also  $q \in \mathbb{S}^m \setminus (S \cup \{N, -N\})$  where  $N$  is the North pole. One can see that hypothesis  $(MD_2)$  holds exactly for *two* points  $s_1 \in S$  and  $s_2 = -s_1 \in S$ ; these points are the intersection points of the equator  $S$  and the plane throughout the points  $q$ ,  $N$  and  $-N$ . Let us fix an order of the points on the great circle:  $N, q, s_1, -N, s_2$ . Then, we have  $\text{dist}_{g_0}(q, S) = d_{g_0}(q, s_1) < \pi/2 < d_{g_0}(q, s_2)$ , thus  $s_1 \in P_S(q)$  but  $s_2 \notin P_S(q)$ .

In spite of Remark 15.2 we have two Moskovitz-Dines type *characterizations*. The first is due to Walter [295] for Hadamard-type Riemannian manifolds.

**Theorem 15.4** *Let  $(M, g)$  be a Hadamard-type Riemannian manifold and let  $S \subset M$  be a nonempty, closed, geodesic convex set. Then  $(MD_1) \Leftrightarrow (MD_2)$ .*

*Proof* Due to Theorem 15.3 we have  $(MD_1) \Rightarrow (MD_2)$ . Now, fix  $z \in S$  arbitrarily. Let  $\sigma : [0, 1] \rightarrow M$  be the unique minimal geodesic with  $\sigma(0) = s$  and  $\sigma(1) = z$ . Since the set  $S$  is geodesic convex, we have  $\sigma([0, 1]) \subseteq S$ . On account of hypotheses  $(MD_2)$ , we have

$$g(\dot{\gamma}(0), \dot{\sigma}(0)) \leq 0. \quad (15.3)$$

Consider now the geodesic triangle determined by vertices  $q, z, s \in M$  and denote by  $\widehat{s}$  the angle belonging to vertex  $s$ . Due to (15.3),  $\cos \widehat{s} \leq 0$ . Thus, applying Theorem 13.3, we have

$$d_g^2(q, s) - d_g^2(q, z) \leq 2d_g(q, s)d_g(z, s) \cos \widehat{s} - d_g^2(z, s) \leq 0,$$

which means  $s \in P_S(q)$ , i.e.,  $(MD_1)$ . This concludes the proof.  $\square$

For not necessarily reversible Minkowski spaces, we may prove the following.

**Theorem 15.5** *Let  $(M, F) = (\mathbb{R}^m, F)$  be a Minkowski space and  $S \subset M$  be a nonempty, closed, geodesic convex set. Then*

- (i)  $(MD_1^+) \Leftrightarrow (MD_2^+)$ ;
- (ii)  $(MD_1^-) \Leftrightarrow (MD_2^-)$ .

*Proof* On account of Theorems 15.2 and 15.3 we have  $(MD_1^+) \Rightarrow (MD_2^+)$  and  $(MD_1^-) \Rightarrow (MD_2^-)$ , respectively. We now prove that  $(MD_2^+) \Rightarrow (MD_1^+)$ ; the implication  $(MD_2^-) \Rightarrow (MD_1^-)$  works in a similar way.

We may assume that  $s \neq q$ . The unique minimal geodesic  $\gamma : [0, 1] \rightarrow M$  from  $\gamma(0) = q$  to  $\gamma(1) = s \in S$  is  $\gamma(t) = q + t(s - q)$ . Fix  $z \in S$ , and define the geodesic segment  $\sigma : [0, 1] \rightarrow M$  by  $\sigma(t) = s + t(z - s)$ . Since  $S$  is geodesic convex, then  $\sigma([0, 1]) \subseteq S$ . Consequently, the inequality from  $(MD_2^+)$  reduces to

$$g_{s-q}(s - q, z - s) \geq 0. \quad (15.4)$$

Relations (15.4) and (13.11) yield

$$\begin{aligned} d_F^2(q, s) &= F^2(s - q) \\ &= g_{s-q}(s - q, s - q) \\ &\leq g_{s-q}(s - q, z - q) \\ &\leq F(s - q) \cdot F(z - q) \\ &= d_F(q, s) \cdot d_F(q, z), \end{aligned}$$

i.e.,  $d_F(q, s) \leq d_F(q, z)$ . Since  $z \in S$  is arbitrarily fixed, we have  $s \in P_S^+(q)$ .  $\square$

## 15.4 Geodesic convexity and Chebyshev sets

If  $(X, \langle \cdot, \cdot \rangle)$  is a finite-dimensional Hilbert space and  $S \subset X$  is a nonempty closed set, the equivalence of the following statements is well-known (see Borwein [44] and Phelps [243]):

- $S$  is convex;
- $S$  is Chebyshev, i.e.,  $\text{card}P_S(x) = 1$  for every  $x \in X$ ;
- $P_S$  is non-expansive, i.e., for every  $x_1, x_2 \in X$ ,

$$\|P_S(x_1) - P_S(x_2)\| \leq \|x_1 - x_2\|.$$

On manifolds, as might be expected, uniqueness of the projection is

influenced not only by the geodesical convexity of the set but also by the *curvature* of the space. The latter is well emphasized by considering the standard  $m$ -dimensional unit sphere  $\mathbb{S}^m$  ( $m \geq 2$ ) with its natural Riemannian metric  $g_0$ , having constant sectional curvature 1. The projection of the North pole  $N \in \mathbb{S}^m$  to a geodesic ball centered at the South pole and radius  $r < \pi/2$  is the *whole* boundary of this geodesic ball. As far as we know, the global Busemann NPC spaces are the unique circumstances where the uniqueness of projections from any point to any geodesic convex set is guaranteed in general, see Jost [152, Section 3.3]. Note that all Hadamard-type Riemannian manifolds belong to this class, see also Udriște [291].

However, uniqueness of projections to generic geodesic convex sets are not known for not necessarily reversible Finsler manifolds with non-positive curvature. In the sequel, we delimit such a class of Finsler manifolds (so, in particular, non-Riemannian manifolds) for which the forward/backward projections are singletons for any geodesic convex set.

Let  $(M, F)$  be a Finsler manifold. A set  $S \subset M$  is a *forward* (resp. *backward*) *Chebyshev* set if  $\text{card}P_S^+(q) = 1$  (resp.  $\text{card}P_S^-(q) = 1$ ) for each  $q \in M$ .

**Theorem 15.6** *Let  $(M, F)$  be a forward (resp. backward) geodesically complete, simply connected Berwald space with non-positive flag curvature, and  $S \subset M$  a nonempty, closed, geodesic convex set. Then,  $S$  is a forward (resp. backward) Chebyshev set.*

*Proof* By Theorem 15.1, we know that  $\text{card}P_S^+(q) \geq 1$  (resp.  $\text{card}P_S^-(q) \geq 1$ ) for every  $q \in M$ . On the other hand, Proposition 13.5 guarantees that  $t \mapsto d_F(\gamma_1(t), \gamma_2(t))$  is convex for any two minimal geodesics  $\gamma_1, \gamma_2 : [0, 1] \rightarrow M$  (with not necessarily a common starting point). In particular, for every  $q \in M$  and any non-constant geodesic  $\gamma : [0, 1] \rightarrow M$ , the function  $t \mapsto d_F^2(q, \gamma(t))$  is strictly convex. Now, assume that  $\text{card}P_S^+(q) > 1$ , i.e., there exists  $s_1, s_2 \in P_S^+(q)$ ,  $s_1 \neq s_2$ . Let  $\gamma : [0, 1] \rightarrow M$  be the minimal geodesic which joins these points. Consequently,  $\text{Im}(\gamma) \subseteq S$  and for every  $0 < t < 1$ , we have

$$d_F^2(q, \gamma(t)) < td_F^2(q, \gamma(1)) + (1-t)d_F^2(q, \gamma(0)) = \text{dist}_F^2(q, S),$$

a contradiction. □

**Remark 15.3** Both Hadamard-type Riemannian manifolds and Minkowski spaces fulfill the hypotheses of Theorem 15.6. Now, we present a typical class of Berwald spaces where Theorem 15.6 applies. Let  $(N, h)$

be an arbitrarily closed hyperbolic Riemannian manifold of dimension at least 2, and  $\varepsilon > 0$ . Let us define the Finsler metric  $F_\varepsilon : T(\mathbb{R} \times N) \rightarrow [0, \infty)$  by

$$F_\varepsilon(t, p; \tau, w) = \sqrt{h_p(w, w) + \tau^2 + \varepsilon \sqrt{h_p^2(w, w) + \tau^4}},$$

where  $(t, p) \in \mathbb{R} \times N$  and  $(\tau, w) \in T_{(t,p)}(\mathbb{R} \times N)$ . Shen [277] pointed out that the pair  $(\mathbb{R} \times N, F_\varepsilon)$  is a reversible Berwald space with non-positive flag curvature which is neither a Riemannian manifold nor a Minkowski space.

**Remark 15.4** There are Finsler manifolds with non-positive flag curvature which are not curved in the sense of Busemann and the function  $t \mapsto d_F(\gamma_1(t), \gamma_2(t))$  is *not* convex; simple examples of these types can be given theoretically on Hilbert geometries (see Socié-Méthou [281]), or numerically on certain Randers spaces (see Shen [275], [276]) by using evolutive programming in the spirit of Kristály-Moroşanu-Róth [175]. Thus, no convexity property of the function  $t \mapsto d_F(\gamma_1(t), \gamma_2(t))$  can be guaranteed in general which raises doubts on the validity of Theorem 15.6 for arbitrary Finsler manifolds with non-positive flag curvature.

**Remark 15.5** The Chebyshev property of a closed set does not imply always its geodesic convexity in arbitrarily Hadamard-type Riemannian manifolds. Indeed, consider the Poincaré upper half-plane model  $\mathbb{H}^2$  (with sectional curvature  $-1$ ) and the closed set  $S = \mathbb{H}^2 \setminus B_{(0,1)}(2)$ , where  $B_{(0,1)}(2)$  is the standard euclidean 2-dimensional open ball with center  $(0, 1) \in \mathbb{R}^2$  and radius 2.  $S$  is a Chebyshev set since every horocycle which is below of the boundary  $\partial S$  and tangent to the same set  $\partial S$ , is situated in the interior of  $B_{(0,1)}(2)$ . However, it is clear that  $S$  is not geodesic convex. For further examples, see Grognet [132, Section 2]. These examples show the converse of Theorem 15.6 does not hold in general. Moreover, the following rigidity result does hold.

**Theorem 15.7** [Busemann [55, p.152]] *Let  $(M, F)$  be a simply-connected, complete, reversible Finsler manifold with non-positive flag curvature. If any Chebyshev set in  $M$  is geodesic convex, then  $(M, F)$  is a Minkowski space.*

The following characterization of geodesic convexity is due to Grognet [132] (see also Udrişte [291] for an alternative proof via the second variation formula of the arc length):

**Theorem 15.8** *Let  $(M, g)$  be a Hadamard-type Riemannian manifold and  $S \subset M$  be a nonempty closed set. The following two statements are equivalent:*

- (i).  $S$  is geodesic convex;
- (ii).  $P_S$  is non-expansive, i.e.,  $d_g(P_S(q_1), P_S(q_2)) \leq d_g(q_1, q_2)$  for any  $q_i \in M$ ,  $i = 1, 2$ .

*Proof* (i) $\Rightarrow$ (ii) This part is contained in Udriște [291]; for completeness we give the proof. First, on account of Theorem 15.6,  $S$  is a Chebyshev set. Fix  $q_1, q_2 \in M$  and let  $\gamma_i : [-1, 1] \rightarrow M$  be the unique geodesic with constant speed from  $\gamma_i(-1) = q_i$  to  $\gamma_i(1) = s_i = P_S(q_i)$ ,  $i = 1, 2$ . Define the variation  $\Sigma : [0, 1] \times [-1, 1] \rightarrow M$  by  $\Sigma(t, u) = \exp_{\gamma_1(u)}(t \exp_{\gamma_1(u)}^{-1}(\gamma_2(u)))$ . Note that

$$\begin{aligned} L_g^2(\Sigma(\cdot, u)) &= \left( \int_0^1 \sqrt{g\left(\frac{\partial \Sigma}{\partial t}(t, u), \frac{\partial \Sigma}{\partial t}(t, u)\right)} dt \right)^2 \\ &= \int_0^1 g\left(\frac{\partial \Sigma}{\partial t}(t, u), \frac{\partial \Sigma}{\partial t}(t, u)\right) dt. \end{aligned}$$

Throughout the first variational formula, we obtain that

$$(L_g^2)'(\Sigma(\cdot, 1)) = 2 \left[ g(\gamma_2'(1), \frac{\partial \Sigma}{\partial t}(1, 1)) - g(\gamma_1'(1), \frac{\partial \Sigma}{\partial t}(0, 1)) \right].$$

Since  $S$  is geodesic convex, then the geodesic  $\Sigma(\cdot, 1)$  belongs to  $S$ . Applying Theorem 15.4, we obtain that

$$g(-\gamma_1'(1), \frac{\partial \Sigma}{\partial t}(0, 1)) \leq 0 \quad \text{and} \quad g(-\gamma_2'(1), -\frac{\partial \Sigma}{\partial t}(1, 1)) \leq 0.$$

Consequently,  $(L_g^2)'(\Sigma(\cdot, 1)) \leq 0$ . Now, using the second variational formula, and taking into account that  $\gamma_1, \gamma_2$  are geodesics, we deduce that

$$(L_g^2)''(\Sigma(\cdot, u)) = 2 \int_0^1 \left[ g\left(\frac{D}{\partial t} \frac{\partial \Sigma}{\partial t}, \frac{D}{\partial t} \frac{\partial \Sigma}{\partial t}\right) - R\left(\frac{\partial \Sigma}{\partial t}, \frac{\partial \Sigma}{\partial u}, \frac{\partial \Sigma}{\partial t}, \frac{\partial \Sigma}{\partial u}\right) \right] dt,$$

where  $R$  is the Riemannian curvature tensor. Since the sectional curvature is non-positive, we obtain  $(L_g^2)''(\Sigma(\cdot, u)) \geq 0$ . Now, the Taylor expansion yields  $L_g(\Sigma(\cdot, 1)) \leq L_g(\Sigma(\cdot, -1))$ , i.e.,  $d_g(s_1, s_2) \leq d_g(q_1, q_2)$ .

(ii) $\Rightarrow$ (i) Assume that  $S$  is not geodesic convex, i.e., there exists two distinct points  $s_1, s_2 \in S$  such that the unique geodesic with constant speed  $\gamma : [0, 1] \rightarrow M$ , with  $\gamma(0) = s_1$  and  $\gamma(1) = s_2$  has the property that  $\gamma(t) \notin S$  for every  $t \in (0, 1)$ . Since  $S$  is proximal, cf. Theorem 15.1, we fix an element  $\tilde{s} \in P_S(\gamma(1/2))$ .

We claim that either  $d_g(s_1, \tilde{s})$  or  $d_g(\tilde{s}, s_2)$  is strictly greater than  $d_g(s_1, s_2)/2$ . If  $\tilde{s} = s_1$  or  $\tilde{s} = s_2$ , the claim is true. Now, if  $s_1 \neq \tilde{s} \neq s_2$ , we assume that  $\max\{d_g(s_1, \tilde{s}), d_g(\tilde{s}, s_2)\} \leq d_g(s_1, s_2)/2$ . Therefore,  $d_g(s_1, s_2) \leq d_g(s_1, \tilde{s}) + d_g(\tilde{s}, s_2) \leq d_g(s_1, s_2)$ , which implies that  $\tilde{s} \in S$  belongs to the geodesic  $\gamma$ , contradicting our initial assumption. Thus, the claim is true. Suppose that  $d_g(s_1, \tilde{s}) > d_g(s_1, s_2)/2$ ; the other case works similarly. Since  $P_S(s_1) = s_1$  and  $d_g(s_1, s_2)/2 = d_g(s_1, \gamma(1/2))$ , relation  $d_g(s_1, \tilde{s}) > d_g(s_1, \gamma(1/2))$  contradicts (ii).  $\square$

**Remark 15.6** In spite of Theorem 15.8, no similar characterization for generic Finsler manifolds with non-positive flag curvature can be given. In the sequel, we give such a simple counterexample on Matsumoto mountain slope  $(\mathbb{R}^2, F_\alpha)$ , see (14.1). More precisely, let  $v = 10$  and  $\alpha = 60^\circ$  in  $(\mathbb{R}^2, F_\alpha)$ , and

$$S = \{(y_1, y_2) \in \mathbb{R}^2 : y_1 + y_2 = 1\}.$$

It is clear that  $S$  is closed and geodesic convex. Let  $q_1 = (0, 0)$  and  $q_2 = (0, 1)$ . Since  $q_2 \in S$ , then  $P_S^+(q_2) = q_2$ . Due to Theorem 15.6,  $\text{card}P_S^+(q_1)=1$ , and from Theorem 15.5,  $s_f \in P_S^+(q_1)$  if and only if  $g_{s_f}(s_f, e) = 0$  where  $e = (1, -1)$  is the direction vector of the straight line  $S$ . Solving the equation  $g_{s_f}(s_f, e) = 0$ , we obtain the unique solution  $s_f = (0.589888935, 0.41011065) \in S$ . Moreover,

$$\begin{aligned} d_{F_\alpha}(P_S^+(q_1), P_S^+(q_2)) &= d_{F_\alpha}(s_f, q_2) = F_\alpha(q_2 - s_f) \\ &= 0.589888935 \cdot F_\alpha(-1, 1) = 0.11923844 \\ &> 0.1 = F_\alpha(q_2 - q_1) \\ &= d_{F_\alpha}(q_1, q_2). \end{aligned}$$

Consequently,  $P_S^+$  is *not* a non-expansive map.

## 15.5 Optimal connection of two submanifolds

Let us assume that two disjoint closed sets  $M_1$  and  $M_2$  are fixed in a (not necessarily reversible) Finsler manifold  $(M, F)$ . Roughly speaking, we are interested in the number of those Finslerian geodesics which connect  $M_1$  and  $M_2$  in an optimal way, i.e., satisfying certain boundary conditions. In order to handle this problem, following Mercuri [212], Caponio, Javaloyes and Masiello [60], we first describe the structure of a special Riemann-Hilbert manifold.

Let  $(M, F)$  be a forward or backward complete Finsler manifold and

let us endow in the same time  $M$  with any complete Riemannian metric  $h$ . Let  $N$  be a smooth submanifold of  $M \times M$ . We consider the collection  $\Lambda_N(M)$  of curves  $c : [0, 1] \rightarrow M$  with  $(c(0), c(1)) \in N$  and having  $H^1$  regularity, that is,  $c$  is absolutely continuous and the integral  $\int_0^1 h(\dot{c}, \dot{c}) ds$  is finite. It is well known that  $\Lambda_N(M)$  is a Hilbert manifold modeled on any of the equivalent Hilbert spaces of  $H^1$  sections, with endpoints in  $TN$ , of the pulled back bundle  $c^*TM$ ,  $c$  being any regular curve in  $\Lambda_N(M)$ . For every  $H^1$  sections  $X$  and  $Y$  of  $c^*TM$  the scalar product is given by

$$\langle X, Y \rangle_1 = \int_0^1 h(X, Y) ds + \int_0^1 h(\nabla_c^h X, \nabla_c^h Y) ds \quad (15.5)$$

where  $\nabla_c^h$  is the usual covariant derivative along  $c$  associated to the Levi-Civita connection of the Riemannian metric  $h$ .

Let  $J : \Lambda_N(M) \rightarrow \mathbb{R}$  be the energy functional defined by

$$J(c) = \frac{1}{2} \int_0^1 F^2(c, \dot{c}) ds. \quad (15.6)$$

For further use, we denote the function  $F^2$  by  $G$ . Note that the functional  $J$  is of class  $C^{2-}$  on the space  $\Lambda_N(M)$ , i.e., it is of class  $C^1$  with locally Lipschitz differential.

**Proposition 15.1** *A curve  $\gamma \in \Lambda_N(M)$  is a constant (nonzero) speed geodesic for the Finsler manifold  $(M, F)$  satisfying the boundary condition*

$$g_{(\gamma(0), \dot{\gamma}(0))}(V, \dot{\gamma}(0)) = g_{(\gamma(1), \dot{\gamma}(1))}(W, \dot{\gamma}(1)), \quad \forall (V, W) \in T_{(\gamma(0), \dot{\gamma}(1))}N, \quad (15.7)$$

*if and only if it is a (non-constant) critical point of  $J$ .*

*Proof* Assume that  $\gamma \in \Lambda_N(M)$  is a critical point of  $J$ . One can prove in a standard way that  $\gamma$  is a smooth regular curve. Now let  $Z \in T_\gamma \Lambda_N(M)$  be a smooth vector field along  $\gamma$  and let  $\Sigma : [0, 1] \times [-\varepsilon, \varepsilon] \rightarrow M$  be a smooth regular variation of  $\gamma$  with variation vector field  $U = \frac{\partial \Sigma}{\partial u}$  having endpoints in  $TN$  and such that  $U(t, 0) = Z(t)$  for all  $t \in [0, 1]$ . Let also  $T = \frac{\partial \Sigma}{\partial t}$ . Since  $G(x, y) = g_{(x, y)}(y, y)$  for any  $(x, y) \in TM \setminus 0$ , from (13.5) we get

$$\frac{d}{du} J(\Sigma) = \frac{1}{2} \int_0^1 \frac{\partial}{\partial u} g_{(\Sigma, T)}(T, T) dt = \int_0^1 g_{(\Sigma, T)}(T, D_U T) dt. \quad (15.8)$$

Since  $\Sigma$  is smooth, we have that  $D_U T = D_T U$  both considered with the

reference vector  $T$ ; therefore, using this equality in (15.8) and evaluating at  $u = 0$  we obtain

$$dJ(\gamma)[Z] = \int_0^1 g_{(\gamma, \dot{\gamma})}(\dot{\gamma}, D_{\dot{\gamma}}Z) dt, \quad (15.9)$$

where  $D_{\dot{\gamma}}Z$  has reference vector  $\dot{\gamma}$ . Now, applying relation (13.4), we have

$$\frac{d}{dt}g_{(\gamma, \dot{\gamma})}(\dot{\gamma}, Z) = g_{(\gamma, \dot{\gamma})}(D_{\dot{\gamma}}\dot{\gamma}, Z) + g_{(\gamma, \dot{\gamma})}(\dot{\gamma}, D_{\dot{\gamma}}Z),$$

which applied to (15.9), gives that

$$\begin{aligned} 0 = dJ(\gamma)[Z] &= - \int_0^1 g_{(\gamma, \dot{\gamma})}(D_{\dot{\gamma}}\dot{\gamma}, Z) dt + g_{(\gamma(1), \dot{\gamma}(1))}(\dot{\gamma}(1), Z(1)) \\ &\quad - g_{(\gamma(0), \dot{\gamma}(0))}(\dot{\gamma}(0), Z(0)). \end{aligned} \quad (15.10)$$

Now, by choosing an endpoints vanishing vector field  $Z$  one can see that  $\gamma$  should verify the equation  $D_{\dot{\gamma}}\dot{\gamma} = 0$ , i.e.,  $\gamma$  is a constant speed geodesic. Consequently, the remaining part of the above relation gives precisely the boundary conditions (15.7).

For the converse, we observe that if  $\gamma$  is a constant non-zero speed geodesic satisfying the boundary conditions (15.7) then (15.10) holds and hence  $\gamma$  is a critical point of  $J$ .  $\square$

**Remark 15.7** Two particular cases is presented concerning the form of the submanifold  $N$ .

(i) Let  $N = \Delta$  be the diagonal in  $M \times M$ . By using the Euler theorem for homogeneous functions, we know that  $\partial_y G(x, y) = 2g_{(x, y)}(\cdot, y)$  for any  $(x, y) \in TM$ . Hence, from  $\gamma(0) = \gamma(1)$  and (15.7) we clearly have

$$\partial_y G(\gamma(0), \dot{\gamma}(0)) = \partial_y G(\gamma(0), \dot{\gamma}(1)).$$

Since the map  $y \mapsto \partial_y G(x, y)$  is an injective map, we necessarily have that  $\dot{\gamma}(0) = \dot{\gamma}(1)$ , i.e., the curve  $\gamma$  in Proposition 15.1 is a closed geodesic. See Mercuri [212].

(ii) Let  $M_1$  and  $M_2$  be two submanifolds of  $M$  and  $N = M_1 \times M_2$ . In (15.7) put  $W = 0$ . Then, for any  $V \in T_{\gamma(0)}M_1$  we get  $g_{(\gamma(0), \dot{\gamma}(0))}(V, \dot{\gamma}(0)) = 0$ . Analogously, taking  $V = 0$ , we have  $g_{(\gamma(1), \dot{\gamma}(1))}(W, \dot{\gamma}(0)) = 0$  for any  $W \in T_{\gamma(1)}M_2$ . When  $(M, F)$  is a Riemannian manifold, these conditions are actually the well-known perpendicularity conditions to  $M_1$  and  $M_2$ , respectively, see Grove [135].

Based on the papers of Caponio-Javaloyes-Masiello [60], Kozma-Krisztály-Varga [162], and Mercuri [212], we sketch the proof of the fact that

$J$  satisfies the (PS)-condition under natural assumptions. This result allows us to give several multiplicity results concerning the number of geodesics joining two different submanifolds in a Finsler manifold.

**Proposition 15.2** *Let  $(M, F)$  be forward (resp. backward) complete and  $N$  be a closed submanifold on  $M \times M$  such that the first projection (resp. the second projection) of  $N$  to  $M$  is compact. Then  $J$  satisfies the (PS)-condition on  $\Lambda_N(M)$ .*

*Proof* (Sketch) We sketch the proof in the forward complete case, the backward being similar.

*Step 1. Differentiable structure on  $\Lambda_N(M)$ .* The manifold  $\Lambda_N(M)$  is a closed submanifold of the complete Hilbert manifold  $\Lambda(M)$ ; the last one is the set of all the  $H^1$  curves in  $M$  parameterized on  $[0, 1]$  with scalar product from (15.5). The differentiable manifold structure on  $\Lambda(M)$  is given by the charts  $\{(O_\omega, \exp_\omega^{-1})\}_{\omega \in C^\infty(M)}$ , where  $\exp_\omega^{-1}$  is the inverse of the map  $\exp_\omega(\xi) = \exp_{\omega(t)} \xi(t)$ , for all  $\xi \in H^1(O_\omega)$ , being  $O_\omega$  a neighborhood of the zero section in  $\omega^*TM$ .

*Step 2. Uniform convergence of a (PS)-sequence.* We consider the sequence  $\{c_n\}_{n \in \mathbb{N}}$  contained in  $\Lambda_N(M)$  which verifies the assumptions from the (PS)-condition. In particular,  $\{J(c_n)\}_n$  is bounded. We shall prove that the sequence  $\{c_n\}_n$  converges uniformly. To do this, we fix a point  $\bar{p} \in \text{pr}_1(N)$ , where  $\text{pr}_1$  is the first projection on  $M \times M$ . Then

$$\begin{aligned} d_F(\bar{p}, c_n(s)) &\leq d_F(\bar{p}, c_n(0)) + d_F(c_n(0), c_n(s)) \\ &\leq d_F(\bar{p}, c_n(0)) + \int_0^1 F(c_n, \dot{c}_n) ds, \end{aligned}$$

for all  $s \in [0, 1]$ ,  $n \in \mathbb{N}$ . Since  $\text{pr}_1(N)$  is compact, there exists a constant  $K$  such that  $d_F(\bar{p}, c_n(0)) \leq K$ . By the Hölder inequality, for every  $s \in [0, 1]$  we have

$$d_F(\bar{p}, c_n(s)) \leq K + \left( \int_0^1 G(c_n, \dot{c}_n) ds \right)^{\frac{1}{2}} = K + \sqrt{J(c_n)} \leq K_1.$$

Consequently, the set  $S = \{c_n(s) : s \in [0, 1], n \in \mathbb{N}\}$  is a forward bounded set; thus, Hopf-Rinow theorem, see Theorem 13.1, shows that there exists a compact subset  $C$  of  $M$  which contains  $S$ . Hence there exist  $k_1, k_2 > 0$  such that

$$k_1|v|^2 \leq G(x, v) \leq k_2|v|^2, \quad \forall x \in C, v \in T_x M.$$

Here, we denoted by  $|\cdot|$  the norm associated to the metric  $h$ . Let  $d_h$

be the usual metric function associated to the Riemannian metric  $h$ . By using the last and Hölder's inequalities, we get

$$\begin{aligned} d_h(c_n(s_1), c_n(s_2)) &\leq \int_{s_1}^{s_2} |\dot{c}_n| ds \leq \sqrt{s_2 - s_1} \left( \int_0^1 |\dot{c}_n|^2 ds \right)^{\frac{1}{2}} \\ &\leq \frac{1}{k_1} \sqrt{s_2 - s_1} \left( \int_0^1 G(c_n, \dot{c}_n) ds \right)^{\frac{1}{2}} \leq K_2 \sqrt{s_2 - s_1}, \end{aligned}$$

with  $s_1 < s_2$  in  $[0, 1]$  and  $K_2 > 0$ , so that  $\{c_n(t)\}$  is relatively compact for every  $t \in [0, 1]$  and uniformly Hölder. Thus we can use the symmetric distance  $d_h$  and the Ascoli-Arzelà theorem to obtain a subsequence, that will be denoted again by  $\{c_n\}$ , converging uniformly to a curve  $\bar{c}$  of class  $C^0$  parameterized in  $[0, 1]$  and having endpoints in  $N$ .

*Step 3. Reduction of the strong convergence of  $\{c_n\}_n$  to an appropriate convergence in  $H^1$ -topology.* For any  $\eta > 0$  small enough the set  $\mathcal{C} = \{\exp_{\bar{c}(s)} v : s \in [0, 1]; v \in \bar{B}_{\bar{c}(s)}(\eta)\}$  is compact in  $M$ . Let  $\text{inj}(p)$  be the injectivity radius of  $p$  in  $(M, h)$  and  $\rho = \inf\{\text{inj}(p) : p \in \mathcal{C}\}$ . Since the injectivity radius is continuous, we have that  $\rho > 0$ ; therefore, there exists a curve  $\omega$  of class  $C^\infty$  such that  $\|\bar{c} - \omega\|_\infty < \min\{\eta, \rho/2\}$ . Let  $[0, 1] \ni t \rightarrow \mathbf{E}(t) = (E_1(t), \dots, E_m(t))$  be a parallel orthonormal frame along  $\omega$  with  $m = \dim M$ , and let  $P_t : \mathbb{R}^m \rightarrow T_{\omega(t)}M$  defined as  $P_t(v_1, \dots, v_m) = v_1 E_1(t) + \dots + v_m E_m(t)$ . Consider the Euclidean open ball of radius  $\rho$ , denoted in the sequel by  $U$ , and the map  $\varphi(t, v) = \exp_{\omega(t)} P_t(v)$ . Since  $\rho \leq \text{inj}(\omega(t))$ , the map  $\varphi_t : U \rightarrow M$ , defined as  $\varphi_t(v) = \varphi(t, v)$ , is locally invertible and injective with invertible differential  $d\varphi_t(v)$  for every  $t \in [0, 1]$  and  $v \in U$ . By taking a smaller open set in  $U$  that contains the closed ball of radius  $\rho/2$  and contained in the closed ball of radius  $2\rho/3$ , we can assume that all the continuous functions involved in the rest of the proof are uniformly bounded in  $[0, 1] \times U$  or in  $\bigcup_{t \in [0, 1]} \{t\} \times \varphi(\{t\} \times U)$ , as for example the norms of  $d\varphi(t, v)$  and  $d\theta(t, x)$ , where  $\theta(t, x) = \varphi_t^{-1}(x)$ . Let  $\mathcal{O}_\omega$  be a neighborhood of  $\omega$  in  $H^1([0, 1], M)$  such that the map  $\varphi_*^{-1} : \mathcal{O}_\omega \rightarrow H^1([0, 1], U)$ , defined as  $\varphi_*^{-1}(x)(t) = \varphi_t^{-1}(x(t))$  is the map of coordinate system centered at  $\omega$ . Observe that the inverse of  $\varphi_*^{-1}$  is the map  $\varphi_*$ , defined by  $\varphi_*(\xi)(t) = \varphi(t, \xi(t))$ . If  $n$  is large enough,  $c_n \in \varphi_*(H^1([0, 1], U))$ , so that we call  $\xi_n = \varphi_*^{-1}(c_n)$ . Consequently, taking into account the above reduction argument inspired by Abbondandolo-Figalli [1, Appendix A.1], the strong convergence of  $\{c_n\}_n$  is equivalent with the strong convergence of  $\{\xi_n\}_n$  in  $H^1([0, 1], U)$ .

*Step 4. Splitting argument.* Taking the vector space of dimension  $2m$

defined by  $V = \{\zeta \in C^\infty([0, 1], \mathbb{R}^m) : \zeta'' - \zeta = 0\}$ , we have the following orthogonal splitting

$$H^1([0, 1], \mathbb{R}^m) = H_0^1([0, 1], \mathbb{R}^m) \oplus V.$$

Therefore, if  $n \in \mathbb{N}$  is large enough there exist  $\xi_n^0 \in H_0^1([0, 1], U)$  and  $\zeta_n \in V$  such that  $\xi_n = \xi_n^0 + \zeta_n$ . Note that  $\{c_n\}$  is a sequence which verifies the assumptions of the (PS)-condition. Moreover, the norm of  $d\varphi_*$  is bounded in  $H^1([0, 1], U)$ ,  $\{\zeta_n\}_n$  is a converging sequence in the  $C^1$  norm (which follows from the  $C^0$  convergence of  $\{\xi_n\}_n$  and the smooth dependence of the solutions of the differential equation defining  $V$  from boundary data) and  $\{\xi_n\}_n$  is a bounded sequence in  $H^1([0, 1], U)$  ( $\xi_n^0$  is a bounded sequence in  $H_0^1([0, 1], U)$ ). Due to the above facts and considering  $J$  defined on the manifold  $\Lambda(M)$ , we have

$$\begin{aligned} & d(J \circ \varphi_*)(\xi_n)[\xi_n - \xi_k] \\ &= d(J \circ \varphi_*)(\xi_n)[\xi_n^0 - \xi_k^0] + d(J \circ \varphi_*)(\xi_n)[\zeta_n - \zeta_k] \\ &= dJ(c_n)[d\varphi_*(\xi_n)[\xi_n^0 - \xi_k^0]] + d(J \circ \varphi_*)(\xi_n)[\zeta_n - \zeta_k] \rightarrow 0. \end{aligned} \quad (15.11)$$

*Step 5. Boundedness of  $\{\xi_n\}_n$  in  $H^1([0, 1], U)$ .* It is enough to observe that

$$\begin{aligned} & \int_0^1 |\dot{\xi}_n|^2 ds = \int_0^1 |d\varphi(s, c_n)[(1, \dot{c}_n)]|^2 ds \\ & \leq K_3 \int_0^1 (1 + h(\dot{c}_n, \dot{c}_n)) ds \leq K_3 + K_4 J(c_n) < K_5 < +\infty, \end{aligned} \quad (15.12)$$

where  $\varphi(s, x) = \varphi_s^{-1}(x)$ , for every  $s \in [0, 1]$  and  $x \in \varphi_s(U)$ , and  $K_3, K_4, K_5$  are positive constants.

*Step 6. (Final) The sequence  $\{\xi_n\}_n$  is fundamental in  $H^1([0, 1], U)$ .* By using similar estimates as (15.11) and (15.12), a technical calculation shows that

$$\int_0^1 |\dot{\xi}_n - \dot{\xi}_k|^2 ds \rightarrow 0$$

as  $n, k \rightarrow \infty$ . This concludes the proof.  $\square$

Now, we are in the position to establish an existence and some multiplicity results concerning the number of geodesics joining two submanifolds in a Finsler manifold.

**Theorem 15.9** *Let  $(M, F)$  be a forward or backward complete Finsler manifold and let  $M_1$  and  $M_2$  be two closed submanifolds of  $M$  such that  $M_1$  or  $M_2$  is compact. Then in any homotopy class of curves from  $M_1$  to  $M_2$  there exists a geodesic with energy smaller than that of any other curve in this class satisfying (15.7). Furthermore, there are at least  $\text{cat}\Lambda_{M_1 \times M_2}(M)$  geodesics joining  $M_1$  and  $M_2$  with the property (15.7).*

*Proof* Since  $\Lambda_{M_1 \times M_2}(M)$  is a complete Hilbert-Riemann manifold and the energy functional satisfies the (PS)-condition, see Proposition 15.2, it follows that the energy integral attains its infimum on any component of  $\Lambda_{M_1 \times M_2}(M)$  and its lower bound, see Theorem 1.4. The infimum points are critical points of  $J$ , while any critical point  $c$  for  $J$  is a geodesic with the property (15.7), see Proposition 15.1. Therefore, the first part is proved. Now, applying Theorem 1.11, we obtain our last claim.  $\square$

**Theorem 15.10** *Let  $(M, F)$  be a compact, connected and simply connected Finsler manifold, and let  $M_1$  and  $M_2$  be two closed disjoint submanifolds of  $M$  with  $M_1$  contractible. Then there are infinitely many geodesics joining  $M_1$  and  $M_2$  with the property (15.7).*

*Proof* Due to Theorem C.3, we have that  $\text{cuplong}\Lambda_{M_1 \times M_2}(M) = \infty$ . Using the inequality  $\text{cat}\Lambda_{M_1 \times M_2}(M) \geq 1 + \text{cuplong}\Lambda_{M_1 \times M_2}(M)$ , from Theorem 15.9 the statement follows.  $\square$

**Theorem 15.11** *Let  $(M, F)$  be a complete, non-contractible Finsler manifold, and let  $M_1$  and  $M_2$  be two closed, disjoint and contractible submanifolds of  $M$  such that  $M_1$  or  $M_2$  is compact. Then there are infinitely many geodesics joining  $M_1$  and  $M_2$  with the property (15.7).*

*Proof* Since  $M_1 \times M_2$  is a submanifold of the product manifold  $M \times M$ , the inclusion  $\Lambda_{M_1 \times M_2}M \hookrightarrow C_{M_1 \times M_2}^0(M) = \{\sigma \in C^0([0, 1], M) : \sigma(0) \in M_1, \sigma(1) \in M_2\}$  is a homotopy equivalence, see Grove [135, Theorem 1.3]. Since  $M_1$  and  $M_2$  are contractible subsets of  $M$ , the sets  $C_{M_1 \times M_2}^0(M)$  and  $M_1 \times M_2 \times \Omega(M)$  are homotopically equivalent, see Fadell-Husseini [111, Proposition 3.2]. Since  $M$  is non-contractible, we have  $\text{cat}\Omega(M) = \infty$ , see Theorem C.1. Therefore, we have that  $\text{cat}\Lambda_{M_1 \times M_2}(M) = \infty$  and we apply again Theorem 15.9 to obtain the desired conclusion.  $\square$

## 15.6 Remarks and perspectives

*A. Remarks.* In this chapter we discussed mainly the best approximation

problem on a non-reversible framework, motivated by some real world applications. Besides of existence and characterization results of the projection, we pointed out some qualitative differences between well-known results of best approximations on normed vector spaces and those in the Finsler context by constructing (counter)examples on the Finslerian-Poincaré disc and on the Matsumoto's mountain slope. The results in Sections 15.2-15.4 are contained in the paper of Kristály [172]. In Section 15.5 we studied the number of Finslerian geodesics connecting 'perpendicularly' two submanifolds in a given Finsler manifold. These results are based on the papers of Caponio-Javaloyes-Masiello [60] and Kozma-Kristály-Varga [162].

*B. Perspectives.* Some important questions arise concerning the best approximation problems on Finsler manifolds. We formulate only a few of them:

- (i). The Moskovitz-Dines type characterization of projections has been proved for Hadamard-type Riemannian manifolds (Theorem 15.4) and Minkowski spaces (Theorem 15.5). We feel quite certain we face a rigidity result: if  $(MD_1^+) \Leftrightarrow (MD_2^+)$  (or,  $(MD_1^-) \Leftrightarrow (MD_2^-)$ ) holds on a Finsler manifold  $(M, F)$  with non-positive flag curvature, then  $(M, F)$  is either a Riemannian manifold or a Minkowski space.
- (ii). We were able to prove the uniqueness result (Theorem 15.6) only on Berwald spaces. Try to extend Theorem 15.6 to any (not necessarily reversible) Finsler manifold with non-positive flag curvature; see also Remark 15.4.
- (iii). Determine those classes of Finsler manifolds with non-positive flag curvature whose (forward and backward) projections are non-expansive for every nonempty, closed, geodesic convex set. We believe the only class fulfilling this property is the class of Hadamard-type Riemannian manifolds. There are three supporting reasons for this fact: (1) non-expansiveness of the projection map holds on Hadamard-type Riemannian manifolds, see Theorem 15.8; (2) in Remark 15.6 we gave a counterexample for non-expansivity of the projection map on a slightly general case than Riemannian manifolds; (3) the non-expansiveness of the projection map holds on a normed vector space if and only if the norm comes from an inner product, see Phelps [243].

# 16

## A Variational Approach to Nash Equilibria

I did have strange ideas during certain periods of time.

---

John F. Nash (b. 1928)

### 16.1 Introduction

Nash equilibrium plays a central role in game theory; it is a concept strategy of a game involving  $n$ -players ( $n \geq 2$ ) in which every player know the equilibrium strategies of the other players, and changing his/her own strategy alone, a player has nothing to gain.

Let  $K_1, \dots, K_n$  ( $n \geq 2$ ) be the nonempty sets of strategies of the players and  $f_i : K_1 \times \dots \times K_n \rightarrow \mathbb{R}$  ( $i \in \{1, \dots, n\}$ ) be the payoff functions. A point  $\mathbf{p} = (p_1, \dots, p_n) \in \mathbf{K}$  is a *Nash equilibrium point* for  $(\mathbf{f}, \mathbf{K})$  if

$$f_i(\mathbf{p}; q_i) \geq f_i(\mathbf{p}) \quad \text{for all } q_i \in K_i, i \in \{1, \dots, n\}.$$

Here and in the sequel, the following notations are used:

$$\begin{aligned} \mathbf{K} &= \prod_{i=1}^n K_i; \quad \mathbf{p} = (p_1, \dots, p_n); \quad (\mathbf{f}, \mathbf{K}) = (f_1, \dots, f_n; K_1, \dots, K_n); \\ (\mathbf{p}; q_i) &= (p_1, \dots, q_i, \dots, p_n). \end{aligned}$$

The most well-known existence result is due to Nash (see [228], [227]) which works for compact and convex subsets  $K_i$  of Hausdorff topological vector spaces, and the continuous payoff functions  $f_i$  are (quasi)convex in the  $i^{\text{th}}$ -variable,  $i \in \{1, \dots, n\}$ . A natural question that arises at this point is:

*How is it possible to guarantee the existence of Nash equilibrium points for a*

family of payoff functions (perhaps set-valued) without any convexity, or even more, when their domains are not convex in the usual sense?

In this chapter we focus our attention to the existence and location of Nash equilibrium points in various context. First, in Section 16.2 we give several formulations of Nash equilibria via variational inequalities of Stampacchia-, and Minty-type following the paper of Cavazzuti-Pappalardo-Passacantando [63], while in Section 16.3 we are dealing with Nash equilibrium points for set-valued maps on vector spaces. In Section 16.4, we treat a case when the domain of the strategy sets are not necessarily convex in the usual sense by applying elements from Riemannian geometry.

## 16.2 Nash equilibria and variational inequalities

We first recall the well-known result of Nash (see [228], [227]), concerning the existence of at least one Nash equilibrium point for a system  $(\mathbf{f}, \mathbf{K})$ .

**Theorem 16.1** *Let  $K_1, \dots, K_n$  be nonempty compact convex subsets of Hausdorff topological vector spaces, and let  $f_i : \mathbf{K} \rightarrow \mathbb{R}$  ( $i \in \{1, \dots, n\}$ ) be continuous functions such that  $q_i \in K_i \mapsto f_i(\mathbf{p}; q_i)$  is quasiconvex for all fixed  $p_j \in K_j$  ( $j \neq i$ ). Then there exists a Nash equilibrium point for  $(\mathbf{f}, \mathbf{K})$ .*

In the sequel, assume that the sets of strategies of the players, i.e.,  $K_1, \dots, K_n$  ( $n \geq 2$ ), are closed convex subsets of a common vector space  $X$ . Moreover, we also assume that the payoff functions  $f_i$  are continuous on  $\mathbf{K}$  and each  $f_i$  is of class  $C^1$  in its  $i^{\text{th}}$  variable on an open set  $D_i \supseteq K_i$ . Let  $\mathbf{p} \in \mathbf{K}$ ; we introduce the notations

$$\nabla \mathbf{f}(\mathbf{p}) = \left( \frac{\partial f_1}{\partial p_1}(\mathbf{p}), \dots, \frac{\partial f_n}{\partial p_n}(\mathbf{p}) \right) \quad \text{and} \quad \langle \nabla \mathbf{f}(\mathbf{p}), \mathbf{q} \rangle = \sum_{i=1}^n \left\langle \frac{\partial f_i}{\partial p_i}(\mathbf{p}), q_i \right\rangle,$$

where  $\langle \cdot, \cdot \rangle$  inside the sum denotes the duality pair between  $X$  and  $X^*$ .

A point  $\mathbf{p} = (p_1, \dots, p_n) \in \mathbf{K}$  is a *Nash-Stampacchia point* for  $(\mathbf{f}, \mathbf{K})$  if

$$\langle \nabla \mathbf{f}(\mathbf{p}), \mathbf{q} - \mathbf{p} \rangle \geq 0 \quad \text{for all } \mathbf{q} \in \mathbf{K}.$$

In a similar way, a point  $\mathbf{p} = (p_1, \dots, p_n) \in \mathbf{K}$  is a *Nash-Minty point* for  $(\mathbf{f}, \mathbf{K})$  if

$$\langle \nabla \mathbf{f}(\mathbf{q}), \mathbf{q} - \mathbf{p} \rangle \geq 0 \quad \text{for all } \mathbf{q} \in \mathbf{K}.$$

**Theorem 16.2** (a) Every Nash equilibrium point for  $(\mathbf{f}, \mathbf{K})$  is a Nash-Stampacchia point for  $(\mathbf{f}, \mathbf{K})$ .

(b) Every Nash-Minty equilibrium point for  $(\mathbf{f}, \mathbf{K})$  is a Nash equilibrium point for  $(\mathbf{f}, \mathbf{K})$ .

(c) If each  $f_i$  is convex in its  $i^{\text{th}}$  variable, then the converses also hold in (a) and (b).

*Proof* The proof easily follows from Lemmas 13.1 and 13.2.  $\square$

**Theorem 16.3** Assume that  $\nabla \mathbf{f}$  is a strictly monotone operator. Then, the following assertions hold true:

(a)  $(\mathbf{f}, \mathbf{K})$  has at most one Nash-Stampacchia point.

(b)  $(\mathbf{f}, \mathbf{K})$  has at most one Nash equilibrium point.

*Proof* (a) Assume that  $\mathbf{p}_1 \neq \mathbf{p}_2 \in \mathbf{K}$  are Nash-Stampacchia points for  $(\mathbf{f}, \mathbf{K})$ . In particular, we clearly have that

$$\langle \nabla \mathbf{f}(\mathbf{p}_1), \mathbf{p}_2 - \mathbf{p}_1 \rangle \geq 0,$$

$$\langle \nabla \mathbf{f}(\mathbf{p}_2), \mathbf{p}_1 - \mathbf{p}_2 \rangle \geq 0.$$

Therefore, by the strict monotonicity assumptions we have

$$0 \leq \langle \nabla \mathbf{f}(\mathbf{p}_1) - \nabla \mathbf{f}(\mathbf{p}_2), \mathbf{p}_2 - \mathbf{p}_1 \rangle < 0,$$

a contradiction.

(b) It follows from Theorem 16.2 (a).  $\square$

Usually, the set of Nash-Stampacchia points and the set of Nash equilibrium points are different. A simple example which support this statement is presented in what follows, see also Kassay-Kolumbán-Páles [155].

**Example 16.1** Let  $K_1 = K_2 = [-1, 1]$ ,  $D_1 = D_2 = \mathbb{R}$ , and  $f_1, f_2 : D_1 \times D_2 \rightarrow \mathbb{R}$  defined by

$$f_1(x_1, x_2) = x_1^2 x_2 + x_1, \quad f_2(x_1, x_2) = -x_1^2 x_2^2 + x_2.$$

One can prove that the set of Nash-Stampacchia points for  $(\mathbf{f}, \mathbf{K})$  is

$$\{(1/2, -1), (-1, 1/2), (-1, -1), (1, -1)\}.$$

However, the unique Nash equilibrium point for  $(\mathbf{f}, \mathbf{K})$  is  $(-1, -1)$ .

In the rest of this section we assume that  $X$  is a prehilbert space. Consequently, the inequality from the definition of Nash-Stampacchia point for  $(\mathbf{f}, \mathbf{K})$  can be written equivalently as

$$\langle (\mathbf{p} - \alpha \nabla \mathbf{f}(\mathbf{p})) - \mathbf{p}, \mathbf{q} - \mathbf{p} \rangle \leq 0 \quad \text{for all } \mathbf{q} \in \mathbf{K}, \alpha > 0.$$

Furthermore, in terms of projections, the Moskovitz-Dines-type relation gives that,

$$\mathbf{p} \in P_{\mathbf{K}}(\mathbf{p} - \alpha \nabla \mathbf{f}(\mathbf{p})) \text{ for all } \alpha > 0,$$

see (15.2). Note that since  $\mathbf{K}$  is convex, it is also a Chebyshev set, i.e., the above relation has the form

$$0 = P_{\mathbf{K}}(\mathbf{p} - \alpha \nabla \mathbf{f}(\mathbf{p})) - \mathbf{p} \text{ for all } \alpha > 0.$$

The last relation motivates the study of the following continuous dynamical system

$$\dot{\mathbf{x}}(t) = P_{\mathbf{K}}(\mathbf{x}(t) - \alpha \nabla \mathbf{f}(\mathbf{x}(t))) - \mathbf{x}(t), \quad (16.1)$$

where  $\alpha > 0$  is fixed. Following the paper of Cavazzuti-Pappalardo-Passacantando [63], we prove

**Theorem 16.4** *Assume that  $\nabla \mathbf{f}(\mathbf{p}) = 0$ . If  $\nabla \mathbf{f}$  is strongly monotone on  $\mathbf{K}$  with constant  $\eta > 0$  and Lipschitz continuous on  $\mathbf{K}$  with constant  $L > 0$ , then there exists  $\alpha_0 > 0$  such that for every  $\alpha \in (0, \alpha_0)$  there exists a constant  $C > 0$  such that for each solution  $\mathbf{x}(t)$  of (16.1), with  $\mathbf{x}(0) \in \mathbf{K}$ , we have*

$$\|\mathbf{x}(t) - \mathbf{p}\| \leq \|\mathbf{x}(0) - \mathbf{p}\| \exp(-Ct), \quad \forall t \geq 0,$$

i.e.,  $\mathbf{x}(t)$  converges exponentially to  $\mathbf{p} \in \mathbf{K}$ .

*Proof* Standard ODE theory shows that there exists a unique maximal semi-flow  $\mathbf{x}$  of (16.1) on an interval  $[0, T)$ . We shall prove that  $T = +\infty$ . First of all, due to the non-expansiveness of the projection operator  $P_{\mathbf{K}}$ , for every  $t \in [0, T)$ , we have that

$$\begin{aligned} & \|P_{\mathbf{K}}(\mathbf{x}(t) - \alpha \nabla \mathbf{f}(\mathbf{x}(t))) - \mathbf{p}\|^2 = \\ & = \|P_{\mathbf{K}}(\mathbf{x}(t) - \alpha \nabla \mathbf{f}(\mathbf{x}(t))) - P_{\mathbf{K}}(\mathbf{p} - \alpha \nabla \mathbf{f}(\mathbf{p}))\|^2 \leq \\ & \leq \|\mathbf{x}(t) - \mathbf{p} - \alpha(\nabla \mathbf{f}(\mathbf{x}(t)) - \nabla \mathbf{f}(\mathbf{p}))\|^2 \\ & = \|\mathbf{x}(t) - \mathbf{p}\|^2 - 2\alpha \langle \nabla \mathbf{f}(\mathbf{x}(t)) - \nabla \mathbf{f}(\mathbf{p}), \mathbf{x}(t) - \mathbf{p} \rangle + \alpha^2 \|\nabla \mathbf{f}(\mathbf{x}(t)) - \nabla \mathbf{f}(\mathbf{p})\|^2 \\ & \leq (1 - 2\alpha\eta + \alpha^2 L^2) \|\mathbf{x}(t) - \mathbf{p}\|^2. \end{aligned}$$

Let  $\alpha_0 = 2\eta/L^2$  and  $\alpha \in (0, \alpha_0)$ . Note that the whole orbit of  $\mathbf{x}(t)$  remains in  $\mathbf{K}$ . We consider the function

$$h(t) = \frac{1}{2} \|\mathbf{x}(t) - \mathbf{p}\|^2, \quad t \in [0, T).$$

Then, we have

$$\begin{aligned} \dot{h}(t) &= \langle \mathbf{x}(t) - \mathbf{p}, P_{\mathbf{K}}(\mathbf{x}(t) - \alpha \nabla \mathbf{f}(\mathbf{x}(t))) - \mathbf{x}(t) \rangle \\ &= \langle \mathbf{x}(t) - \mathbf{p}, P_{\mathbf{K}}(\mathbf{x}(t) - \alpha \nabla \mathbf{f}(\mathbf{x}(t))) - \mathbf{p} \rangle - \|\mathbf{x}(t) - \mathbf{p}\|^2 \\ &\leq \|\mathbf{x}(t) - \mathbf{p}\| \cdot \|P_{\mathbf{K}}(\mathbf{x}(t) - \alpha \nabla \mathbf{f}(\mathbf{x}(t))) - \mathbf{p}\| - \|\mathbf{x}(t) - \mathbf{p}\|^2 \\ &\leq (\sqrt{1 - 2\alpha\eta} + \alpha^2 L^2 - 1) \|\mathbf{x}(t) - \mathbf{p}\|^2 \\ &= -2Ch(t), \end{aligned}$$

where  $C = 1 - \sqrt{1 - 2\alpha\eta} + \alpha^2 L^2} > 0$ . Consequently, we have

$$\dot{h}(t) \leq -2Ch(t),$$

with  $h(0) = \frac{1}{2} \|\mathbf{x}(0) - \mathbf{p}\|^2$ . In particular, the function  $t \mapsto h(t) \exp(2Ct)$  is non-increasing on the interval  $[0, T)$ . In particular,  $\mathbf{x}$  can be extended toward to  $T > 0$  which contradicts the maximality of the interval  $[0, T)$ . Therefore,  $T = +\infty$ . After an integration, we have that

$$\|\mathbf{x}(t) - \mathbf{p}\| \leq \|\mathbf{x}(0) - \mathbf{p}\| \exp(-Ct), \quad \forall t \geq 0,$$

which completes the proof. □

### 16.3 Nash equilibria for set-valued maps

In some cases the objective/payoff functions are not single-valued, i.e., the values of a payoff functions are not real numbers but some sets. Set-valued versions of the Nash equilibrium problem can be founded in the literature, see for example Guillerme [136] and Luo [199]. In this section we propose a new approach in the study of the existence of Nash equilibria for set-valued maps which is based on the contingent derivative.

Let  $K_1, \dots, K_n$  be nonempty subsets of a real normed space  $X$ , and let  $F_i : \mathbf{K} \rightsquigarrow \mathbb{R}$  ( $i \in \{1, \dots, n\}$ ) be set-valued maps with nonempty compact values. A point  $\mathbf{p} = (p_1, \dots, p_n) \in \mathbf{K}$  is a *Nash equilibrium point* for  $(\mathbf{F}, \mathbf{K}) = (F_1, \dots, F_n; K_1, \dots, K_n)$  if  $\mathbf{p}$  is a Nash equilibrium point for  $(\min \mathbf{F}, \mathbf{K})$ .

The following result is an easy consequence of Theorem 16.1.

**Theorem 16.5** *Let  $K_1, \dots, K_n$  be nonempty compact convex subsets of a real normed space  $X$ , and let  $F_i : \mathbf{K} \rightsquigarrow \mathbb{R}$  ( $i \in \{1, \dots, n\}$ ) be continuous set-valued maps on  $\mathbf{K}$  with nonempty compact values such that  $q_i \in K_i \rightsquigarrow F_i(\mathbf{p}; q_i)$  is convex on  $K_i$  for all fixed  $p_j \in K_j$  ( $j \neq i$ ). Then there exists a Nash equilibrium point for  $(\mathbf{F}, \mathbf{K})$ .*

*Proof* Let  $f_i = \min F_i$  ( $i \in \{1, \dots, n\}$ ). It is easy to prove that the functions  $f_i$  are continuous and convex in the  $i^{\text{th}}$  variable. It remains to apply Theorem 16.1.  $\square$

Let  $K = K_1 = K_2 = [-1, 1]$ ,  $X = \mathbb{R}$ , and  $F_1, F_2 : K \times K \rightsquigarrow \mathbb{R}$  be defined by

$$F_1(x_1, x_2) = [\max\{|x_1|, |x_2|\} - 1, 0]; \quad (16.2)$$

$$F_2(x_1, x_2) = [1 - \max\{|x_1|, |x_2|\}, 2 - x_1^2 - x_2^2]. \quad (16.3)$$

Clearly,  $F_1$  and  $F_2$  are continuous on  $K \times K$ , but  $x_2 \in K \rightsquigarrow F_2(0, x_2)$  is not convex on  $K$ . Therefore, Theorem 16.5 can not be applied. However, we will see in Example 16.2 that there are Nash equilibrium points for  $(\mathbf{F}, \mathbf{K}) = (F_1, F_2; K, K)$ . To determine these points we elaborate a method by using the contingent derivative of set-valued maps.

Let  $K_1, \dots, K_n$  be nonempty convex subsets of a real normed space  $X$ , and let  $F_i : K_1 \times \dots \times X \times \dots \times K_n \rightsquigarrow \mathbb{R}$  ( $i \in \{1, \dots, n\}$ ) be set-valued maps with compact, nonempty values. Here,  $X$  is in the  $i^{\text{th}}$  position. Let  $\mathbf{p} \in \mathbf{K}$  be a fixed element. We can define the (partial) contingent derivative of  $F_i$  in the  $i^{\text{th}}$  variable at the point  $(\mathbf{p}, \min F_i(\mathbf{p}))$ , i.e., the contingent derivative of  $F_i(p_1, \dots, p_{i-1}, \cdot, p_{i+1}, \dots, p_n)$  at the point  $(p_i, \min F_i(\mathbf{p}))$ , see Appendix, Definition E.4.

**Definition 16.1** A point  $\mathbf{p} = (p_1, \dots, p_n) \in \mathbf{K}$  is said to be a Nash contingent point for  $(\mathbf{F}, \mathbf{K})$  if

$$D_i F_i(\mathbf{p}, \min F_i(\mathbf{p}))(q_i - p_i) \subseteq \mathbb{R}_+$$

for all  $q_i \in K_i$  and all  $i \in \{1, \dots, n\}$ .

**Proposition 16.1** Let  $\mathbf{p} = (p_1, \dots, p_n) \in \mathbf{K}$  be a Nash equilibrium point for  $(\mathbf{F}, \mathbf{K})$ . If each map  $q_i \rightsquigarrow F_i(\mathbf{p}; q_i)$  ( $i \in \{1, \dots, n\}$ ) is  $K_i$ -locally Lipschitz, then  $\mathbf{p}$  is a Nash contingent point for  $(\mathbf{F}, \mathbf{K})$ .

*Proof* Since  $\mathbf{p}$  is a Nash equilibrium point for  $(\mathbf{F}, \mathbf{K})$ , we have

$$F_i(\mathbf{p}; q_i) - \min F_i(\mathbf{p}) \subseteq \mathbb{R}_+ \quad (16.4)$$

for all  $q_i \in K_i$  and all  $i \in \{1, \dots, n\}$ .

Let  $i \in \{1, \dots, n\}$ ,  $q_i \in K_i$  be fixed elements and we fix arbitrarily  $c_i \in D_i F_i(\mathbf{p}, \min F_i(\mathbf{p}))(q_i - p_i)$ . We prove that  $c_i \geq 0$ . From (E.2) we have that

$$\liminf_{t \rightarrow 0^+} \text{dist} \left( c_i, \frac{F_i(\mathbf{p}; p_i + t(q_i - p_i)) - \min F_i(\mathbf{p})}{t} \right) = 0, \quad (16.5)$$

since the map  $v_i \rightsquigarrow F_i(\mathbf{p}; v_i)$  is  $K_i$ -locally Lipschitz. Since  $K_i$  is convex, for  $t > 0$  small enough,  $y_i^t = p_i + t(q_i - p_i) \in K_i$ . Using (16.4), we have that

$$\frac{F_i(\mathbf{p}; y_i^t) - \min F_i(\mathbf{p})}{t} \subseteq \mathbb{R}_+.$$

Suppose that  $c_i < 0$ . Then

$$0 < |c_i| = \text{dist}(c_i, \mathbb{R}_+) \leq \text{dist}\left(c_i, \frac{F_i(\mathbf{p}; y_i^t) - \min F_i(\mathbf{p})}{t}\right),$$

which contradicts (16.5). □

The above result allows us to select the Nash equilibrium points among Nash contingent points. In most of the cases the study of contingent Nash points is much easier than those of the Nash equilibrium points. In the sequel we present two such examples. We will use the notation

$$[a, b] = [\min\{a, b\}, \max\{a, b\}], \text{ where } a, b \in \mathbb{R}.$$

**Example 16.2** Let us consider the set-valued maps from (16.2) and (16.3) in extended forms, i.e.  $K = K_1 = K_2 = [-1, 1]$ ,  $X = \mathbb{R}$  and let  $F_1 : X \times K \rightsquigarrow \mathbb{R}$ ,  $F_2 : K \times X \rightsquigarrow \mathbb{R}$  be defined by

$$F_1(x_1, x_2) = [\max\{|x_1|, |x_2|\} - 1, 0];$$

$$F_2(x_1, x_2) = [1 - \max\{|x_1|, |x_2|\}, 2 - x_1^2 - x_2^2].$$

It is easy to verify that  $u \in K \rightsquigarrow F_1(u, x_2)$  and  $v \in K \rightsquigarrow F_2(x_1, v)$  are  $K$ -locally Lipschitz maps ( $x_1, x_2 \in K$  being fixed points). We find those points  $(x_1, x_2) \in K \times K$  which fulfill

$$D_1 F_1((x_1, x_2), \min F_1(x_1, x_2))(v_1 - x_1) \subseteq \mathbb{R}_+ \text{ for all } v_1 \in K; \tag{16.6}$$

$$D_2 F_2((x_1, x_2), \min F_2(x_1, x_2))(v_2 - x_2) \subseteq \mathbb{R}_+ \text{ for all } v_2 \in K. \tag{16.7}$$

Using the geometric meaning of the contingent derivative (see relation (E.1)), after an elementary discussion and computation, the points which satisfy the inclusions (16.6) and (16.7) respectively, are

$$CNP_1 = \{(x_1, x_2) \in K \times K : |x_1| \leq |x_2|\};$$

$$CNP_2 = \{(x_1, x_2) \in K \times K : |x_1| > |x_2|\} \cup \{(x_1, x_2) \in K \times K : |x_2| = 1\}.$$

Therefore, the contingent Nash points for  $(\mathbf{F}, \mathbf{K}) = (F_1, F_2; K, K)$  are

$$CNP = CNP_1 \cap CNP_2 = \{(x_1, x_2) \in K \times K : |x_2| = 1\}.$$

A direct computation shows that the set of Nash equilibrium points for  $(\mathbf{F}, \mathbf{K})$  coincides with the set  $CNP$ .

**Example 16.3** Let  $K = [-1, 1]$ , and let  $F_1 : \mathbb{R} \times K \rightsquigarrow \mathbb{R}$  and  $F_2 : K \times \mathbb{R} \rightsquigarrow \mathbb{R}$  be defined by

$$F_1(x_1, x_2) = \begin{cases} [2, |x_1 - 1|], & \text{if } x_2 \geq 0, \\ [2, |x_1 + 1|], & \text{if } x_2 < 0, \end{cases}$$

and

$$F_2(x_1, x_2) = \begin{cases} [2, |x_2 + 1|], & \text{if } x_1 \geq 0, \\ [2, |x_2 - 1|], & \text{if } x_1 < 0. \end{cases}$$

The points which satisfy the corresponding inclusions from (16.6) and (16.7) respectively, are

$$CNP_1 = \{(1, x_2) : x_2 \in [0, 1]\} \cup \{(-1, x_2) : x_2 \in [-1, 0)\};$$

$$CNP_2 = \{(x_1, -1) : x_1 \in [0, 1]\} \cup \{(x_1, 1) : x_1 \in [-1, 0)\}.$$

The maps  $u \in K \rightsquigarrow F_1(u, x_2)$  and  $v \in K \rightsquigarrow F_2(x_1, v)$  are  $K$ -locally Lipschitz ( $x_1, x_2 \in K$  being fixed points), however the set of Nash contingent points for  $(\mathbf{F}, \mathbf{K}) = (F_1, F_2; K, K)$  is  $CNP = CNP_1 \cap CNP_2$ , which is empty. Hence the set of Nash equilibrium points is empty too, due to Proposition 16.1.

Note that the set-valued maps from Examples 16.2 and 16.3 are not continuous on  $K \times K$ . In the sequel, we give sufficient conditions to obtain Nash contingent points, assuming some continuity hypotheses on the set-valued maps. Before to do this, we can state the converse of Proposition 16.1 by taking a convexity assumption for the corresponding set-valued maps. Let  $X$  be a normed space, and let  $K$  be a nonempty subset of  $X$ .

**Definition 16.2** The set-valued map  $F : X \rightsquigarrow \mathbb{R}$  is called  $K$ -pseudoconvex at  $(u, c) \in \text{Graph}(F)$  if

$$F(u') \subseteq c + DF(u, c)(u' - u) \text{ for all } u' \in K.$$

In the case when  $K = \text{Dom}(F)$ , the above definition reduces to [17, Definition 5.1.1]. The following result can be easily proved.

**Proposition 16.2** Let  $K_1, \dots, K_n$  be nonempty subsets of a real normed space  $X$ , and let  $F_i : K_1 \times \dots \times X \times \dots \times K_n \rightsquigarrow \mathbb{R}$  ( $i \in \{1, \dots, n\}$ ) be set-valued maps with nonempty compact values. Let  $\mathbf{p} = (p_1, \dots, p_n) \in \mathbf{K}$  be a Nash contingent point for  $(\mathbf{F}, \mathbf{K})$ . If  $q_i \rightsquigarrow F_i(\mathbf{p}; q_i)$  ( $i \in \{1, \dots, n\}$ ) is

$K_i$ -pseudo-convex at  $(p_i, \min F_i(\mathbf{p}))$ , then  $\mathbf{p}$  is a Nash equilibrium point for  $(\mathbf{F}, \mathbf{K})$ .

We give an existence theorem concerning the Nash contingent points for a system  $(\mathbf{F}, \mathbf{K})$  by using a Fan-type result.

**Theorem 16.6** [Kristály-Varga [179]] *Let  $\tilde{X}$  be a real normed space, let  $K$  be a nonempty convex compact subset of  $\tilde{X}$ , and let  $F : K \times K \rightsquigarrow \mathbb{R}$  be a set-valued map satisfying the following conditions:*

- (i) for all  $v \in K$ ,  $u \in K \rightsquigarrow F(u, v)$  is lower semicontinuous on  $K$ ;
- (ii) for all  $u \in K$ ,  $v \in K \rightsquigarrow F(u, v)$  is convex on  $K$ ;
- (iii) for all  $u \in K$ ,  $F(u, u) \subseteq \mathbb{R}_+$ .

Then there exists an element  $\bar{u} \in K$  such that

$$F(\bar{u}, v) \subseteq \mathbb{R}_+ \text{ for all } v \in K. \quad (16.8)$$

Let  $K_i$  and  $F_i$  ( $i \in \{1, \dots, n\}$ ) be as in Proposition 16.2. We denote by  $F_i|_{\mathbf{K}}$  the restriction of  $F_i$  to  $\mathbf{K} = K_1 \times \dots \times K_i \times \dots \times K_n$ .

**Definition 16.3**  $F_i|_{\mathbf{K}}$  is called  $i$ -lower semicontinuously differentiable if

$$\text{Graph}(F_i|_{\mathbf{K}}) \times X \ni (\mathbf{p}, c, h) \rightsquigarrow D_i F_i(\mathbf{p}, c)(h)$$

is lower semicontinuous.

Now, we are in the position to give an existence result concerning the Nash contingent points.

**Theorem 16.7** *Let  $K_1, \dots, K_n$  be nonempty compact convex subsets of a real normed space  $X$  and let  $F_i : K_1 \times \dots \times X \times \dots \times K_n \rightsquigarrow \mathbb{R}$  ( $i \in \{1, \dots, n\}$ ) be set-valued maps with nonempty compact values and with closed graph. Suppose that for each  $i \in \{1, \dots, n\}$  the map  $F_i$  is  $K_i$ -locally Lipschitz in the  $i^{\text{th}}$  variable,  $F_i|_{\mathbf{K}}$  is continuous on  $\mathbf{K}$  and  $i$ -lower semicontinuously differentiable. Then there exists a Nash contingent point for  $(\mathbf{F}, \mathbf{K})$ .*

*Proof* Let  $\tilde{X} = X \times \dots \times X$ . Clearly,  $\mathbf{K}$  is a compact, convex subset of  $\tilde{X}$ . We define the map  $G : \mathbf{K} \times \mathbf{K} \rightsquigarrow \mathbb{R}$  by

$$G(\mathbf{p}, \mathbf{q}) = \sum_{i=1}^n D_i F_i(\mathbf{p}, \min F_i(\mathbf{p}))(q_i - p_i),$$

where  $\mathbf{p} = (p_1, \dots, p_n)$  and  $\mathbf{q} = (q_1, \dots, q_n)$ . We will verify the hypotheses from Theorem 16.6.

(i) Let us fix  $\mathbf{q} \in \mathbf{K}$ . Since the sum of finite lower semicontinuous maps is lower semicontinuous, it is enough to prove that  $\mathbf{p} \in \mathbf{K} \rightsquigarrow D_i F_i(\mathbf{p}, \min F_i(\mathbf{p}))(q_i - p_i)$  is lower semicontinuous on  $\mathbf{K}$  for each  $i \in \{1, \dots, n\}$ . To do this, let us fix an  $\mathbf{p} \in \mathbf{K}$ . Now, let

$$c_i \in D_i F_i(\mathbf{p}, \min F_i(\mathbf{p}))(q_i - p_i)$$

and  $\{\mathbf{p}^m\}$  be a sequence from  $\mathbf{K}$  which converges to  $\mathbf{p}$ . Since  $F_i|_{\mathbf{K}}$  is continuous on  $\mathbf{K}$ , the function  $\min(F_i|_{\mathbf{K}})$  is also continuous. Therefore  $\min F_i(\mathbf{p}^m) \rightarrow \min F_i(\mathbf{p})$  as  $m \rightarrow \infty$ . Using the fact that  $F_i|_{\mathbf{K}}$  is  $i$ -lower semicontinuously differentiable, there exists a sequence  $c_i^m \in D_i F_i(\mathbf{p}^m, \min F_i(\mathbf{p}^m))(q_i - p_i^m)$  such that  $c_i^m \rightarrow c_i$  as  $m \rightarrow \infty$ .

(ii) Let us consider  $\mathbf{p} \in \mathbf{K}$ , fixed. We will prove that  $\mathbf{q} \in \mathbf{K} \rightsquigarrow G(\mathbf{p}, \mathbf{q})$  is convex on  $\mathbf{K}$ . Since the sum of convex maps is also convex, it is enough to prove that  $\mathbf{q} \in \mathbf{K} \rightsquigarrow D_i F_i(\mathbf{p}, \min F_i(\mathbf{p}))(q_i - p_i)$  is convex. Using the fact that  $F_i|_{\mathbf{K}}$  is  $i$ -lower semicontinuously differentiable (in particular  $q_i \rightsquigarrow F_i(\mathbf{p}; q_i)$  is sleek at  $(p_i, \min F_i(\mathbf{p}))$ ) and Remark E.2, the above set-valued map is a closed convex process.

(iii) Let  $\mathbf{p} \in \mathbf{K}$  and  $i \in \{1, \dots, n\}$  be fixed. We prove that every element  $c_i$  from  $D_i F_i(\mathbf{p}, \min F_i(\mathbf{p}))(0)$  is non-negative. In fact, since  $q_i \rightsquigarrow F_i(\mathbf{p}; q_i)$  is  $K_i$ -locally Lipschitz, from (E.2) we have

$$\liminf_{t \rightarrow 0^+} \text{dist} \left( c_i, \frac{F_i(\mathbf{p}; p_i + t \cdot 0) - \min F_i(\mathbf{p})}{t} \right) = 0.$$

Since

$$\frac{F_i(\mathbf{p}) - \min F_i(\mathbf{p})}{t} \subseteq \mathbb{R}_+ \text{ for all } t > 0,$$

we obtain that  $c_i \geq 0$ . Therefore,

$$G(\mathbf{p}, \mathbf{p}) = \sum_{i=1}^n D_i F_i(\mathbf{p}, \min F_i(\mathbf{p}))(0) \subseteq \mathbb{R}_+.$$

By Theorem 16.6 we obtain an element  $\bar{\mathbf{p}} = (\bar{p}_1, \dots, \bar{p}_n) \in \mathbf{K}$  such that

$$\sum_{i=1}^n D_i F_i(\bar{\mathbf{p}}, \min F_i(\bar{\mathbf{p}}))(q_i - \bar{p}_i) \subseteq \mathbb{R}_+ \text{ for all } \mathbf{q} \in \mathbf{K}. \quad (16.9)$$

Let  $i \in \{1, \dots, n\}$  be fixed. We may choose  $q_j = \bar{p}_j$ ,  $j \neq i$ . Since the

(partial) contingent derivatives are closed convex process, we clearly have that  $0 \in D_j F_j(\bar{\mathbf{p}}, \min F_j(\bar{\mathbf{p}}))(0)$ ,  $j \neq i$ . From (16.9), we obtain

$$D_i F_i(\bar{\mathbf{p}}, \min F_i(\bar{\mathbf{p}}))(q_i - \bar{p}_i) \subseteq \mathbb{R}_+ \text{ for all } q_i \in K_i.$$

The proof is complete.  $\square$

Using Remark E.1, we immediately obtain a consequence of the above theorem. A similar result was obtained by Kassay-Kolumbán-Páles [155].

**Corollary 16.1** *Let  $K_1, \dots, K_n$  be nonempty compact convex subsets of a real normed space  $X$  and let  $f_i : K_1 \times \dots \times X \times \dots \times K_n \rightarrow \mathbb{R}$  ( $i \in \{1, \dots, n\}$ ) be continuous functions. Suppose that there exist open convex sets  $D_i \subseteq X$  such that  $K_i \subseteq D_i$  and  $f_i$  is continuously differentiable in the  $i^{\text{th}}$  variable on  $D_i$  and  $\partial_i f_i$  is continuous on  $K_1 \times \dots \times D_i \times \dots \times K_n$ . Then there exists an element  $\mathbf{p} \in \mathbf{K}$  such that*

$$\langle \partial_i f_i(\mathbf{p}), q_i - p_i \rangle \geq 0 \text{ for all } q_i \in K_i \text{ and all } i \in \{1, \dots, n\}. \quad (16.10)$$

**Example 16.4** Let  $K = K_1 = K_2 = [-1, 1]$ ,  $X = D_1 = D_2 = \mathbb{R}$ , and let  $f_1, f_2 : D_1 \times D_1 \rightarrow \mathbb{R}$  be defined by

$$f_1(x_1, x_2) = x_1 x_2 + x_1^2, \quad f_2(x_1, x_2) = x_2 - 3x_1 x_2^2.$$

Clearly, the above sets and functions satisfy the assumptions from Corollary 16.1, hence the existence of a solution for the system (16.10) is guaranteed. It is easy to observe that, for  $i \in \{1, 2\}$ , we have  $\partial_i f_i(x_1, x_2)(v_i - x_i) \geq 0$  for all  $v_i \in K$  if and only if

- $\partial_i f_i(x_1, x_2) = 0$  if  $-1 < x_i < 1$ ;
- $\partial_i f_i(x_1, x_2) \geq 0$ , if  $x_i = -1$  and
- $\partial_i f_i(x_1, x_2) \leq 0$  if  $x_i = 1$ .

Discussing all the cases, we establish that the set of Nash contingent points for  $(\mathbf{f}, \mathbf{K}) = (f_1, f_2; K, K)$  is  $CNP = \{(\frac{1}{2}, -1)\}$ . This point will be also a Nash equilibrium point for  $(\mathbf{f}, \mathbf{K})$ .

## 16.4 Lack of convexity: a Riemannian approach

In this section we are going to treat a case when the strategy sets  $K_i$  are *not* necessarily convex in the usual sense. In order to handle this problem a geometrical method will be exploited: we assume that one can find suitable Riemannian manifolds  $(M_i, g_i)$  such that the set  $K_i$  becomes a geodesic convex set in  $M_i$ . Note that the choice of such Riemannian structures does not influence the Nash equilibrium points for  $(\mathbf{f}, \mathbf{K})$ .

Moreover, after fixing these manifolds if the payoff functions  $f_i$  become convex on  $K_i$  (i.e.,  $f_i \circ \gamma_i : [0, 1] \rightarrow \mathbb{R}$  is convex in the usual sense for every geodesic  $\gamma_i : [0, 1] \rightarrow K_i$ ), the following existence result may be stated.

**Theorem 16.8** *Let  $(M_i, g_i)$  be finite-dimensional Riemannian manifolds,  $K_i \subset M_i$  be nonempty, compact, geodesic convex sets, and  $f_i : \mathbf{K} \rightarrow \mathbb{R}$  be continuous functions such that  $K_i \ni q_i \mapsto f_i(\mathbf{p}; q_i)$  is convex on  $K_i$  for every  $\mathbf{p} \in \mathbf{K}$ ,  $i \in \{1, \dots, n\}$ . Then, there exists a Nash equilibrium point for  $(\mathbf{f}, \mathbf{K})$ .*

*Proof* We apply Theorem 13.5 by choosing  $X = \mathbf{K} = \Pi_{i=1}^n K_i$  and  $h : X \times X \rightarrow \mathbb{R}$  defined by

$$h(\mathbf{q}, \mathbf{p}) = \sum_{i=1}^n [f_i(\mathbf{p}; q_i) - f_i(\mathbf{p})].$$

First of all, note that the sets  $K_i$  are ANRs, being closed subsets of finite-dimensional manifolds (thus, locally contractible spaces). Moreover, since a product of a finite family of ANRs is an ANR (see Bessage-Pelczyński [38, p.69]), it follows that  $X$  is an ANR. Due to Proposition 13.2,  $X$  is contractible, thus acyclic.

Note that the function  $h$  is continuous, and  $h(\mathbf{p}, \mathbf{p}) = 0$  for every  $\mathbf{p} \in X$ . Consequently, the set  $\{(\mathbf{q}, \mathbf{p}) \in X \times X : 0 > h(\mathbf{q}, \mathbf{p})\}$  is open.

It remains to prove that  $S_{\mathbf{p}} = \{\mathbf{q} \in X : 0 > h(\mathbf{q}, \mathbf{p})\}$  is contractible or empty for all  $\mathbf{p} \in X$ . Assume that  $S_{\mathbf{p}} \neq \emptyset$  for some  $\mathbf{p} \in X$ . Fix  $\mathbf{q}^j = (q_1^j, \dots, q_n^j) \in S_{\mathbf{p}}$ ,  $j \in \{1, 2\}$ , and let  $\gamma_i : [0, 1] \rightarrow K_i$  be the unique geodesic joining the points  $q_i^1 \in K_i$  and  $q_i^2 \in K_i$  (note that  $K_i$  is geodesic convex),  $i \in \{1, \dots, n\}$ . Let  $\gamma : [0, 1] \rightarrow \mathbf{K}$  defined by  $\gamma(t) = (\gamma_1(t), \dots, \gamma_n(t))$ . Due to the convexity of the function  $K_i \ni q_i \mapsto f_i(\mathbf{p}; q_i)$ , for every  $t \in [0, 1]$ , we have

$$\begin{aligned} h(\gamma(t), \mathbf{p}) &= \sum_{i=1}^n [f_i(\mathbf{p}; \gamma_i(t)) - f_i(\mathbf{p})] \\ &\leq \sum_{i=1}^n [t f_i(\mathbf{p}; \gamma_i(1)) + (1-t) f_i(\mathbf{p}; \gamma_i(0)) - f_i(\mathbf{p})] \\ &= t h(\mathbf{q}^2, \mathbf{p}) + (1-t) h(\mathbf{q}^1, \mathbf{p}) \\ &< 0. \end{aligned}$$

Consequently,  $\gamma(t) \in S_{\mathbf{p}}$  for every  $t \in [0, 1]$ , i.e.,  $S_{\mathbf{p}}$  is a geodesic convex set in the product manifold  $\mathbf{M} = \Pi_{i=1}^n M_i$  endowed with its natural (warped-)product metric (with the constant weight functions 1),

see O'Neill [233, p.208]. Now, Proposition 13.2 implies that  $S_{\mathbf{p}}$  is contractible. Alternatively, we may exploit the fact that the projections  $\text{pr}_i S_{\mathbf{p}}$  are geodesic convex, thus contractible sets,  $i \in \{1, \dots, n\}$ .

On account of Theorem 13.5, there exists  $\mathbf{p} \in \mathbf{K}$  such that  $0 = h(\mathbf{p}, \mathbf{p}) \leq h(\mathbf{q}, \mathbf{p})$  for every  $\mathbf{q} \in \mathbf{K}$ . In particular, putting  $\mathbf{q} = (\mathbf{p}; q_i)$ ,  $q_i \in K_i$  fixed, we obtain that  $f_i(\mathbf{p}; q_i) - f_i(\mathbf{p}) \geq 0$  for every  $i \in \{1, \dots, n\}$ , i.e.,  $\mathbf{p}$  is a Nash equilibrium point for  $(\mathbf{f}, \mathbf{K})$ .  $\square$

Our further investigation is motivated by the following two questions:

- What about the case when some payoff functions  $f_i$  are *not* convex on  $K_i$  in spite of the geodesic convexity of  $K_i$  on  $(M_i, g_i)$ ?
- Even for convex payoff functions  $f_i$  on  $K_i$ , how can we *localize* the Nash equilibrium points for  $(\mathbf{f}, \mathbf{K})$ ?

The following concept is destined for simultaneously handling the above questions.

Let  $(M_i, g_i)$  be complete finite-dimensional Riemannian manifolds,  $K_i \subset M_i$  be nonempty, geodesic convex sets, and  $f_i : (\mathbf{K}; D_i) \rightarrow \mathbb{R}$  functions such that  $D_i \ni q_i \mapsto f_i(\mathbf{p}; q_i)$  is of class  $C^1$  for every  $\mathbf{p} \in \mathbf{K}$ , where  $(\mathbf{K}; D_i) = K_1 \times \dots \times D_i \times \dots \times K_n$ , with  $D_i$  open and geodesic convex, and  $K_i \subseteq D_i \subseteq M_i$ ,  $i \in \{1, \dots, n\}$ .

A point  $\mathbf{p} = (p_1, \dots, p_n) \in \mathbf{K}$  is a *Nash critical point* for  $(\mathbf{f}, \mathbf{K})$  if

$$g_i(\partial_i f_i(\mathbf{p}), \exp_{p_i}^{-1}(q_i)) \geq 0 \text{ for all } q_i \in K_i, i \in \{1, \dots, n\}.$$

Here,  $\partial_i f_i(\mathbf{p})$  denotes the  $i^{\text{th}}$  partial derivative of  $f_i$  at the point  $p_i \in K_i$ .

A useful relation between Nash equilibrium points and Nash critical points is established by the following result.

**Proposition 16.3** *Any Nash equilibrium point for  $(\mathbf{f}, \mathbf{K})$  is a Nash critical point for  $(\mathbf{f}, \mathbf{K})$ . In addition, if  $D_i \ni q_i \mapsto f_i(\mathbf{p}; q_i)$  is convex for every  $\mathbf{p} \in \mathbf{K}$ ,  $i \in \{1, \dots, n\}$ , the converse also holds.*

*Proof* Let  $\mathbf{p} \in \mathbf{K}$  be a Nash equilibrium point for  $(\mathbf{f}, \mathbf{K})$ . In particular, for every  $q_i \in K_i$  with  $i \in \{1, \dots, n\}$  fixed, the geodesic segment  $t \mapsto \exp_{p_i}(t \exp_{p_i}^{-1}(q_i))$ ,  $t \in [0, 1]$  joining the points  $p_i$  and  $q_i$ , belongs entirely to  $K_i$ ; thus,

$$f_i(\mathbf{p}; \exp_{p_i}(t \exp_{p_i}^{-1}(q_i))) - f_i(\mathbf{p}) \geq 0 \text{ for all } t \in [0, 1]. \tag{16.11}$$

Combining relation (16.11) with

$$g_i(\partial_i f_i(\mathbf{p}), \exp_{p_i}^{-1}(q_i)) = \lim_{t \rightarrow 0^+} \frac{f_i(\mathbf{p}; \exp_{p_i}(t \exp_{p_i}^{-1}(q_i))) - f_i(\mathbf{p})}{t},$$

we indeed have that  $\mathbf{p} \in \mathbf{K}$  is a Nash critical point for  $(\mathbf{f}, \mathbf{K})$ .

Now, let us assume that  $\mathbf{p} \in \mathbf{K}$  is a Nash critical point for  $(\mathbf{f}, \mathbf{K})$ , i.e.,

$$\begin{aligned} 0 &\leq g_i(\partial_i f_i(\mathbf{p}), \exp_{p_i}^{-1}(q_i)) \\ &= \lim_{t \rightarrow 0^+} \frac{f_i(\mathbf{p}; \exp_{p_i}(t \exp_{p_i}^{-1}(q_i))) - f_i(\mathbf{p})}{t}. \end{aligned} \quad (16.12)$$

The function

$$g(t) = \frac{f_i(\mathbf{p}; \exp_{p_i}(t \exp_{p_i}^{-1}(q_i))) - f_i(\mathbf{p})}{t}$$

is well-defined on the whole interval  $(0, 1]$ ; indeed,  $t \mapsto \exp_{p_i}(t \exp_{p_i}^{-1}(q_i))$  is the minimal geodesic joining the points  $p_i \in K_i$  and  $q_i \in K_i$  which belongs to  $K_i \subset D_i$ . Moreover, a standard computation shows that  $t \mapsto g(t)$  is non-decreasing on  $(0, 1]$  due to the convexity of  $D_i \ni q_i \mapsto f_i(\mathbf{p}; q_i)$ . Consequently, (16.12) implies that

$$\begin{aligned} 0 &\leq \lim_{t \rightarrow 0^+} g(t) \\ &\leq g(1) = f_i(\mathbf{p}; \exp_{p_i}(\exp_{p_i}^{-1}(q_i))) - f_i(\mathbf{p}) \\ &= f_i(\mathbf{p}; q_i) - f_i(\mathbf{p}), \end{aligned}$$

which completes the proof.  $\square$

In order to state our existence result concerning Nash critical points, we consider the hypothesis

- (H)  $K_i \ni q_i \mapsto g_i(\partial_i f_i(\mathbf{p}), \exp_{p_i}^{-1}(q_i))$  is convex for every  $\mathbf{p} \in \mathbf{K}$  and  $i \in \{1, \dots, n\}$ .

**Remark 16.1** Let  $I_1, I_2 \subseteq \{1, \dots, n\}$  be such that  $I_1 \cup I_2 = \{1, \dots, n\}$ . Hypothesis (H) holds, for instance, when

- (a)  $(M_i, g_i)$  is Euclidean,  $i \in I_1$ ;  
 (b)  $K_i = \text{Im} \gamma_i$  where  $\gamma_i : [0, 1] \rightarrow M_i$  is a minimal geodesic,  $i \in I_2$ .

(a) We have  $\exp_{p_i} = p_i + \text{id}_{\mathbb{R}^{\dim M_i}}$ . Therefore, the function  $K_i \ni q_i \mapsto \langle \partial_i f_i(\mathbf{p}), q_i - p_i \rangle$  becomes affine, so convex.

(b) In this case, if  $\sigma_i : [0, 1] \rightarrow M_i$  is a geodesic segment joining the points  $\sigma_i(0) = \gamma_i(\tilde{t}_0)$  with  $\sigma_i(1) = \gamma_i(\tilde{t}_1)$  ( $0 \leq \tilde{t}_0 < \tilde{t}_1 \leq 1$ ), then  $\text{Im}(\sigma_i) \subseteq \text{Im}(\gamma_i) = K_i$ . Fix  $p_i = \gamma_i(t_i) \in K_i$  ( $0 \leq \tilde{t}_i \leq 1$ ). Let  $a_0, a_1 \in \mathbb{R}$  ( $a_0 < a_1$ ) such that  $\exp_{p_i}(a_0 \gamma_i'(t_i)) = \gamma_i(\tilde{t}_0)$  and  $\exp_{p_i}(a_1 \gamma_i'(t_i)) = \gamma_i(\tilde{t}_1)$ . Then,  $\sigma_i(t) = \exp_{p_i}((a_0 + (a_1 - a_0)t) \gamma_i'(t_i))$ . Now, the claim easily follows since the convexity of function from (H) reduces to the affinity of  $t \mapsto g_i(\partial_i f_i(\mathbf{p}), (a_0 + (a_1 - a_0)t) \gamma_i'(t_i))$ .

**Theorem 16.9** *Let  $(M_i, g_i)$  be complete finite-dimensional Riemannian manifolds,  $K_i \subset M_i$  be nonempty, compact, geodesic convex sets and  $f_i : (\mathbf{K}; D_i) \rightarrow \mathbb{R}$  continuous functions such that  $D_i \ni q_i \mapsto f_i(\mathbf{p}; q_i)$  is of class  $C^1$  for every  $\mathbf{p} \in \mathbf{K}$ ,  $i \in \{1, \dots, n\}$ . If (H) holds, there exists a Nash critical point for  $(\mathbf{f}, \mathbf{K})$ .*

*Proof* The proof is similar to that of Theorem 16.8; we show only the differences. Let  $X = \mathbf{K} = \prod_{i=1}^n K_i$  and  $h : X \times X \rightarrow \mathbb{R}$  defined by

$$h(\mathbf{q}, \mathbf{p}) = \sum_{i=1}^n g_i(\partial_i f_i(\mathbf{p}), \exp_{p_i}^{-1}(q_i)).$$

It is clear that  $h(\mathbf{p}, \mathbf{p}) = 0$  for every  $\mathbf{p} \in X$ .

First of all, the (upper-semi)continuity of  $h(\cdot, \cdot)$  on  $X \times X$  implies the fact that the set  $\{(\mathbf{q}, \mathbf{p}) \in X \times X : 0 > h(\mathbf{q}, \mathbf{p})\}$  is open.

Now, let  $\mathbf{p} \in X$  such that  $S_{\mathbf{p}} = \{\mathbf{q} \in X : 0 > h(\mathbf{q}, \mathbf{p})\}$  is not empty. Now, let us fix  $\mathbf{q}^j = (q_1^j, \dots, q_n^j) \in S_{\mathbf{p}}$ ,  $j \in \{1, 2\}$ , and let  $\gamma_i : [0, 1] \rightarrow K_i$  be the unique geodesic joining the points  $q_i^1 \in K_i$  and  $q_i^2 \in K_i$ . Let also  $\gamma : [0, 1] \rightarrow \mathbf{K}$  defined by  $\gamma(t) = (\gamma_1(t), \dots, \gamma_n(t))$ . Due to hypotheses (H), the convexity of the function  $[0, 1] \ni t \mapsto h(\gamma(t), \mathbf{p})$ ,  $t \in [0, 1]$  easily follows. Therefore,  $\gamma(t) \in S_{\mathbf{p}}$  for every  $t \in [0, 1]$ , i.e.,  $S_{\mathbf{p}}$  is a geodesic convex set, thus contractible.

Theorem 13.5 implies the existence of  $\mathbf{p} \in \mathbf{K}$  such that  $0 = h(\mathbf{p}, \mathbf{p}) \leq h(\mathbf{q}, \mathbf{p})$  for every  $\mathbf{q} \in \mathbf{K}$ . In particular, if  $\mathbf{q} = (\mathbf{p}; q_i)$ ,  $q_i \in K_i$  fixed, we obtain that  $g_i(\partial_i f_i(\mathbf{p}), \exp_{p_i}^{-1}(q_i)) \geq 0$  for every  $i \in \{1, \dots, n\}$ , i.e.,  $\mathbf{p}$  is a Nash critical point for  $(\mathbf{f}, \mathbf{K})$ . The proof is complete.  $\square$

**Remark 16.2** Proposition 16.3 and Theorem 16.9 give together a possible answer to the location of Nash equilibrium points. Indeed, on account of Theorem 16.9 we are able to find explicitly the Nash critical points for  $(\mathbf{f}, \mathbf{K})$ ; then, due to Proposition 16.3, among Nash critical points for  $(\mathbf{f}, \mathbf{K})$  we may choose the Nash equilibrium points for  $(\mathbf{f}, \mathbf{K})$ .

**Remark 16.3** The set of Nash critical points and the set of Nash equilibrium points are usually different. Indeed, let  $f_1, f_2 : [-1, 1]^2 \rightarrow \mathbb{R}$  be defined by  $f_1(x_1, x_2) = f_2(x_1, x_2) = x_1^3 + x_2^3$ . Note that  $(0, 0)$ ,  $(0, -1)$ ,  $(-1, 0)$  and  $(-1, -1)$  are Nash critical points, but only  $(-1, -1)$  is a Nash equilibrium point for  $(f_1, f_2; [-1, 1], [-1, 1])$ .

The following examples show the applicability of our results.

**Example 16.5** Let  $K_1 = [-1, 1]$ ,  $K_2 = \{(\cos t, \sin t) : t \in [\pi/4, 3\pi/4]\}$ ,

and  $f_1, f_2 : K_1 \times K_2 \rightarrow \mathbb{R}$  defined for every  $x \in K_1, (y_1, y_2) \in K_2$  by

$$f_1(x, (y_1, y_2)) = |x|y_1^2 - y_2, \quad f_2(x, (y_1, y_2)) = (1 - |x|)(y_1^2 - y_2^2).$$

Note that  $K_1 \subset \mathbb{R}$  is convex in the usual sense, but  $K_2 \subset \mathbb{R}^2$  is not. However, if we consider the Poincaré upper-plane model  $(\mathbb{H}^2, g_{\mathbb{H}})$ , the set  $K_2 \subset \mathbb{H}^2$  is geodesic convex with respect to the metric  $g_{\mathbb{H}}$ , being the image of a geodesic segment from  $(\mathbb{H}^2, g_{\mathbb{H}})$ . It is clear that  $f_1(\cdot, (y_1, y_2))$  is a convex function on  $K_1$  in the usual sense for every  $(y_1, y_2) \in K_2$ . Moreover,  $f_2(x, \cdot)$  is also a convex function on  $K_2 \subset \mathbb{H}^2$  for every  $x \in K_1$ . Indeed, the latter fact reduces to the convexity of the function  $t \mapsto (1 - |x|)\cos(2t)$ ,  $t \in [\pi/4, 3\pi/4]$ . Therefore, Theorem 16.8 guarantees the existence of at least one Nash equilibrium point for  $(f_1, f_2; K_1, K_2)$ . Using Proposition 16.3, a simple calculation shows that the set of Nash equilibrium (=critical) points for  $(f_1, f_2; K_1, K_2)$  is  $K_1 \times \{(0, 1)\}$ .

**Example 16.6** Let  $K_1 = [-1, 1]^2$ ,  $K_2 = \{(y_1, y_2) : y_2 = y_1^2, y_1 \in [0, 1]\}$ , and  $f_1, f_2 : K_1 \times K_2 \rightarrow \mathbb{R}$  defined for every  $(x_1, x_2) \in K_1, (y_1, y_2) \in K_2$  by

$$f_1((x_1, x_2), (y_1, y_2)) = -x_1^2 y_2 + x_2 y_1,$$

$$f_2((x_1, x_2), (y_1, y_2)) = x_1 y_2^2 + x_2 y_1^2.$$

The set  $K_1 \subset \mathbb{R}^2$  is convex, but  $K_2 \subset \mathbb{R}^2$  is not in the usual sense. However,  $K_2$  may be considered as the image of a geodesic segment on the paraboloid of revolution  $p_{\text{rev}}(u, v) = (v \cos u, v \sin u, v^2)$  endowed with its natural Riemannian structure having the coefficients

$$g_{11}(u, v) = v^2, \quad g_{12}(u, v) = g_{21}(u, v) = 0, \quad g_{22}(u, v) = 1 + 4v^2.$$

More precisely,  $K_2$  becomes geodesic convex on  $\text{Im}(p_{\text{rev}})$ , being actually a compact part of a meridian on  $\text{Im}(p_{\text{rev}})$ . Note that neither  $f_1(\cdot, (y_1, y_2))$  nor  $f_2((x_1, x_2), \cdot)$  is convex (the convexity of the latter function being considered on  $K_2 \subset \text{Im}(p_{\text{rev}})$ ); thus, Theorem 16.8 is not applicable. In view of Remark 16.1, Theorem 16.9 can be applied in order to determine the set of Nash critical points. This set is nothing but the set of solutions in the form  $((\tilde{x}_1, \tilde{x}_2), (\tilde{y}, \tilde{y}^2)) \in K_1 \times K_2$  of the system

$$\begin{cases} -2\tilde{x}_1 \tilde{y}^2 (x_1 - \tilde{x}_1) + \tilde{y} (x_2 - \tilde{x}_2) \geq 0, & \forall (x_1, x_2) \in K_1, \\ \tilde{y} (2\tilde{y}^2 \tilde{x}_1 + \tilde{x}_2) (y - \tilde{y}) \geq 0, & \forall y \in [0, 1]. \end{cases} \quad (\text{System}_{\text{NCP}})$$

Note that the second inequality is obtained from

$$\partial_2 f_2((x_1, x_2), (y_1, y_2)) = \left( \frac{2x_2 y_1}{y_2^2}, \frac{2x_1 y_2}{1 + 4y_2^2} \right)$$

and

$$\exp_{(\tilde{y}, \tilde{y}^2)}^{-1}(y, y^2) = (y - \tilde{y}, 2\tilde{y}(y - \tilde{y})), \quad \tilde{y}, y \in [0, 1].$$

We distinguish three cases: (a)  $\tilde{y} = 0$ ; (b)  $\tilde{y} = 1$ ; and (c)  $0 < \tilde{y} < 1$ .

(a)  $\tilde{y} = 0$ . Then, any  $((\tilde{x}_1, \tilde{x}_2), (0, 0)) \in K_1 \times K_2$  solves  $(\text{System}_{\text{NCP}})$ .

(b)  $\tilde{y} = 1$ . After an easy computation, we obtain that  $((-1, -1), (1, 1)) \in K_1 \times K_2$  and  $((0, -1), (1, 1)) \in K_1 \times K_2$  solve  $(\text{System}_{\text{NCP}})$ .

(c)  $0 < \tilde{y} < 1$ . The unique situation when  $(\text{System}_{\text{NCP}})$  is solvable is  $\tilde{y} = \sqrt{2}/2$ . In this case,  $(\text{System}_{\text{NCP}})$  has a unique solution  $((1, -1), (\sqrt{2}/2, 1/2)) \in K_1 \times K_2$ .

Consequently, the set of Nash critical points for  $(f_1, f_2; K_1, K_2)$ , denoted in the sequel by  $S_{\text{NCP}}$ , is the union of the points from (a), (b) and (c), respectively.

Let us denote by  $S_{\text{NEP}}$  the set of Nash equilibrium points for  $(f_1, f_2; K_1, K_2)$ .

Due to Proposition 16.3, we may select the elements of  $S_{\text{NEP}}$  from  $S_{\text{NCP}}$ .

Therefore, the elements of  $S_{\text{NEP}}$  are the solutions  $((\tilde{x}_1, \tilde{x}_2), (\tilde{y}, \tilde{y}^2)) \in S_{\text{NCP}}$  of the system

$$\begin{cases} -x_1^2\tilde{y}^2 + x_2\tilde{y} \geq -\tilde{x}_1^2\tilde{y}^2 + \tilde{x}_2\tilde{y}, & \forall (x_1, x_2) \in K_1, \\ \tilde{x}_1y^4 + \tilde{x}_2y^2 \geq \tilde{x}_1\tilde{y}^4 + \tilde{x}_2\tilde{y}^2, & \forall y \in [0, 1]. \end{cases} \quad (\text{System}_{\text{NEP}})$$

We consider again the above three cases.

(a)  $\tilde{y} = 0$ . Among the elements  $((\tilde{x}_1, \tilde{x}_2), (0, 0)) \in K_1 \times K_2$  which solve  $(\text{System}_{\text{NCP}})$ , only those are solutions for  $(\text{System}_{\text{NEP}})$  which fulfill the condition  $\tilde{x}_2 \geq \max\{-\tilde{x}_1, 0\}$ .

(b)  $\tilde{y} = 1$ . We have  $((-1, -1), (1, 1)) \in S_{\text{NEP}}$ , but  $((0, -1), (1, 1)) \notin S_{\text{NEP}}$ .

(c)  $0 < \tilde{y} < 1$ . We have  $((1, -1), (\sqrt{2}/2, 1/2)) \in S_{\text{NEP}}$ .

## 16.5 Historical comments and perspectives

*A. Historical comments.* Although many extensions and applications can be found of Nash's result (see, for instance, Chang [67], Georgiev [127], Guillermo [136], Kulpa-Szymanski [184], Luo [199], Morgan-Scalzo [218], Yu-Zhang [300], and references therein), to the best of our knowledge, only a few works can be found in the literature dealing with the question addressed in the Introduction, i.e., the location of Nash equilibrium points for a functions with non-usual properties (for instance, set-valued maps) which are defined on sets with non-standard structures. Nessah-Kerstens [229] characterize the existence of Nash equilibrium points on

non-convex strategy sets via a modified diagonal transfer convexity notion; Tala-Marchi [288] also treat games with non-convex strategies reducing the problem to the convex case via certain homeomorphism; Kassay-Kolumbán-Páles [155] and Ziad [304] considered Nash equilibrium points on convex domains for non-convex payoff functions having suitable regularity instead of their convexity.

In this chapter we treated the existence and location of Nash equilibrium points in various context. Via variational inequalities and dynamical systems, we established stability results concerning Nash critical points following the paper of Cavazzuti-Pappalardo-Passacantando [63]. In Section 16.3 we studied Nash equilibrium points for set-valued maps on vector spaces. The purpose of the last section was to initiate a new approach concerning the study of Nash equilibria, where the payoff functions are defined on sets which are not necessarily convex in the usual sense, by embedding these sets into suitable Riemannian manifolds.

*B. Perspectives.* The main challenging question is to study the (exponential) stability of Nash critical points for functions which are defined on sets embedded into certain Riemannian manifolds. Since the non-expansiveness and the validity of Moskovitz-Dines property of the projection operator are indispensable in such an argument (see Theorem 16.4), we believe that this fact is possible only for those cases where the strategy sets can be embedded into Hadamard-type Riemannian manifolds (see Theorems 15.4 and 15.8).

# 17

## Problems to Part III

God exists since mathematics  
is consistent, and the Devil  
exists since we cannot prove it.

---

André Weil (1906–1998)

**Problem 17.1** Prove that  $(\text{Int}(M_C), d_H)$  is a metric space, where  $C$  is a simple, closed and convex curve and  $d_H$  is the Hilbert distance. [Hint: See Section 13.2.]

**Problem 17.2** Let  $C$  be an ellipse with semi-axes  $a > b > 0$ , and consider three points  $P_1 = (a/2, 0)$ ,  $P_2 = (-a/2, 0)$  and  $P_3 = (a/2, -b/2)$  in the interior of the ellipse  $\text{Int}(M_C)$ . Determine the minimum points of the total cost function  $C(P_i, 3, s) = \sum_{i=1}^3 d_H^s(\cdot, P_i)$  on  $\text{Int}(M_C)$  for  $s = 1$  and  $s = 2$ , respectively.

**Problem 17.3** Let  $M = \mathbb{R}^m$ ,  $m \geq 2$ , be endowed with the natural Euclidean metric, and  $P_i \in M$  ( $i \in \{1, \dots, n\}$ ) given points. Prove that the unique minimum point  $T_f = T_b$  (i.e., the place of the deposit) for  $C_f(P_i, n, s) = C_b(P_i, n, s)$  is the center of gravity of points  $\{P_1, \dots, P_n\}$  if and only if  $s = 2$ . [Hint: See Example 14.1.]

**Problem 17.4** There is a ship which moves on a closed curve and there are  $n$  fixed ambulance stations. Suppose that at a certain moment, the ship's crew asks for first-aid continuing their trip. Determine the position of ambulance station from where the team assistance reaches the ship within minimal time. [Hint: Follow Problem 14.1.]

**Problem 17.5** If  $M_0$  is a submanifold of the Riemannian manifold  $(M, g)$  and  $\gamma : [0, 1] \rightarrow M$  is a geodesic of  $M$  with the properties that

$\gamma(1) \in M_0$  and  $d_g(\gamma(0), y) \geq L_g(\gamma)$  for every  $y \in M_0$ , then  $g(\dot{\gamma}(1), X) = 0$  for every  $X \in T_{\gamma(1)}M_0$ , i.e.,  $\gamma$  is perpendicular to  $M_0$  at the point of contact. [Hint: Use the first variational formula and follow Proposition 15.1.]

**Problem 17.6** If  $M_1$  and  $M_2$  are submanifolds of the Riemannian manifold  $(M, g)$  and  $\gamma : [0, 1] \rightarrow M$  is a geodesic of  $M$  with the properties that  $x_0 = \gamma(0) \in M_1$ ,  $y_0 = \gamma(1) \in M_2$ , and  $d_g(x, y) \geq d_g(x_0, y_0) = L_g(\gamma)$  for every  $(x, y) \in M_1 \times M_2$ , then  $g(\dot{\gamma}(0), X) = 0$  and  $g(\dot{\gamma}(1), Y) = 0$  for every  $(X, Y) \in T_{\gamma(0)}M_1 \times T_{\gamma(1)}M_2$ , i.e.,  $\gamma$  is a common perpendicular of  $M_1$  and  $M_2$ . [Hint: Use the first variational formula and follow Proposition 15.1.]

**Problem 17.7** Let  $(M, g)$  be a simply connected Riemannian manifold with non-positive sectional curvature and let  $\alpha, \beta : \mathbb{R} \rightarrow M$  be two geodesics. We allow that  $\beta(t) = \text{constant}$ , and  $\|\dot{\alpha}\| = 1$ . The distance function  $f(t) = d_g(\alpha(t), \beta(t))$  has the following properties:

- 1) the equation  $f(t) = 0$  has at most one solution;
- 2)  $f$  is a function of class  $C^\infty$  for all  $t \in \mathbb{R}$  with  $f(t) \neq 0$ .

[Hint: Use the first and second variational formula.]

**Problem 17.8** Let  $K_1 = [-5, 3]$ ,  $K_2 = \{(\cos t, \sin t) : t \in [\pi/4, 3\pi/4]\}$ , and  $f_1, f_2 : K_1 \times K_2 \rightarrow \mathbb{R}$  defined for every  $x \in K_1$ ,  $(y_1, y_2) \in K_2$  by  $f_1(x, (y_1, y_2)) = x^2y_1^3 - y_2^2$ ,  $f_2(x, (y_1, y_2)) = (16 - x^2)(y_1^2 - y_2^2)$ . Determine the set of Nash equilibrium points for  $(\mathbf{f}, \mathbf{K})$ .

**Problem 17.9** Let  $K_1 = \{(\sin t, \cos t) : t \in [\pi/8, 3\pi/7]\}$ ,  $K_2 = \{(y_1, y_2) : y_2 = y_1^2 + 2, y_1 \in [0, 2]\}$ , and  $f_1, f_2 : K_1 \times K_2 \rightarrow \mathbb{R}$  defined for every  $(x_1, x_2) \in K_1$ ,  $(y_1, y_2) \in K_2$  by  $f_1((x_1, x_2), (y_1, y_2)) = 3x_1y_2^2 + 2x_2^3y_1^3$ ,  $f_2((x_1, x_2), (y_1, y_2)) = 4x_1^2y_2^3 + 5x_2^2y_1$ . Determine the set of Nash equilibrium points for  $(\mathbf{f}, \mathbf{K})$ .

**Problem 17.10** Let  $K = K_1 = K_2 = [-2, 3]$ ,  $X = \mathbb{R}$ , and define the set-valued maps  $F_1, F_2 : K \times K \rightsquigarrow \mathbb{R}$  defined by  $F_1(x_1, x_2) = [\max\{x_1^2, x_2^3\} - 1, 0]$  and  $F_2(x_1, x_2) = [1 - \max\{|x_1|, x_2^2\}, 6 - x_1 - x_2^2]$ . Determine the set of Nash contingent points as well as the Nash equilibrium points for  $(\mathbf{F}, \mathbf{K}) = (F_1, F_2; K, K)$ .

**Problem 17.11** (Location of Post Office) Let  $P_1 = (1, 0) \in \mathbb{R}^2$  and  $P_2 = (0, 1) \in \mathbb{R}^2$  be two Post Offices, and let two players moving on the set  $\mathbf{K} = K \times K$  where  $K = (-\infty, 0]$ . The purpose of the  $i^{\text{th}}$  player is to minimize the distance between  $(x_1, x_2) \in \mathbf{K}$  and his favorite goal  $P_i$ . What about the case when the players moving area is subject to the

social constraint  $S = \{(x_1, x_2) \in \mathbf{K} : x_1 + x_2 \leq -2\}$ ? [Hint: Use the Nash point notions from Section 16.2.]



## Appendices



# A

## Elements of convex analysis

By plucking her petals, you do  
not gather the beauty of the  
flower.

---

Rabindranath Tagore  
(1861–1941)

### A.1 Convex sets and convex functions

Mathematically, the notion of *convex set* can be made precise by defining the *segment* joining any two points  $x$  and  $y$  in a vector space to be the set of all points having the form  $\lambda x + (1 - \lambda)y$  for  $0 \leq \lambda \leq 1$ . A set is convex if and only if it contains all the segments joining any two of its points. The geometric nature of this definition is depicted in Figures A.1 and A.2.

Given a collection of convex sets, it is always possible to create other convex sets by the operations of *dilation*, *sum* or *intersection*. More precisely, if  $A$  and  $B$  are convex sets in a linear space then the sets

$$\alpha A := \{\alpha x; x \in A\} \quad \text{where } \alpha \in \mathbb{R}$$

and

$$A + B := \{a + b; a \in A, b \in B\}$$

are convex, too. Moreover, any intersection of convex sets is convex.

The notion of convex function was introduced in the first part of the twentieth century, though it was implicitly used earlier by Gibbs and Maxwell in order to describe relationships between thermodynamic variables.




Figure A.1 Geometric illustration of a convex set.




Figure A.2 Shape of a nonconvex set.

**Definition A.1** Let  $C$  be a convex set in a linear space. A function  $f : C \rightarrow \mathbb{R}$  is convex if and only if

$$f(\lambda x + (1 - \lambda)y) \leq \lambda f(x) + (1 - \lambda)f(y) \quad \text{for any } \lambda \in [0, 1].$$




Figure A.3 Geometric illustration of a convex function.




Figure A.4 Geometric illustration of the epigraph of a function  $f : [a, b] \rightarrow \mathbb{R}$ .

Thus, a functional is convex if the segment connecting two points on its graph lies above the graph of the function, as shown in Figure A.3. Now, this definition is often expressed concisely by defining the set of all points that lie above the graph of a functional to be the *epigraph* (see Figure A.4). This set plays an important role in optimization theory and is usually referred in economics as an *upper contour set*.

**Definition A.2** The epigraph of a functional  $f$  acting on a vector space  $X$  is the subset

$$\text{epi}(f) := \{(x, a) \in X \times \mathbb{R}; f(x) \leq a\}.$$

The following result makes clear how the notions of convexity of a set and convexity of a functional are related.

**Proposition A.1** Suppose  $f$  is a functional defined on a convex set. Then  $f$  is a convex functional if and only if the set  $\text{epi}(f)$  is convex.

We have seen in this volume that convexity plays an important role in many optimization problems. In addition, some classes of inequality constraints couched in terms of *convex cones* are amenable to Lagrange multiplier methods for abstract spaces. Convex cones are also used to define orderings on vector spaces, which in turn facilitate the introduction of inequality constraints.




Figure A.5 A convex closed cone with vertex at the origin in  $\mathbb{R}^2$ .

**Definition A.3** Let  $X$  be a vector space. A set  $C_0 \subset X$  is a cone with vertex at the origin if

$$x \in X \quad \text{implies} \quad \alpha x \in C_0,$$

for all  $\alpha \geq 0$ . A cone with vertex at a point  $v \in X$  is defined to be the translation  $v + C$  of a cone  $C_0$  with vertex at the origin:

$$C = \{v + x; x \in C_0\}.$$

A positive cone  $C_0^+ \subset X$  is defined by

$$C_0^+ := \{x; x \in C_0, x \geq 0\}.$$

A cone with vertex at the origin need not be convex. We also point out that a cone need not be closed, in general. Examples of cones (either convex or not convex) are depicted in Figures A.5, A.6, and A.7.

The next theorem is a result due to Efimov and Stechkin in the theory of best approximation. Roughly speaking, this property assures that in a certain class of Banach spaces, every sequentially weakly closed Chebyshev set is convex.

**Theorem A.1** (Lemma 1 of [101], Theorem 2 of [290]) *Let  $X$  be a uniformly convex Banach space with strictly convex topological dual. Assume that  $M$  is a sequentially weakly closed non-convex subset of  $X$ .*

*Then, for any convex dense subset  $S$  of  $X$ , there exists  $x_0 \in S$  such*




Figure A.6 A nonconvex closed cone with vertex at the origin in  $\mathbb{R}^2$ .




Figure A.7 A nonconvex open cone with vertex at the origin in  $\mathbb{R}^2$ .

that the set

$$\{y \in M : \|y - x_0\| = d(x_0, M)\}$$

contains at least two points.

## A.2 Convex analysis in Banach spaces

Throughout this section we assume that  $(X, \|\cdot\|)$  is a Banach space.

**Definition A.4** Let  $f : X \rightarrow (-\infty, +\infty]$  be a function and let  $x \in X$  be a point where  $f$  is finite. The one-sided directional derivative of  $f$  at  $x$  with respect to a vector  $y \in X$  is

$$f'(x; y) = \lim_{\lambda \searrow 0} \frac{f(x + \lambda y) - f(x)}{\lambda},$$

if this limit exists.

**Lemma A.1** Let  $f : X \rightarrow (-\infty, +\infty]$  be a convex function, and let  $x \in X$  be a point where  $f(x)$  is finite. For each  $y \in X$ , the difference quotient in the definition of  $f'(x; y)$  is a non-decreasing function of  $\lambda > 0$ , so that  $f'(x; y)$  exists and

$$f'(x; y) = \lim_{\lambda \searrow 0} \frac{f(x + \lambda y) - f(x)}{\lambda}.$$

Moreover,  $f'(x; y)$  is a positively homogeneous convex of  $y \in X$ .

**Definition A.5** Let  $f : X \rightarrow (-\infty, +\infty]$  be a convex function. An element  $x^* \in X^*$  is said to be a subgradient of the convex function  $f$  at a point  $x \in X$  (see Fig. A.8) provided that for any  $z \in X$ ,

$$f(z) \geq f(x) + \langle x^*, z - x \rangle.$$

The set of all subgradient of  $f$  at  $x$  is called the subdifferential of  $f$  at  $x$  and is denoted by  $\partial f(x)$ .




Figure A.8 Subgradients of a convex function at some point.

**Lemma A.2** Let  $f : X \rightarrow (-\infty, +\infty]$  be a convex function and let  $x \in X$  be a point where  $f(x)$  is finite. Then  $x^* \in \partial f(x)$  if and only if for all  $y \in X$ ,  $f'(x; y) \geq \langle x^*, y \rangle$ .

Now, let  $S \subset X$  be a non-empty closed convex set and denote by  $j_S : X \rightarrow (-\infty, +\infty]$  the indicator function associated to  $S$ , that is,

$$j_S(x) = \begin{cases} 0, & \text{if } x \in S \\ +\infty, & \text{if } x \in X \setminus S. \end{cases} \quad (\text{A.1})$$

In this case,

$$\partial j_S(x) = \{x^* \in X^* ; \langle x^*, z - x \rangle \leq 0, \text{ for all } z \in S\},$$

hence  $\partial j_S(x)$  coincide with the normal cone associated to  $S$  at the point  $x \in S$ . Thus,  $\partial j_S(x) = N_S(x)$ .

Let  $S \subset X$  be a nonempty set. We denote the *distance function*  $\text{dist}_S : X \rightarrow \mathbb{R}$  defined by

$$\text{dist}_S(x) = \inf\{\|x - y\| : y \in S\}.$$

It is clear that  $\text{dist}_S$  is a globally Lipschitz function. Consequently, one may define its *directional derivative* in the sense of Clarke at the point  $x \in S$  with direction  $v \in X$ , denoted by  $\text{dist}_S^0(x; v)$ .

**Definition A.6** A vector  $v \in X$  is tangent to  $S$  at some point  $x \in S$  if  $\text{dist}_S^0(x; v) = 0$ . The set of tangent vectors of  $S$  at  $x$  is denoted by  $T_S(x)$ .

Note that  $T_S(x)$  is a closed convex cone in  $X$ ; in particular,  $0 \in T_S(x)$ .

**Definition A.7** Let  $x \in S$ . The normal cone to  $S$  at  $x$  is

$$N_S(x) = \{x^* \in X^* : \langle x^*, v \rangle \leq 0 \text{ for all } v \in T_S(x)\}.$$

**Proposition A.2** Let  $X = X_1 \times X_2$ , where  $X_1, X_2$  are Banach spaces, and let  $x = (x_1, x_2) \in S_1 \times S_2$ , where  $S_1, S_2$  are subsets of  $X_1, X_2$ , respectively. Then

$$T_{S_1 \times S_2}(x) = T_{S_1}(x_1) \times T_{S_2}(x_2); \quad N_{S_1 \times S_2}(x) = N_{S_1}(x_1) \times N_{S_2}(x_2).$$

# B

## Function spaces

All the effects of Nature are  
only the mathematical  
consequences of a small  
number of immutable laws.

---

Pierre-Simon Laplace  
(1749–1827)

In this part we recall some basic facts on Lebesgue and Sobolev spaces. A central place is dedicated to the main theorems in these basic function spaces.

### B.1 Lebesgue spaces

Let  $\Omega \subset \mathbb{R}^N$  be an open set. We recall that the Lebesgue spaces are defined by

$$L^p(\Omega) := \left\{ u : \Omega \rightarrow \mathbb{R}; u \text{ is measurable and } \int_{\Omega} |u(x)|^p dx < \infty \right\}$$

if  $1 \leq p < \infty$ , and

$$L^\infty(\Omega) := \{ u : \Omega \rightarrow \mathbb{R}; u \text{ is measurable and } \operatorname{ess\,sup}_{x \in \Omega} |u(x)| < \infty \} .$$

For any  $1 \leq p \leq \infty$  we define the space

$$L^p_{\text{loc}}(\Omega) := \{ u : \Omega \rightarrow \mathbb{R}; u \in L^p(\omega) \text{ for each } \omega \subset\subset \Omega \} .$$

We have used in this volume the following basic results on Lebesgue spaces.

**Theorem B.1** (*Fatou's Lemma*). Assume that for any  $n \geq 1$ ,  $u_n \in L^1(\Omega)$  and  $u_n \geq 0$  a.e. on  $\Omega$ . Then

$$\int_{\Omega} \liminf_{n \rightarrow \infty} u_n \, dx \leq \liminf_{n \rightarrow \infty} \int_{\Omega} u_n \, dx.$$

The next theorem is due to Brezis and Lieb [49] provides us with a correction term that changes Fatou's lemma from an inequality to an equality.

**Theorem B.2** (*Brézis–Lieb Lemma*). Let  $(u_n)$  be a sequence of functions defined on  $\Omega$  that converges pointwise a.e. to a function  $u$ . Assume that for some  $1 \leq p < \infty$ , there exists a positive constant  $C$  such that for all  $x \in \Omega$ ,

$$\sup_{n \geq 1} \int_{\Omega} |u_n(x)|^p \, dx < C.$$

Then

$$\lim_{n \rightarrow \infty} \int_{\Omega} [|u_n(x)|^p - |u_n(x) - u(x)|^p] \, dx = \int_{\Omega} |u(x)|^p \, dx.$$

**Theorem B.3** (*Lebesgue's Dominated Convergence Theorem*). Assume that for any  $n \geq 1$ ,  $u_n \in L^1(\Omega)$ . Suppose that  $u_n \rightarrow u$  a.e. on  $\Omega$  and  $|u_n| \leq v$  a.e., for some function  $v \in L^1(\Omega)$ . Then

$$\lim_{n \rightarrow \infty} \int_{\Omega} u_n(x) \, dx = \int_{\Omega} u(x) \, dx.$$

## B.2 Sobolev spaces

Let  $u \in L^1_{\text{loc}}(\Omega)$  be a function. A function  $v_{\alpha} \in L^1_{\text{loc}}(\Omega)$  such that

$$\int_{\Omega} u(x) D_{\alpha} \varphi(x) \, dx = (-1)^{|\alpha|} \int_{\Omega} v_{\alpha}(x) \varphi(x) \, dx$$

for any  $\varphi \in C^{\infty}_0(\Omega)$  is called the weak distributional derivative of  $u$  and is denoted by  $D^{\alpha}u$ . Here  $\alpha = (\alpha_1, \dots, \alpha_N)$ , where  $\alpha_i$  ( $i = 1, \dots, N$ ) are nonnegative integers and  $D^{\alpha}u = \frac{\partial^{|\alpha|}}{\partial x_1^{\alpha_1} \dots \partial x_N^{\alpha_N}}$ , with  $|\alpha| = \alpha_1 + \dots + \alpha_N$ .

It is clear that if such a  $v_{\alpha}$  exists, it is unique up to sets of measure zero.

To define a Sobolev space we introduce a functional  $\|\cdot\|_{m,p}$ , where  $m$

is a nonnegative integer and  $1 \leq p \leq \infty$ , as follows

$$\|u\|_{m,p} = \left( \sum_{0 \leq |\alpha| \leq m} \int_{\Omega} |D^{\alpha}u(x)|^p dx \right)^{1/p},$$

for  $1 \leq p < \infty$  and

$$\|u\|_{m,\infty} = \max_{0 \leq |\alpha| \leq m} \sup_{\Omega} |D^{\alpha}u(x)|$$

if  $p = \infty$ . It is obvious that  $\|\cdot\|_{m,p}$  and  $\|\cdot\|_{m,\infty}$  define norms on any vector space of functions for which values of these functionals are finite, provided functions are identified in the space if they are equal almost everywhere.

For any integer  $m \geq 1$  and any  $1 \leq p \leq \infty$  we can define the following spaces

$$H^{m,p}(\Omega) = \text{the completion of } \{u \in C^m(\Omega) : \|u_{m,p}\| < \infty\}$$

with respect to the norm  $\|\cdot\|_{m,p}$ ;

$$W^{m,p}(\Omega) = \{u \in L^p(\Omega) : D^{\alpha}u \in L^p(\Omega) \text{ for } 0 \leq |\alpha| \leq m\},$$

where  $D^{\alpha}$  denotes the weak partial derivative of  $u$ ;

$$W_0^{m,p}(\Omega) = \text{the closure of } C_0^{\infty}(\Omega) \text{ in the space } W^{m,p}(\Omega).$$

The above spaces equipped with the norms  $\|\cdot\|_{m,p}$  are called Sobolev spaces over  $\Omega$ .

It is clear that  $W^{0,p} = W_0^{0,p} = L^p$ . It is also known that  $H^{m,p}(\Omega) = W^{1,p}(\Omega)$  for every domain  $\Omega \subset \mathbb{R}^N$ . The spaces  $W^{m,p}(\Omega)$  and  $W_0^{m,p}(\Omega)$  are Banach spaces and these spaces are reflexive if and only if  $1 < p < \infty$ . Moreover,  $W^{k,2}(\Omega)$  and  $W_0^{k,2}(\Omega)$  are Hilbert spaces with scalar product

$$\langle u, v \rangle = \sum_{|\alpha| \leq k} \int_{\Omega} D^{\alpha}u(x) D^{\alpha}v(x) dx,$$

where  $D^0u = u$ . Also,  $W^{m,p}(\Omega)$  and  $W_0^{m,p}(\Omega)$  are separable for  $1 \leq p < \infty$ . For any  $1 \leq p < \infty$  we denote  $p' = p/(p-1)$  (if  $p = 1, p' = \infty$ ) and we denote by  $W^{-m,p'}(\Omega)$  the dual space of  $W_0^{m,p}(\Omega)$  and we write

$$W^{-m,p'}(\Omega) = (W_0^{m,p}(\Omega))^*.$$

Then every element  $T \in W^{-m,p'}(\Omega)$  has the form

$$T(u) = \sum_{|\alpha| \leq k} \int_{\Omega} D^{\alpha}u(x) v_{\alpha}(x) dx,$$

where  $v_\alpha$  are suitable elements in  $L^{p'}(\Omega)$ .

### B.3 Compact embedding results

Let  $(X, \|\cdot\|_X), (Y, \|\cdot\|_Y)$  be Banach spaces. We say that  $X$  is continuously embedded into  $Y$  if there exists an injective linear map  $i : X \rightarrow Y$  and a constant  $C$  such that  $\|i(x)\|_Y \leq C\|x\|_X$  for all  $x \in X$ . We identify  $X$  with the image  $i(X)$ . We say that  $X$  is compactly embedded into  $Y$  if  $i$  is a compact map, that is,  $i$  maps bounded subsets of  $X$  into relatively compact subset of  $Y$ .

The following theorem has a particular importance in the variational and qualitative analysis of differential and partial differential equations, due to the control over the nonlinear terms.

**Theorem B.4** (Sobolev embedding theorem) *Suppose that  $\Omega \subset \mathbb{R}^N$  is a bounded domain with Lipschitz boundary. Then*

- (i) *If  $kp < N$ , then  $W^{k,p}(\Omega)$  is continuously embedded into  $L^q(\Omega)$  for each  $1 \leq q \leq \frac{Np}{N-kp}$ ; this embedding is compact, if  $q < \frac{Np}{N-kp}$ .*
- (ii) *If  $0 \leq m < k - \frac{N}{p} < m + 1$ , then  $W^{k,p}(\Omega)$  is continuously embedded into  $C^{m,\alpha}(\Omega)$ , for  $0 \leq \alpha \leq k - m - \frac{m}{p}$ , this embedding is compact if  $\alpha < k - m - \frac{m}{p}$ .*

If  $\Omega \subset \mathbb{R}^N$  is a bounded domain with Lipschitz boundary then the Poincaré inequality holds for all  $u \in W_0^{1,p}(\Omega)$ :

$$\int_{\Omega} u^2 dx \leq C \int_{\Omega} |\nabla u|^2 dx,$$

where  $C = C(\Omega)$  is a constant, possibly depending on  $\Omega$  but independent of  $u$ .

As a consequence of the Poincaré inequality it follows that

$$\|u\| = \left( \int_{\Omega} |\nabla u|^2 dx \right)^{1/2}$$

is a norm equivalent to the standard one in  $W_0^{1,p}(\Omega)$ .

Let  $C^{m,\beta}(\Omega)$  denote the set of all functions  $u$  belonging to  $C^m(\Omega)$  and whose partial derivatives  $D^\alpha u$ , with  $|\alpha| = m$ , are Hölder continuous with exponent  $0 < \beta < 1$ .

For a bounded domain  $\Omega$ ,  $C^{m,\beta}(\Omega)$  becomes a Banach space with a norm

$$\|u\|_{C^{m,\beta}} = \sum_{|\alpha| \leq m} \|D^\alpha u\|_\infty + \sum_{|\alpha|=m} \sup_{x \neq y \in \Omega} \frac{|D^\alpha u(x) - D^\alpha u(y)|}{|x - y|^\beta}.$$

For an arbitrary domain  $\Omega \subset \mathbb{R}^N$  and  $u \in W_0^{k,p}(\Omega)$  we have

$$K \|u\|_{p^*}^p \leq \sum_{|\alpha|=k} \int_{\Omega} |D^\alpha u(x)|^p dx, \quad (\text{B.1})$$

where  $p^* = \frac{Np}{N-kp}$  and  $K = K(k, n, p)$  is the best Sobolev constant.

This inequality says that  $W_0^{k,p}$  is continuously embedded in  $L^{p^*}(\Omega)$ .

If  $u \in W^{k,p}(\mathbb{R}^N)$ , then the best Sobolev constant  $S$  is given by

$$K(k, n, p) = \pi^{-1/2} N^{-1/p} \left( \frac{p-1}{N-p} \right)^{(p-1)/p} \left\{ \frac{\Gamma\left(1 + \frac{N}{2}\right) \Gamma(N)}{\Gamma\left(\frac{N}{p}\right) \Gamma\left(1 + N - \frac{N}{p}\right)} \right\}^{1/N},$$

where  $\Gamma(\cdot)$  denotes Euler's Gamma function. Moreover, the equality holds in (B.1) if and only if  $u$  has the form

$$u(x) = \left[ a + b|x|^{\frac{p}{p-1}} \right]^{1 - \frac{N}{p}}, \quad (\text{B.2})$$

where  $a$  and  $b$  are positive constants.

If  $k = 1, p = 2$ , then the sharp constant will be denoted by  $K_n$  and we have

$$K_n = \sqrt{\frac{4}{n(n-2)\omega_n^{\frac{n}{2}}}}, \quad (\text{B.3})$$

where  $\omega_n$  is the volume of the unit sphere.

We denote by  $D^{k,p}(\mathbb{R}^N)$  completion of  $C_0^\infty(\mathbb{R}^N)$  with respect to the norm

$$\|u\|_{D^{k,p}}^p = \sum_{|\alpha|=k} \int_{\Omega} |D^\alpha u(x)|^p dx.$$

According to the inequality (B.1), the function space  $D^{k,p}(\mathbb{R}^N)$  is continuously embedded in  $L^{p^*}(\mathbb{R}^N)$ .

Let  $\Omega$  be an unbounded domain in  $\mathbb{R}^N$  with smooth boundary  $\partial\Omega$  and let  $G$  be a subgroup of  $O(N)$  whose elements leave  $\Omega$  invariant, that is,  $g(\Omega) = \Omega$  for all  $g \in G$ . We assume that  $\Omega$  is compatible with  $G$ , that is, for some  $r > 0$

$$m(y, r, G) \rightarrow \infty, \quad \text{as } \text{dist}(y, \Omega) \leq r, \quad |y| \rightarrow \infty \quad (\text{B.4})$$

where

$$m(y, r, G) = \sup \left\{ n \in \mathbb{N} : \begin{array}{l} \exists g_1, g_2, \dots, g_n \in G \text{ s.t.} \\ B(g_j y, r) \cap B(g_k y, r) = \emptyset \text{ if } j \neq k \end{array} \right\}.$$

Let  $X = W_0^{1,p}(\Omega)$  and define a representation of  $G$  over  $X$  as follows:

$$(\pi(g)u)(x) = u(g^{-1}x), \quad g \in G, \quad u \in X \quad x \in \Omega.$$

As usual we shall write  $gu$  in place of  $\pi(g)u$ .

A function  $u$  defined on  $\Omega$  is said to be  $G$ -invariant if

$$u(gx) = u(x), \quad \forall g \in G, \text{ a.e. } x \in \Omega.$$

Then  $u \in X$  is  $G$ -invariant if and only if

$$u \in X^G = W_{0,G}^{1,p}(\Omega) := \{u \in X : gu = u, \forall g \in G\}.$$

On  $X$  we take the norm

$$\|u\| = \left\{ \int_{\Omega} (|\nabla u|^p + |u|^p) \right\}^{1/p}.$$

The next result is essentially given by Willem [297] and in this form was proved by Kobayashi-Ôtani [160].

**Theorem B.5** *If  $\Omega$  is compatible with  $G$ , then the embeddings*

$$X^G = W_{0,G}^{1,p}(\Omega) \hookrightarrow L^q(\Omega), \quad p < q < p^* = \frac{Np}{N-p}$$

*are compact.*

Now, we consider the space

$$H^1(\mathbb{R}^N) := \{ u \in L^2(\mathbb{R}^N) : \nabla u \in L^2(\mathbb{R}^N) \}$$

with the inner product

$$\langle u, v \rangle = \int_{\mathbb{R}^N} [\nabla u \nabla v + uv].$$

The space  $(H^1(\mathbb{R}^N), \langle \cdot, \cdot \rangle)$  becomes a Hilbert space.

Let  $G$  be a subgroup of  $O(N)$ . We define

$$H_G^1(\mathbb{R}^N) = \{u \in H^1 : gu = u, \forall g \in G\},$$

where  $gu(x) := u(g^{-1}x)$ .

The following result establishes a compact embedding in the case of lack of compactness.

**Theorem B.6** (Lions [193]) *Let  $N_j \geq 2, j = 1, \dots, k, \sum_{j=1}^k N_j = N$  and*

$$G := O(N_1) \times O(N_2) \times \dots \times O(N_k).$$

*Then the following embedding is compact:*

$$H_G^1(\mathbb{R}^N) \hookrightarrow L^p(\mathbb{R}^N), \quad 2 < p < 2^*.$$

In the following we consider  $\Omega = \tilde{\Omega} \times \mathbb{R}^N, N - m \geq 2, \tilde{\Omega} \subset \mathbb{R}^m (m \geq 1)$  is open bounded and  $1 \leq p \leq N$ . We consider the space  $W_0^{1,p}(\Omega)$  with the norm  $\|u\| = \left( \int_{\Omega} |\nabla u|^p \right)^{\frac{1}{p}}$ . Let  $G$  be a subgroup of  $O(N)$  defined by  $G = id^m \times O(N - m)$ . The action of  $G$  on  $W_0^{1,p}(\Omega)$  is given by  $gu(x_1, x_2) = u(x_1, g_1 x_2)$  for every  $(x_1, x_2) \in \tilde{\Omega} \times \mathbb{R}^{N-m}$  and  $g = id^m \times g_1 \in G$ . The subspace of invariant functions is defined by

$$W_{0,G}^{1,p} = \{u \in X : gu = u, \forall g \in G\}.$$

The action of  $G$  on  $W_0^{1,p}(\Omega)$  is isometric, that is

$$\|gu\| = \|u\|, \quad \forall g \in G.$$

We have the following result.

**Theorem B.7** (Lions [193]) *If  $2 \leq p \leq N$ , then the embedding*

$$W_{0,G}^{1,p} \hookrightarrow L^s(\Omega), \quad p < s < p^*$$

*is compact.*

We end this section with a result given by Esteban and Lions [110]. For this we consider  $\tilde{\Omega} \subset \mathbb{R}^m$  a bounded open set  $\Omega = \tilde{\Omega} \times \mathbb{R}$  and let

$$H_0^1(\Omega) = \{u \in L^2(\Omega) : \nabla u \in L^2(\Omega), u = 0 \text{ on } \partial\Omega = \partial\tilde{\Omega} \times \mathbb{R}\}.$$

We denote by  $K$  the cone of  $H_0^1(\Omega)$  defined by

$$\begin{aligned} \mathcal{K} = \{u \in H_0^1(\omega \times \mathbb{R}) : & u \text{ is nonnegative,} \\ & y \mapsto u(x, y) \text{ is nonincreasing for } x \in \omega, y \geq 0, \text{ and} \\ & y \mapsto u(x, y) \text{ is nondecreasing for } x \in \omega, y \leq 0\}, \end{aligned}$$

**Theorem B.8** (Esteban and Lions [110]) *The embeddings  $K \hookrightarrow L^q(\Omega)$  are compact for  $q \in (2, 2_m^*)$ , where  $2_m^* = \frac{2(m+1)}{m-1}$  if  $m > 1$ , and  $2_m^* = \infty$  if  $m = 1$ .*

### B.4 Sobolev spaces on Riemann manifolds

Let  $(M, g)$  be a Riemannian manifold of dimension  $n$ . For  $k$  an integer and  $k$  an integer and  $u \in C^\infty(M)$ ,  $\nabla^k u$  denotes the  $k$ -th covariant derivative of  $u$  (with the convection  $\nabla^0 u = u$ .) The component of  $\nabla u$  in the local coordinates  $(x^1, \dots, x^n)$  are given by

$$(\nabla^2)_{ij} = \frac{\partial^2 u}{\partial x^i \partial x^j} - \Gamma_{ij}^k \frac{\partial u}{\partial x^k}.$$

By definition on has

$$|\nabla^k u|^2 = g^{i_1 j_1} \dots g^{i_k j_k} (\nabla^k u)_{i_1 \dots i_k} (\nabla^k u)_{j_1 \dots j_k}.$$

For  $k \in \mathbb{N}$  and  $p \geq 1$  real, we denote by  $\mathcal{C}_k^p(M)$  the space of smooth functions  $u \in C^\infty(M)$  such that  $|\nabla^j u| \in L^p(M)$  for any  $j = 0, \dots, k$ . Hence,

$$\mathcal{C}_k^p = \left\{ u \in C^\infty(M) : \forall j = 0, \dots, k, \int_M |\nabla^j u|^p dv(g) < \infty \right\}$$

where, in local coordinates,  $dv(g) = \sqrt{\det(g_{ij})} dx$ , and where  $dx$  stands for the Lebesgue's volume element of  $\mathbb{R}^n$ . If  $M$  is compact, on has that  $\mathcal{C}_k^p(M) = C^\infty(M)$  for all  $k$  and  $p \geq 1$ .

**Definition B.1** The Sobolev space  $H_k^p(M)$  is the completion of  $\mathcal{C}_k^p(M)$  with respect the norm

$$\|u\|_{H_k^p} = \sum_{j=0}^k \left( \int_M |\nabla^j u|^p dv(g) \right)^{\frac{1}{p}}.$$

More precisely, one can look at  $H_k^p(M)$  as the space of functions  $u \in L^p(M)$  which are limit in  $L^p(M)$  of a Cauchy sequence  $(u_m) \subset \mathcal{C}_k$ , and define the norm  $\|u\|_{H_k^p}$  as above where  $|\nabla^j u|, 0 \leq j \leq k$ , is now the limit in  $L^p(M)$  of  $|\nabla^j u_m|$ . These space are Banach space, and if  $p > 1$ , then  $H_k^p$  is reflexive. We note that, if  $M$  is compact,  $H_k^p(M)$  does not depend on the Riemannian metric. If  $p = 2$ ,  $H_k^2(M)$  is a Hilbert space when equipped with the equivalent norm

$$\|u\| = \sqrt{\sum_{j=0}^k \int_M |\nabla^j u|^2 dv(g)}. \tag{B.5}$$

The scalar product  $\langle \cdot, \cdot \rangle$  associated to  $\| \cdot \|$  is defined by

$$\langle u, v \rangle = \sum_{m=0}^k \int_M (g^{i_1 j_1} \dots g^{i_m j_m} (\nabla^m u)_{i_1 \dots i_m} (\nabla^m v)_{j_1 \dots j_m}) dv(g). \quad (\text{B.6})$$

We denote by  $C^k(M)$  the set of  $k$  times continuously differentiable functions, for which the norm

$$\|u\|_{C^k} = \sum_{i=1}^n \sup_M |\nabla^i u|$$

is finite. The Hölder space  $C^{k,\alpha}(M)$  is defined for  $0 < \alpha < 1$  as the set of  $u \in C^k(M)$  for which the norm

$$\|u\|_{C^{k,\alpha}} = \|u\|_{C^k} + \sup_{x,y} \frac{|\nabla^k u(x) - \nabla^k u(y)|}{|x - y|^\alpha}$$

is finite, where the supremum is over all  $x \neq y$  such that  $y$  is contained in a normal neighborhood of  $x$ , and  $\nabla^k u(y)$  is taken to mean the tensor at  $x$  obtained by parallel transport along the radial geodesics from  $x$  to  $y$ .

As usual,  $C^\infty(M)$  and  $C_0^\infty(M)$  denote the spaces of smooth functions and smooth compactly supported function on  $M$  respectively.

**Definition B.2** The Sobolev space  $\mathring{H}_k^p(M)$  is the closure of  $C_0^\infty(M)$  in  $H_k^p(M)$ .

If  $(M, g)$  is a complete Riemannian manifold, then for any  $p \geq 1$ , we have  $\mathring{H}_k^p(M) = H_k^p(M)$ .

We finish this section with the Sobolev embedding theorem and the Rellich–Kondrachov result for compact manifold without and with boundary.

**Theorem B.9** (*Sobolev embedding theorems for compact manifolds*)  
Let  $M$  be a compact Riemannian manifold of dimension  $n$ .

- If  $\frac{1}{r} \geq \frac{1}{p} - \frac{k}{n}$ , then the embedding  $H_k^p(M) \hookrightarrow L^r(M)$  is continuous.
- (Rellich–Kondrachov theorem) Suppose that the inequality in a) is strict, then the embedding  $H_k^p(M) \hookrightarrow L^r(M)$  is compact.
- Suppose  $0 < \alpha < 1$  and  $\frac{1}{p} \leq \frac{k-\alpha}{n}$ , then the embedding  $H_k^p(M) \hookrightarrow C^\alpha(M)$  is continuous.

**Theorem B.10** Let  $(M, g)$  be a compact  $n$ -dimensional Riemannian manifold with boundary  $\partial M$ .

- a) The embedding  $H_1^p(M) \hookrightarrow L^q(M)$  is continuous, if  $p \leq q \leq \frac{np}{n-p}$  and compact for  $p \leq q < \frac{np}{n-p}$ .
- b) If  $\partial M \neq \emptyset$ , then the embedding  $H_1^p(M) \hookrightarrow L^q(\partial M)$  is continuous, if  $p \leq q \leq \frac{p(n-1)}{n-p}$  and compact for  $p \leq q < \frac{p(n-1)}{n-p}$ .

**Theorem B.11** For any smooth compact Riemannian manifold  $(M, g)$  of dimension  $n \geq 3$ , there exists  $B > 0$  such that for any  $u \in H_1^2(M)$ ,

$$\left( \int_M |u|^{2^*} dv_g \right)^{2/2^*} \leq K_n^2 \int_M |\nabla u|^2 dv_g + B \int_M u^2 dv_g \quad (\text{B.7})$$

and the inequality is sharp.

**Theorem B.12** (Global elliptic regularity) Let  $M$  be a compact Riemann manifold, and suppose that  $u \in L_{\text{loc}}^1(M)$  is a weak solution to  $\Delta_g u = f$ .

- a) If  $f \in H_k^p(M)$ , then  $u \in H_{k+2}^p(M)$ , and

$$\|u\|_{H_{k+2}^p} \leq C(\|\Delta_g u\|_{H_k^p} + \|u\|_{L^p}).$$

- b) If  $f \in C^{k,\alpha}(M)$ , then  $f \in C^{k+2,\alpha}(M)$ , and

$$\|u\|_{C_{k+2}^p} \leq C(\|\Delta_g u\|_{C_k^p} + \|u\|_{C^\alpha}).$$

We conclude this part by stating the following version of the Hopf's maximum principle.

**Proposition B.1** Let  $f : M \times \mathbb{R} \rightarrow \mathbb{R}$  be a continuous function and  $u \in C^2(M)$  such that

$$-\Delta_g u \geq u(x)f(x, u(x)),$$

then either  $u > 0$  or  $u = 0$ .

# C

## Category and genus

Every human activity, good or bad, except mathematics, must come to an end.

---

Paul Erdős (1913–1996)

Topological tools play a central role in the study of variational problems. Though this approach was foreshadowed in the works of H. Poincaré and G. Birkhoff, the force of these ideas was realized in the first decades of the 20th century, in the pioneering works of Ljusternik and Schnirelmann [197] and Morse [220, 221]. In this section we recall the notions of Ljusternik-Schnirelmann category and Krasnoselski genus as well as some basic properties of them.

**Definition C.1** Let  $M$  be a topological space and  $A \subset M$  a subset. The continuous map  $\eta : A \times [0, 1] \rightarrow M$  is called a deformation of  $A$  in  $M$  if  $\eta(u, 0) = u$  for every  $u \in A$ . The set  $A$  is said to be contractible in  $M$  if there exists a deformation  $\eta : A \times [0, 1] \rightarrow M$  with  $\eta(A, 1) = \{p\}$  for some  $p \in M$ .

**Definition C.2** Let  $M$  be a topological space. A set  $A \subset M$  is said to be of Ljusternik-Schnirelmann category  $k$  in  $M$  (denoted  $\text{cat}_M(A) = k$ ) if it can be covered by  $k$  but not by  $k-1$  closed sets which are contractible to a point in  $M$ . If such  $k$  does not exist, then  $\text{cat}_M(A) = +\infty$ . We define  $\text{cat}(M) = \text{cat}_M(M)$ .

The Ljusternik-Schnirelmann category has the following basic properties.

**Proposition C.1** *If  $A$  and  $B$  are subsets of the topological space  $M$ , then the following properties hold true.*

- (a) If  $A \subseteq B$ , then  $\text{cat}_M(A) \leq \text{cat}_M(B)$ ;
- (b)  $\text{cat}_M(A \cup B) \leq \text{cat}_M(A) + \text{cat}_M(B)$ .
- (c) If  $A$  is closed in  $X$  and  $\eta : A \times [0, 1] \rightarrow M$  a deformation of  $A$  in  $X$ , then  $\text{cat}_M(A) \leq \text{cat}_M(\eta(A, 1))$ .
- (d) If  $M$  is an ANR (in particular, it is a Banach-Finsler manifold of class  $C^1$ ) and  $A \subset M$ , then there exist a neighborhood of  $A$  in  $M$ , such that  $\text{cat}_M(\bar{U}) = \text{cat}_M(A)$ .

**Theorem C.1** [111] *Let  $M$  be a non-contractible topological space and let  $\Omega(M)$  denote the space of based loops in  $M$ . Then  $\text{cat}(\Omega(M)) = \infty$ .*

**Definition C.1** Let  $M$  be an arcwise connected topological space. Then  $\text{cuplong}(M)$  is the greatest non-vanishing iterated cup-product of singular cocycles of  $M$  such that each component is of dimension greater than zero.

**Theorem C.2** [272] *If  $M$  is an arcwise connected metric space, then  $\text{cat}(M) \geq \text{cuplong}(M) + 1$ .*

**Theorem C.3** [273] *Let  $M$  be a compact, connected, simply connected and non-contractible Riemannian manifold. If  $\Omega(M)$  denotes the space of the free loops on  $M$ , then  $\text{cuplong}(\Omega) = \infty$ .*

The word *genus* hails from biology, where it is used to designate a collection of organisms having common characteristics. In mathematics this word is also used to group objects with common characteristics. We give in what follows a rigorous mathematical description of this notion.

Let  $X$  be a real Banach space and denote

$$\mathcal{A} = \{A \subset X \setminus \{0\} : A = -A, A \text{ is closed}\}.$$

**Definition C.3** We say that the positive integer  $k$  is the Krasnoselski genus of  $A \in \mathcal{A}$ , if there exists an odd map  $\varphi \in C(A, \mathbb{R}^k \setminus \{0\})$  and  $k$  is the smallest integer with this property. The genus of the set  $A$  is denoted by  $\gamma(A) = k$ . When there does not exist a finite such  $k$ , set  $\gamma(A) = \infty$ . Finally, set  $\gamma(\emptyset) = 0$ .

**Example C.1** Suppose  $A \subset X \setminus \{0\}$  is closed and  $A \cap (-A) = \emptyset$ . Let  $\tilde{A} = A \cup (-A)$ . Then  $\gamma(\tilde{A}) = 1$  since the function  $\varphi(x) = 1$  for  $x \in A$  and  $\varphi(x) = -1$  for  $x \in -A$  is odd and lies in  $C(\tilde{A}, \mathbb{R} \setminus \{0\})$ .

For  $A \in \mathcal{A}$  and  $\delta > 0$  we denote by  $N_\delta(A)$  the uniform  $\delta$ -neighborhood of  $A$ , that is,  $N_\delta(A) = \{x \in X : \text{dist}_A(x) \leq \delta\}$ .

**Proposition C.2** *Let  $A, B \in \mathcal{A}$ . Then the following properties hold true.*

- 1° *Normalization: If  $x \neq 0$ ,  $\gamma(\{x\} \cup \{-x\}) = 1$ ;*
- 2° *Mapping property: If there exists an odd map  $f \in C(A, B)$ , then  $\gamma(A) \leq \gamma(B)$ ;*
- 3° *Monotonicity property: If  $A \subset B$ , then  $\gamma(A) \leq \gamma(B)$ ;*
- 4° *Subadditivity:  $\gamma(A \cup B) \leq \gamma(A) + \gamma(B)$ ;*
- 5° *Continuity property: If  $A$  is compact then  $\gamma(A) < \infty$  and there is  $\delta > 0$  such that  $N_\delta(A) \in \mathcal{A}$ . In such a case,  $\gamma(N_\delta(A)) = \gamma(A)$ .*

**Example C.2** (a) If  $S^{k-1}$  is the  $(k-1)$ -dimensional unit sphere in  $\mathbb{R}^k$ ,  $k \in \mathbb{N}$ , then  $\gamma(S^{k-1}) = k$ .

(b) If  $S$  is the unit sphere in an infinite dimensional and separable Banach space, then  $\gamma(S) = +\infty$ .

# D

## Clarke and Degiovanni gradients

Profound study of nature is  
the most fertile source of  
mathematical discoveries.

---

Baron Jean Baptiste Joseph  
Fourier (1768–1830)

In many time independent problems arising in applications, the solutions of a given problem are critical points of an appropriate energy functional  $f$ , which is usually supposed to be real and of class  $C^1$  (or even differentiable) on a real Banach space. One may ask what happens if  $f$ , which often is associated to the original equation in a canonical way, fails to be  $C^1$  or differentiable. In this case the gradient of  $f$  must be replaced by a generalized one, in a sense which is described in this Appendix, either for locally Lipschitz or for continuous/lower semi-continuous functionals. In the latter case we use in the general framework of *metric spaces*.

### D.1 Locally Lipschitz functionals

Let  $X$  will denote a real Banach space. Let  $X^*$  be its dual and, for every  $x \in X$  and  $x^* \in X^*$ , let  $\langle x^*, x \rangle$  denote the duality pairing between  $X^*$  and  $X$ .

**Definition D.1** A functional  $f : X \rightarrow \mathbb{R}$  is said to be locally Lipschitz provided that, for every  $x \in X$ , there exists a neighbourhood  $V$  of  $x$  and a positive constant  $k = k(V)$  depending on  $V$  such that for all  $y, z \in V$ ,

$$|f(y) - f(z)| \leq k \|y - z\|.$$

**Definition D.2** Let  $f : X \rightarrow \mathbb{R}$  be a locally Lipschitz functional and  $x, v \in X$ . We call the Clarke generalized directional derivative of  $f$  in  $x$  with respect to the direction  $v$  the number

$$f^0(x, v) = \limsup_{\substack{y \rightarrow x \\ \lambda \searrow 0}} \frac{f(y + \lambda v) - f(y)}{\lambda}.$$

If  $f$  is a locally Lipschitz functional, then  $f^0(x, v)$  is a finite number and

$$|f^0(x, v)| \leq k\|v\|.$$

Moreover, if  $x \in X$  is fixed, then the mapping  $v \mapsto f^0(x, v)$  is positive homogeneous and subadditive, so convex continuous. By the Hahn-Banach theorem, there exists a linear map  $x^* : X \rightarrow \mathbb{R}$  such that for every  $v \in X$ ,

$$x^*(v) \leq f^0(x, v).$$

**Definition D.3** Let  $f : X \rightarrow \mathbb{R}$  be a locally Lipschitz functional and  $x \in X$ . The generalized gradient (Clarke subdifferential) of  $f$  at the point  $x$  is the nonempty subset  $\partial f(x)$  of  $X^*$  which is defined by

$$\partial f(x) = \{x^* \in X^*; f^0(x, v) \geq \langle x^*, v \rangle, \text{ for all } v \in X\}.$$

We also point out that if  $f$  is convex, then  $\partial f(x)$  coincides with the subdifferential of  $f$  in  $x$  in the sense of the convex analysis, that is,

$$\partial f(x) = \{x^* \in X^*; f(y) - f(x) \geq \langle x^*, y - x \rangle, \text{ for all } y \in X\}.$$

We list in what follows the main properties of the Clarke gradient of a locally Lipschitz functional. We refer to [72], [73] for further details and proofs.

- a) For every  $x \in X$ ,  $\partial f(x)$  is a convex and  $\sigma(X^*, X)$ -compact set.
- b) For every  $x, v \in X$  the following holds

$$f^0(x, v) = \max\{\langle x^*, v \rangle; x^* \in \partial f(x)\}.$$

c) The multivalued mapping  $x \mapsto \partial f(x)$  is upper semicontinuous, in the sense that for every  $x_0 \in X$ ,  $\varepsilon > 0$  and  $v \in X$ , there exists  $\delta > 0$  such that, for any  $x^* \in \partial f(x)$  satisfying  $\|x - x_0\| < \delta$ , there is some  $x_0^* \in \partial f(x_0)$  satisfying  $|\langle x^* - x_0^*, v \rangle| < \varepsilon$ .

- d) The functional  $f^0(\cdot, \cdot)$  is upper semi-continuous.
- e) If  $x$  is an extremum point of  $f$ , then  $0 \in \partial f(x)$ .
- f) The mapping

$$\lambda(x) = \min_{x^* \in \partial f(x)} \|x^*\|$$

exists and is lower semicontinuous.

g)  $\partial(-f)(x) = -\partial f(x)$ .

h) Lebourg’s mean value theorem: if  $x$  and  $y$  are two distinct point in  $X$  then there exists a point  $z$  situated in the open segment joining  $x$  and  $y$  such that

$$f(y) - f(x) \in \langle \partial f(z), y - x \rangle$$

i) If  $f$  has a Gâteaux derivative  $f'$  which is continuous in a neighbourhood of  $x$ , then  $\partial f(x) = \{f'(x)\}$ . If  $X$  is finite dimensional, then  $\partial f(x)$  reduces at one point if and only if  $f$  is Fréchet-differentiable at  $x$ .

**Definition D.4** A point  $x \in X$  is said to be a critical point of the locally Lipschitz functional  $f : X \rightarrow \mathbb{R}$  if  $0 \in \partial f(x)$ , that is  $f^0(x, v) \geq 0$ , for every  $v \in X$ . A number  $c$  is a critical value of  $f$  provided that there exists a critical point  $x \in X$  such that  $f(x) = c$ .

We remark that a minimum point is also a critical point. Indeed, if  $x$  is a local minimum point, then for every  $v \in X$ ,

$$0 \leq \limsup_{\lambda \searrow 0} \frac{f(x + \lambda v) - f(x)}{\lambda} \leq f^0(x, v).$$

**Definition D.5** If  $f : X \rightarrow \mathbb{R}$  is a locally Lipschitz functional and  $c$  is a real number, we say that  $f$  satisfies the Palais-Smale condition at the level  $c$  (in short,  $(PS)_c$ ) if any sequence  $(x_n)$  in  $X$  satisfying  $f(x_n) \rightarrow c$  and  $\lambda(x_n) \rightarrow 0$ , contains a convergent subsequence. The mapping  $f$  satisfies the Palais-Smale condition (in short,  $(PS)$ ) if every sequence  $(x_n)$  which satisfies  $(f(x_n))$  is bounded and  $\lambda(x_n) \rightarrow 0$ , has a convergent subsequence.

## D.2 Continuous or lower semi-continuous functionals

We are now interested in the main properties of the corresponding properties in the case where  $f$  is a continuous or even a lower semi-continuous functional defined on a metric space.

**Definition D.6** Let  $(X, d)$  be a metric space and let  $f : X \rightarrow \mathbb{R}$  be a continuous function. Let  $u \in X$  be a fixed element. We denote by  $|df|(u)$  the supremum of the  $\sigma \in [0, \infty[$  such that there exist  $\delta > 0$  and a continuous map  $\mathcal{H} : B(u, \delta) \times [0, \delta] \rightarrow X$  such that for all  $v \in B(u, \delta)$  and  $t \in [0, \delta]$  we have

- a)  $d(\mathcal{H}(v, t), v) \leq t$ ;  
 b)  $f(\mathcal{H}(v, t)) \leq f(v) - \sigma t$ .

The extended real number  $|df|(u)$  is called the *weak slope* of  $f$  at  $u$ .

**Definition D.7** Let  $(X, d)$  be a metric space and  $f : X \rightarrow \mathbb{R}$  be a continuous function. A point  $u \in X$  is said to be a critical point of  $f$  if  $|df|(u) = 0$ .

**Definition D.8** Let  $(X, d)$  be a metric space. A continuous function  $f : X \rightarrow \mathbb{R}$  satisfies the Palais-Smale condition at level  $c \in \mathbb{R}$  (shortly,  $(PS)_c$ -condition) if every sequence  $\{u_n\} \subset X$  such that  $\lim_{n \rightarrow \infty} f(u_n) = c$  and  $\lim_{n \rightarrow \infty} |df|(u_n) = 0$ , possesses a convergent subsequence.

For every  $c \in \mathbb{R}$  we set

$$K_c = \{u \in X : |df|(u) = 0, f(u) = c\};$$

$$f^c = \{u \in X : f(u) \leq c\}.$$

**Remark D.1** Note that  $|df|(u) = \|f'(u)\|$  whenever  $X$  is a Finsler manifold of class  $C^1$  and  $f$  is of class  $C^1$ , see [85]. In particular, when  $X$  is a Banach space, Definition D.8 reduces to Definition 1.1 (a).

**Theorem D.1** [Deformation lemma; [75, Theorem 2.3]] *Let  $(X, d)$  be a complete metric space and let  $f : X \rightarrow \mathbb{R}$  be a continuous function, and  $c \in \mathbb{R}$ . Assume that  $f$  satisfies the  $(PS)_c$ -condition. Then, given  $\bar{\varepsilon} > 0$ ,  $\mathcal{O}$  a neighborhood of  $K_c$  (if  $K_c = \emptyset$ , we choose  $\mathcal{O} = \emptyset$ ) and  $\lambda > 0$ , there exist  $\varepsilon > 0$  and a continuous map  $\eta : X \times [0, 1] \rightarrow X$  such that*

- (i)  $d(\eta(u, t), u) \leq \lambda t$  for all  $u \in X$  and  $t \in [0, 1]$ ;  
 (ii)  $f(\eta(u, t)) \leq f(u)$  for all  $u \in X$  and  $t \in [0, 1]$ ;  
 (iii)  $\eta(u, t) = u$  for all  $u \notin f^{-1}(]c - \bar{\varepsilon}, c + \bar{\varepsilon}[)$  and  $t \in [0, 1]$ ;  
 (iv)  $\eta(f^{c+\varepsilon} \setminus \mathcal{O}, 1) \subseteq f^{c-\varepsilon}$ .

These notions can be extended to *arbitrary* functions defined on a metric space  $(X, d)$ . Let  $f : X \rightarrow \mathbb{R}$  be a function. We denote by  $B_r(u)$  the open ball of center  $u$  and radius  $r$  and we set

$$\text{epi}(f) = \{(u, \lambda) \in X \times \mathbb{R} : f(u) \leq \lambda\}.$$

In the following,  $X \times \mathbb{R}$  will be endowed with the metric

$$d((u, \lambda), (v, \mu)) = (d(u, v)^2 + (\lambda - \mu)^2)^{1/2}$$

and  $\text{epi}(f)$  with the induced metric.

**Definition D.9** For every  $u \in X$  with  $f(u) \in \mathbb{R}$ , we denote by  $|df|(u)$  the supremum of the  $\sigma$ 's in  $[0, +\infty[$  such that there exist  $\delta > 0$  and a continuous map

$$\mathcal{H} : (B_\delta(u, f(u)) \cap \text{epi}(f)) \times [0, \delta] \rightarrow X$$

satisfying

$$d(\mathcal{H}((w, \mu), t), w) \leq t, \quad f(\mathcal{H}((w, \mu), t)) \leq \mu - \sigma t,$$

whenever  $(w, \mu) \in B_\delta(u, f(u)) \cap \text{epi}(f)$  and  $t \in [0, \delta]$ .

The extended real number  $|df|(u)$  is called the weak slope of  $f$  at  $u$ .

Define a function  $\mathcal{G}_f : \text{epi}(f) \rightarrow \mathbb{R}$  by  $\mathcal{G}_f(u, \lambda) = \lambda$ . Of course,  $\mathcal{G}_f$  is Lipschitz continuous of constant 1.

**Proposition D.1** For every  $u \in X$  with  $f(u) \in \mathbb{R}$ , we have  $f(u) = \mathcal{G}_f(u, f(u))$  and

$$|df|(u) = \begin{cases} \frac{|d\mathcal{G}_f|(u, f(u))}{\sqrt{1 - |d\mathcal{G}_f|(u, f(u))^2}} & \text{if } |d\mathcal{G}_f|(u, f(u)) < 1, \\ +\infty & \text{if } |d\mathcal{G}_f|(u, f(u)) = 1. \end{cases}$$

We refer to [59, Proposition 2.3] for a complete proof of this result.

The previous proposition allows us to reduce, at some extent, the study of the general function  $f$  to that of the continuous function  $\mathcal{G}_f$ . Moreover, if  $f$  is continuous then Definition D.9 reduces to Definition D.6.

Two important notions which are very well related to the notion of weak slope are those of *generalized directional derivative* and *generalized gradient* in the sense of Degiovanni. For every  $u \in X$  with  $f(u) \in \mathbb{R}$ ,  $v \in X$  and  $\varepsilon > 0$ , let  $f_\varepsilon^0(u; v)$  be the infimum of  $r \in \mathbb{R}$  such that there are  $\delta > 0$  and a continuous map

$$\mathcal{V} : (B_\delta(u, f(u)) \cap \text{epi}(f)) \times (0, \delta] \rightarrow B_\varepsilon(v)$$

satisfying

$$f(w + t\mathcal{V}((w, \mu), t)) \leq \mu + rt,$$

whenever  $(w, \mu) \in B_\delta(u, f(u)) \cap \text{epi}(f)$  and  $t \in (0, \delta]$ . Under these assumptions, we define the *generalized directional derivative* of  $f$  at  $u$  with respect to  $v$  by

$$f^0(u; v) := \sup_{\varepsilon > 0} f_\varepsilon^0(u; v).$$

We also define the *generalized gradient* of  $f$  at  $u$  by

$$\partial f(u) := \{\varphi \in X^*; f^0(u; v) \geq \varphi(v) \text{ for every } v \in X\}.$$

For a function of one variable,  $\partial f(x_0)$  is the collection of angular coefficients  $m$  for which the lines  $f(x_0) + m(x - x_0)$  lie under the graph of  $f$ .

**Theorem D.2** *If  $u \in X$  and  $f(u) \in \mathbb{R}$ , the following facts hold:*

- (a)  $|df|(u) < +\infty \iff \partial f(u) \neq \emptyset$ ;
- (b)  $|df|(u) < +\infty \implies \partial f(u) \geq \min \{\|u^*\| : u^* \in \partial f(u)\}$ .

We point out that  $f^0(u; v)$  is greater or equal than the directional derivative in the sense of Clarke-Rockafellar, hence  $\partial f(u)$  contains the subdifferential of  $f$  at  $u$  in the sense of Clarke. However, if  $f : X \rightarrow \mathbb{R}$  is locally Lipschitz, these notions agree with those of Clarke. Thus, in such a case,  $f^0(u; \cdot)$  is also Lipschitz continuous and we have for all  $u, v \in X$ ,

$$f^0(u; v) = \limsup_{\substack{z \rightarrow u, w \rightarrow v \\ t \rightarrow 0^+}} \frac{f(z + tw) - f(z)}{t}, \quad (\text{D.1})$$

$$\{(u, v) \mapsto f^0(u; v)\} \text{ is upper semicontinuous on } X \times X. \quad (\text{D.2})$$

By means of the weak slope, we can now introduce the two main notions of critical point theory.

**Definition D.10** We say that  $u \in X$  is a (lower) critical point of  $f$ , if  $f(u) \in \mathbb{R}$  and  $|df|(u) = 0$ . We say that  $c \in \mathbb{R}$  is a (lower) critical value of  $f$ , if there exists a (lower) critical point  $u \in X$  of  $f$  with  $f(u) = c$ .

**Definition D.11** Let  $c \in \mathbb{R}$ . A sequence  $(u_n)$  in  $X$  is said to be a Palais-Smale sequence at level  $c$  ( $(PS)_c$ -sequence, for short) for  $f$ , if  $f(u_n) \rightarrow c$  and  $|df|(u_n) \rightarrow 0$ .

We say that  $f$  satisfies the Palais-Smale condition at level  $c$  ( $(PS)_c$ , for short), if every  $(PS)_c$ -sequence  $(u_n)$  for  $f$  admits a convergent subsequence  $(u_{n_k})$  in  $X$ .

The main feature of the weak slope is that it allows to prove natural extensions of the classical critical point theory for general continuous functions defined on complete metric spaces. Moreover, one can try to reduce the study of a lower semicontinuous function  $f$  to that of the continuous function  $\mathcal{G}_f$ . Actually, Proposition D.1 suggests to exploit

the bijective correspondence between the set where  $f$  is finite and the graph of  $f$ .

Assuming that  $(X, d)$  is a metric space and  $f : X \rightarrow \mathbb{R}$  is continuous, for any  $a \in \overline{\mathbb{R}} := \mathbb{R} \cup \{-\infty, +\infty\}$  we set

$$f^a := \{u \in X; f(u) \leq a\}.$$

**Definition D.12** Let  $a, b \in \overline{\mathbb{R}}$  with  $a \leq b$ . The pair  $(f^b, f^a)$  is said to be trivial if for every neighborhoods  $[\alpha', \alpha'']$  of  $a$  and  $[\beta', \beta'']$  of  $b$  in  $\overline{\mathbb{R}}$ , there exists a continuous function  $\mathcal{H} : f^{\beta'} \times [0, 1] \rightarrow f^{\beta''}$  such that for all  $x \in f^{\beta'}$ ,  $\mathcal{H}(x, 0) = x$ ,  $\mathcal{H}(f^{\beta'} \times \{1\}) \subset f^{\alpha''}$ , and  $\mathcal{H}(f^{\alpha'} \times [0, 1]) \subset f^{\alpha''}$ .

**Definition D.13** A real number  $c$  is said to be an essential value of  $f$  if for every  $\varepsilon > 0$  there exist  $a, b \in (c - \varepsilon, c + \varepsilon)$  with  $a < b$  such that the pair  $(f^b, f^a)$  is not trivial.

**Example D.1** Let  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  defined by  $f(x, y) = e^x - y^2$ . Then 0 is an essential value of  $f$ , but not a critical value of  $f$ . On the other hand, condition  $(PS)_0$  is not satisfied for  $f$

**Definition D.14** Let  $X$  be a normed space and  $f : X \rightarrow \overline{\mathbb{R}}$  an even function with  $f(0) < +\infty$ . For every  $(0, \lambda) \in \text{epi}(f)$  we denote by  $|d_{\mathbb{Z}_2} \mathcal{G}_f|(0, \lambda)$  the supremum of the  $\sigma$ 's in  $[0, +\infty)$  such that there exist  $\delta > 0$  and a continuous map

$$\mathcal{H} = (\mathcal{H}_1, \mathcal{H}_2) : (B_\delta(0, \lambda) \cap \text{epi}(f)) \times [0, \delta] \rightarrow \text{epi}(f)$$

satisfying

$$d(\mathcal{H}((w, \mu), t), (w, \mu)) \leq t, \quad \mathcal{H}_2((w, \mu), t) \leq \mu - \sigma t,$$

$$\mathcal{H}_1((-w, \mu), t) = -\mathcal{H}_1((w, \mu), t),$$

whenever  $(w, \mu) \in B_\delta(0, \lambda) \cap \text{epi}(f)$  and  $t \in [0, \delta]$ .

The following result is the *saddle point* theorem for nondifferentiable functions.

**Theorem D.3** Let  $X$  be a Banach space and assume that  $f : X \rightarrow \mathbb{R} \cup \{+\infty\}$  is an even lower semicontinuous function. Assume that there exists a strictly increasing sequence  $(V_h)$  of finite-dimensional subspaces of  $X$  with the following properties:

- (a) there exist a closed subspace  $Z$  of  $X$ ,  $\varrho > 0$  and  $\alpha > f(0)$  such that  $X = V_0 \oplus Z$  and

$$\forall u \in Z : \|u\| = \varrho \implies f(u) \geq \alpha;$$

- (b) there exists a sequence  $(R_h)$  in  $]\varrho, +\infty[$  such that

$$\forall u \in V_h : \|u\| \geq R_h \implies f(u) \leq f(0);$$

- (c) for every  $c \geq \alpha$ , the function  $f$  satisfies  $(PS)_c$  and  $(epi)_c$ ;

- (d) we have  $|d_{\mathbb{Z}_2} \mathcal{G}_f|(0, \lambda) \neq 0$  whenever  $\lambda \geq \alpha$ .

Then there exists a sequence  $(u_h)$  of critical points of  $f$  with  $f(u_h) \rightarrow +\infty$ .

# E

## Elements of set-valued analysis

To myself I am only a child  
playing on the beach, while  
vast oceans of truth lie  
undiscovered before me.

---

Sir Isaac Newton (1642–1727)

Let  $X$  and  $Y$  be metric spaces, and let  $F : X \rightsquigarrow Y$  be a set-valued map with nonempty values.

**Definition E.1** (i)  $F : X \rightsquigarrow Y$  is upper semi-continuous at  $u \in X$  (usc at  $u$ ) if for any neighborhood  $U$  of  $F(u)$  there exists  $\eta > 0$  such that for every  $u' \in B_X(u, \eta)$  we have  $F(u') \subseteq U$ .

(ii)  $F : X \rightsquigarrow Y$  is lower semi-continuous at  $u \in X$  (lsc at  $u$ ) if for any  $y \in F(u)$  and for any sequence of elements  $\{u_n\}$  in  $X$  converging to  $u$  there exists a sequence  $\{y_n\}$  converging to  $y$  and  $y_n \in F(u_n)$ .

(iii)  $F : X \rightsquigarrow Y$  is upper (resp. lower) semi-continuous on  $X$  if  $F$  is upper (resp. lower) semi-continuous at every point  $u \in X$ .

(iv)  $F : X \rightsquigarrow Y$  is continuous at  $u \in X$  if it is both upper semi-continuous and lower semi-continuous at  $u$ , and that it is continuous on  $X$  if and only if it is continuous at every point of  $X$ .

Let  $M$  be a subset of  $Y$ . We set

$$F^{-1}(M) = \{x \in X : F(x) \cap M \neq \emptyset\};$$

$$F^{+1}(M) = \{x \in X : F(x) \subseteq M\}.$$

The subset  $F^{-1}(M)$  is called the *inverse image* of  $M$  by  $F$  and  $F^{+1}(M)$  is called the *core* of  $M$  by  $F$ . The following characterization of the upper (resp. lower) semicontinuity of  $F$  on  $X$  can be given.

**Proposition E.1** *Let  $X$  and  $Y$  be metric spaces. A set-valued map  $F : X \rightsquigarrow Y$  with nonempty values is upper semicontinuous on  $X$  if and only if the inverse image of any closed subset  $M \subseteq Y$  is closed, and it is lower semicontinuous on  $X$  if and only if the core of any closed subset  $M \subseteq Y$  is closed.*

Let  $X$  and  $Y$  be vector spaces, let  $F : X \rightsquigarrow Y$  be a set-valued map, and let  $K$  be a convex subset of  $X$ .

**Definition E.2** (i)  $F$  is a process if  $\lambda F(u) = F(\lambda u)$  for all  $u \in X$ ,  $\lambda > 0$ , and  $0 \in F(0)$ .

(ii) We say that  $F$  is convex on  $K$  if

$$\lambda F(x_1) + (1 - \lambda)F(x_2) \subseteq F(\lambda x_1 + (1 - \lambda)x_2)$$

for all  $x_1, x_2 \in K$  and all  $\lambda \in [0, 1]$ .

Let  $X$  be a normed space,  $K$  be a subset of  $X$ , and let  $u \in \overline{K}$ , where  $\overline{K}$  is the closure of  $K$ . The *contingent cone*  $T_K(u)$  is defined by

$$T_K(u) = \{v \in X : \liminf_{t \rightarrow 0^+} \frac{\text{dist}(u + tv, K)}{t} = 0\}.$$

In particular, if  $K$  is convex, then  $K \subseteq u + T_K(u)$ .

Let  $F : X \rightsquigarrow \mathbb{R}$  be a set-valued map with nonempty compact values.

**Definition E.3** (i) We say that  $F$  is Lipschitz around  $u \in X$  if there exist a positive constant  $L$  and a neighborhood  $U$  of  $u$  such that

$$F(u_1) \subseteq F(u_2) + L\|u_1 - u_2\| \cdot [-1, 1] \text{ for all } u_1, u_2 \in U.$$

(ii) Let  $K \subseteq X$ .  $F$  is  $K$ -locally Lipschitz if it is Lipschitz around all  $u \in K$ .

**Proposition E.2** *Let  $X$  be a normed space. If  $F : X \rightsquigarrow \mathbb{R}$  is  $K$ -locally Lipschitz, then the restriction  $F|_K : K \rightsquigarrow \mathbb{R}$  is continuous on  $K$ .*

**Definition E.4** The contingent derivative  $DF(u, c)$  of  $F : X \rightsquigarrow \mathbb{R}$  at  $(u, c) \in \text{Graph}F$  is the set-valued map from  $X$  to  $\mathbb{R}$  defined by

$$\text{Graph}(DF(u, c)) = T_{\text{Graph}F}(u, c), \quad (\text{E.1})$$

where  $T_{\text{Graph}F}(u, c)$  is the contingent cone at  $(u, c)$  to the  $\text{Graph}F$ .

We can characterize the contingent derivative by a limit of a differential quotient. Let  $(u, c) \in \text{Graph}F$  and suppose that  $F$  is Lipschitz

around  $u$ . We have

$$v \in DF(u, c)(h) \iff \liminf_{t \rightarrow 0^+} \text{dist}(v, \frac{F(u + th) - c}{t}) = 0 \quad (\text{E.2})$$

(see Aubin and Frankowska [17, Proposition 5.1.4, p. 186]). If we introduce the *Kuratowski upper limit*

$$\text{Limsup}_{u' \rightarrow u} F(u') = \left\{ c \in \mathbb{R} : \liminf_{u' \rightarrow u} \text{dist}(c, F(u')) = 0 \right\},$$

then (E.2) can be written in the following form:

$$DF(u, c)(h) = \text{Limsup}_{t \rightarrow 0^+} \frac{F(u + th) - c}{t}.$$

**Remark E.1** Let us consider the case when  $F$  is single-valued, that is,  $F(u) = \{f(u)\}$  for all  $u \in X$ . Suppose that  $f : X \rightarrow \mathbb{R}$  is continuously differentiable. From Aubin and Frankowska [17, Proposition 5.1.3, p. 184], we have that

$$DF(u, f(u))(h) = f'(u)(h) \text{ for all } h \in X. \quad (\text{E.3})$$

**Definition E.5** (i) We say that  $F : X \rightsquigarrow \mathbb{R}$  is sleek at  $(u, c) \in \text{Graph } F$  if the map

$$\text{Graph } F \ni (u', c') \rightsquigarrow \text{Graph}(DF(u', c'))$$

is lower semicontinuous at  $(u, c)$ .  $F$  is said to be sleek if it is sleek at every point  $(u, c) \in \text{Graph } F$ .

(ii)  $F : X \rightsquigarrow \mathbb{R}$  is lower semicontinuously differentiable if the map

$$(u, c, h) \in \text{Graph } F \times X \rightsquigarrow DF(u, c)(h)$$

is lower semicontinuous.

The lower semicontinuous differentiability of a set-valued map clearly implies its sleekness.

**Remark E.2** If  $F : X \rightsquigarrow \mathbb{R}$  is a set-valued map with nonempty compact values and closed graph, and it is sleek at  $(u, c) \in \text{Graph } F$ , then its contingent derivative at  $(u, c)$  is a closed convex process (see [17, Theorem 4.1.8, p. 130]).

## References

- [1] Abbondandolo, A. and Figalli, A. (2007). High action orbits for Tonelli Lagrangians and superlinear Hamiltonians on compact configuration spaces, *J. Differential Equations* **234**, 626–653.
- [2] Akagi, G., Kobayashi, J., and Ôtani, M. (2003). Principle of symmetric criticality and evolution equations, Dynamical systems and differential equations (Wilmington, NC, 2002), *Discrete Contin. Dyn. Syst.*, suppl., 1–10.
- [3] Alama, S. and Tarantello, G. (1993). On semilinear elliptic equations with indefinite nonlinearities, *Calc. Var. Partial Differential Equations* **1**, 439–475.
- [4] Alama, S. and Tarantello, G. (1996). Elliptic problems with nonlinearities indefinite in sign, *J. Funct. Anal.* **141**, 159–215.
- [5] Allegretto, W. and Huang, Y.X. (1995). Eigenvalues of the indefinite weight  $p$ -Laplacian in weighted  $\mathbb{R}^N$  spaces, *Funkc. Ekvac.* **38**, 233–242.
- [6] Ambrosetti, A., Brézis, H., and Cerami, G. (1994). Combined effects of concave and convex nonlinearities in some elliptic problems, *J. Funct. Anal.* **122**, 519–543.
- [7] Ambrosetti, A. and Malchiodi, A. (2006). *Perturbation Methods and Semilinear Elliptic Problems on  $\mathbb{R}^n$*  (Birkhäuser Verlag, Basel).
- [8] Ambrosetti, A. and Rabinowitz, P. (1973). Dual variational methods in critical point theory and applications, *J. Funct. Anal.* **14**, 349–381.
- [9] Amick, C.J. (1984). Semilinear elliptic eigenvalue problems on an infinite strip with an application to stratified fluids, *Ann. Scuola Norm. Sup. Pisa Cl. Sci.* **11**, 441–499.
- [10] Amick, C.J. and Toland, J.F. (1983). Nonlinear elliptic eigenvalue problems on an infinite strip-global theory of bifurcation and asymptotic bifurcation, *Math. Ann.* **262**, 313–342.
- [11] Ammann, B., Dahl, M., and Humbert E. (2008). Smooth Yamabe invariant and surgery, *arXiv:0804.1418v3*.
- [12] Anane, A. (1987). Simplicité et isolation de la première valeur propre du  $p$ -laplacien, *C.R. Acad. Sci. Paris, Sér. I Math.* **305**, 725–728.
- [13] Anello, G. and Cordaro, G. (2007). Perturbation from Dirichlet problem involving oscillating nonlinearities, *J. Differential Equations* **234**, 80–90.

- [14] Aronsson, G. and Janfalk, U. (1992). On Hele-Shaw flow of power-law fluids, *European J. Appl. Math.* **3**, 343–366.
- [15] Asplund, E. (1967). Averaged norms, *Israel J. Math.* **5**, 227–233.
- [16] Asplund, E. (1968). Topics in the theory of convex functions, *Proc. NATO Institute Venice*, Oderisi Gubio.
- [17] Aubin, J.-P. and Frankowska, H. (1990). *Set-valued Analysis* (Birkhäuser, Boston).
- [18] Aubin, T. (1976). Équations différentielle non linéaires et problème de Yamabe concernant la courbure scalaire, *J. Math. Pures Appl.* **55**, 269–296.
- [19] Aubin, T. (1982). *Nonlinear Analysis on Manifold. Monge–Ampère Equations* (Springer-Verlag, Heidelberg).
- [20] Aubin, T. (1998). *Some Nonlinear Problems in Riemannian Geometry* (Springer Verlag, Berlin).
- [21] Azagra, D. and Ferrera, J. (2005). Proximal calculus on Riemannian manifolds, *Mediterr. J. Math.* **2**, 437–450.
- [22] Azagra, D., Ferrera, J., and López-Mesas, J.F. (2005). Nonsmooth analysis and Hamilton-Jacobi equations on Riemannian manifolds, *J. Funct. Anal.* **220**, 304–361.
- [23] Bao, D., Chern, S.S., and Shen, Z. (2000). *An Introduction to Riemann–Finsler Geometry*, Graduate Texts in Mathematics, vol. 200 (Springer Verlag, Berlin).
- [24] Barbu, V. (1976). *Nonlinear Semigroups and Differential Equations in Banach Spaces* (Noordhoff, Leyden).
- [25] Bartolo, P., Benci, V., and Fortunato, D. (1983). Abstract critical point theorems and applications to some nonlinear problems with “strong” resonance at infinity, *Nonlinear Analysis TMA* **7**, 981–1012.
- [26] Bartsch, T. (1993). *Topological Methods for Variational Problems with Symmetries* (Springer Verlag, Berlin).
- [27] Bartsch, T. (1993). Infinitely many solutions of a symmetric Dirichlet problems, *Nonlinear Analysis TMA* **20**, 1205–1216.
- [28] Bartsch, T. and Figueiredo, D.G. de (1999). Infinitely many solutions of nonlinear elliptic systems, *Progr. Nonlinear Differential Equations Appl.* **35**, 51–67.
- [29] Bartsch, T., Liu, Z., and Weth, T. (2004). Sign changing solutions of superlinear Schrödinger equations, *Comm. Partial Differential Equations* **29**, 25–42.
- [30] Bartsch, T., Pankov, A., and Wang, Z.-Q. (2001). Nonlinear Schrödinger equations with steep potential well, *Commun. Contemp. Math.* **4**, 549–569.
- [31] Bartsch, T., Schneider, M., and Weth, T. (2004). Multiple solutions of a critical polyharmonic equation, *J. Reine Angew. Math.* **571**, 131–143.
- [32] Bartsch, T. and Wang, Z.-Q. (1995). Existence and multiplicity results for some superlinear elliptic problems in  $\mathbb{R}^N$ , *Comm. Partial Differential Equations* **20**, 1725–1741.
- [33] Bartsch, T. and Willem, M. (1993). Infinitely many non-radial solutions of an Euclidian scalar field equation, *J. Funct. Anal.* **117**, 447–460.

- [34] Bartsch, T. and Willem, M. (1993). Infinitely many radial solutions of a semilinear problem in  $\mathbb{R}^N$ , *Arch. Ration. Mech. Anal.* **124**, 261–276.
- [35] Benci, V., Fortunato, D., and Pisani, L. (1998). Solitons like solutions of a Lorentz invariant equation in dimension 3, *Rev. Math. Phys.* **10**, 315–344.
- [36] Berestycki, H., Cappuzzo Dolcetta, I., and Nirenberg, L. (1995). Variational methods for indefinite superlinear homogeneous elliptic problems, *Nonlinear Differential Equations Appl. (NoDEA)* **2**, 553–572.
- [37] Berestycki, H. and Lions, P.-L. (1983). Nonlinear scalar field equations I, II, *Arch. Ration. Mech. Anal.* **82**, 313–345, 347–375.
- [38] Bessaga, C. and Pelczyński, A. (1975). *Selected Topics in Infinite-Dimensional Topology* (Polish Scientific Publishers, Warszawa).
- [39] Bhattacharya, T., DiBenedetto, E., and Manfredi, J. (1989). Limits as  $p \rightarrow \infty$  of  $\Delta_p u_p = f$  and related extremal problems, *Rend. Sem. Mat. Univ. Pol. Torino, Fascicolo Speciale*, 15–68.
- [40] Bidaut-Véron, M.-F. and Véron, L. (1991). Nonlinear elliptic equations on compact Riemannian manifolds and asymptotics of Emden equations, *Invent. Math.* **106**, 489–539.
- [41] Bonanno, G. (2002). A minimax inequality and its applications to ordinary differential equations, *J. Math. Anal. Appl.* **270**, 210–229.
- [42] Bonanno, G. (2003). Some remarks on a three critical points theorem, *Nonlinear Analysis, TMA* **54**, 651–665.
- [43] Borenshtein, O.Y. and Shul’man, V.S. (1991). A minimax theorem, *Math. Notes* **50**, 752–754.
- [44] Borwein, J.M. (2007). Proximality and Chebyshev sets, *Optim. Lett.* **1**, 21–32.
- [45] Borwein, J.M. and Preiss, D. (1987). A smooth variational principle with applications to subdifferentiability and to differentiability of convex functions, *Trans. Amer. Math. Soc.* **303**, 517–527.
- [46] Breckner, B. and Varga, Cs. Infinitely many solutions for a class of systems of differential inclusions, preprint.
- [47] Brézis, H. (1983). *Analyse Fonctionnelle. Théorie et Applications* (Masson, Paris).
- [48] Brézis, H. and Browder, F. (1998). Partial differential equations in the 20th century, *Adv. Math.* **135**, 76–144.
- [49] Brézis, H. and Lieb, E.H. (1983). A relation between pointwise convergence of functions and convergence of functionals, *Proc. Amer. Math. Soc.* **88**, 486–490.
- [50] Brézis, H. and Nirenberg, L. (1983). Positive solutions of nonlinear elliptic equations involving critical Sobolev exponents, *Comm. Pure Appl. Math.*, **36**, 437–477.
- [51] Brézis, H. and Nirenberg, L. (1991). Remarks on finding critical points, *Comm. Pure Appl. Math.* **64**, 939–963.
- [52] Brézis, H. and Nirenberg, L. (1993).  $H^1$  versus  $C^1$  local minimizers, *C.R. Acad. Sci. Paris, Sér. I Math.* **317**, 465–472.
- [53] Bridston, M. and Haefliger, A. (1999). *Metric Spaces of Non-positive Curvature*, (Springer Verlag, Berlin).

- [54] Burton, G.R. (1985). Semilinear elliptic equations on unbounded domains, *Math. Z.* **190**, 519–525.
- [55] Busemann, H. (1955). *The Geometry of Geodesics* (Academic Press, New York).
- [56] Busemann, H. and Mayer, W. (1941). On the foundations of the calculus of variations, *Trans. Amer. Math. Soc.* **49**, 173–198.
- [57] Cammaroto, F., Chinni, A. and Di Bella, B. (2006). Some multiplicity results for quasilinear Neumann problems, *Arch. Math. (Basel)* **86**, 154–162.
- [58] Cammaroto, F., Chinni, A., and Di Bella, B. (2007). Multiple solutions for a quasilinear elliptic variational system on strip-like domains, *Proc. Edinb. Math. Soc.* **50**, 597–603.
- [59] Campa, I. and Degiovanni, M. (2000). Subdifferential calculus and nonsmooth critical point theory, *SIAM J. Optim.* **10**, 1020–1048.
- [60] Caponio, E., Javaloyes, M.A. and Masiello, A., On the energy functional on Finsler manifolds and applications to stationary spacetimes, preprint.
- [61] Carrião P.C., and Miyagaki, O. (2002). Existence of non-trivial solutions of elliptic variational systems in unbounded domains, *Nonlinear Analysis TMA* **51**, 155–169.
- [62] Cartan, E. (1928). *Leçons sur la Géométrie des Espaces de Riemann* (Gauthiers–Villars, Paris).
- [63] Cavazzuti, E., Pappalardo, M., and Passacantando, M. (2002). Nash equilibria, variational inequalities, and dynamical systems, *J. Optim. Theory Appl.* **114**, 491–506.
- [64] Cerami, G. (1978). Un criterio di esistenza per i punti critici su varietà illimitate, *Rend. Sci. Mat. Fis. Chim. Geol.* **112**, 332–336.
- [65] Chabrowski, J. (1997). Elliptic variational problems with indefinite nonlinearities, *Topol. Meth. Nonlin. Anal.* **9**, 221–231.
- [66] Chabrowski, J. (1997). *Variational Methods for Potential Operator Equations* (De Gruyter, Berlin–New York).
- [67] Chang, S.-Y. (2004). Maximal elements in noncompact spaces with application to equilibria, *Proc. Amer. Math. Soc.* **132**, 535–541.
- [68] Cheney, W. (2001). *Analysis for Applied Mathematics* (Springer, New York).
- [69] Cherrier, P. (1984). Problèmes de Neumann nonlinéaires sur les variétés Riemanniennes, *J. Funct. Anal.* **57**, 154–207.
- [70] Choquet, G. (1969). *Lectures on Analysis*, Vol. I (Math. Lectures Notes Ser., New York–Amsterdam).
- [71] Cîrstea, F.-C. and Rădulescu, V. (2001). On a class of quasilinear eigenvalue problems on unbounded domains, *Arch. Math. (Basel)* **77**, 337–346.
- [72] Clarke, F.H. (1983). *Nonsmooth Analysis* (John Wiley & Sons, New York).
- [73] Clarke, F.H. (1990). *Optimization and Nonsmooth Analysis*, Classics in Applied Mathematics, vol. 5 (Society for Industrial and Applied Mathematics (SIAM), Philadelphia).

- [74] Conti, M., Merizzi, L., and Terracini, S. (2000). Radial solutions of superlinear equations on  $\mathbb{R}^N$ . A global variational approach, *Arch. Ration. Mech. Anal.* **153**, 291–316.
- [75] Corvellec, J.-N., Degiovanni, M., and Marzocchi, M. (1993). Deformation properties for continuous functionals and critical point theory, *Topol. Methods Nonlinear Anal.* **1**, 151–171.
- [76] Costa, D.G. (1994). On a class of elliptic system in  $\mathbb{R}^N$ , *Electronic J. Differential Equations* **111**, 103–122.
- [77] Costa, D.G. and Magalhães, C.A. (1996). A unified approach to a class of strongly indefinite functionals, *J. Differential Equations* **125**, 521–548.
- [78] Coti Zelati, V. and Rabinowitz, P.H. (1992). Homoclinic type solutions for a semilinear elliptic PDE on  $\mathbb{R}^n$ , *Comm. Pure Appl. Math.* **45**, 1217–1269.
- [79] Cotsiolis, A. and Iliopoulos, D. (1991). Équations elliptiques non linéaires sur  $S^n$ . Le problème de Nirenberg, *C.R. Acad. Sci. Paris, Sér. I Math.* **313**, 607–609.
- [80] Cotsiolis, A. and Iliopoulos, D. (1995). Équations elliptiques non linéaires à croissance de Sobolev sur-critique, *Bull. Sci. Math.* **119**, 419–431.
- [81] Daneš, J. (1985). Equivalence of some geometric and related results of nonlinear functional analysis, *Comment. Math. Univ. Carolin.* **26**, 443–454.
- [82] Dautray, R. and Lions, J.-L. (1985). *Mathematical Analysis and Numerical Methods for Science and Technology, vol. 1: Physical Origins and Classical Methods* (Springer Verlag, Berlin).
- [83] Degiovanni, M. and Lancelotti, S. (1995). Perturbations of even nonsmooth functionals, *Differential Integral Equations* **8**, 981–992.
- [84] Degiovanni, M., Marino, A., and Tosques, M. (1985). Evolution equations with lack of convexity, *Nonlinear Anal.* **9**, 1401–1443.
- [85] Degiovanni, M. and Marzocchi, M. (1994). A critical point theory for nonsmooth functionals, *Ann. Mat. Pura Appl.* **167**, 73–100.
- [86] Degiovanni, M. and Schuricht, F. (1998). Buckling of nonlinearly elastic rods in the presence of obstacles treated by nonsmooth critical point theory, *Math. Ann.* **311**, 675–728.
- [87] Degiovanni, M. and Rădulescu, V. (1999). Perturbation of nonsmooth symmetric nonlinear eigenvalue problems, *C.R. Acad. Sci. Paris Sér. I* **329**, 281–286.
- [88] Di Falco, A.G. (2003). Infinitely many solutions to the Neumann problem for quasilinear elliptic systems, *Matematiche (Catania)* **LVIII**, 117–130.
- [89] Di Falco, A.G. (2003). Infinitely many solutions to the Neumann problem for quasilinear elliptic systems, *Matematiche (Catania)* **LX**, 163–179.
- [90] Diaz, J.-I. (1986). *Nonlinear Partial Differential Equations and Free Boundaries. Elliptic Equations*, Research Notes in Mathematics, Vol. 106 (Pitman, Boston-London-Melbourne).

- [91] Ding, W. (1986). On a conformally invariant elliptic equation on  $\mathbb{R}^n$ , *Comm. Math. Phys.* **107**, 331–335.
- [92] Dinu, T.-L. (2006). Entire solutions of Schrödinger elliptic systems with discontinuous nonlinearity and sign-changing potential, *Math. Modelling and Analysis* **11**, 1–14.
- [93] Do Carmo, M.P. (1992). *Riemannian Geometry* (Birkhäuser, Boston).
- [94] Drábek, P. (1995). Nonlinear eigenvalue problems for the  $p$ -Laplacian in  $\mathbb{R}^N$ , *Math. Nachr.* **173**, 131–139.
- [95] Drábek, P., Kufner, A., and Nicolosi, F. (1997). *Quasilinear Elliptic Equations with Degenerations and Singularities*, de Gruyter Series in Nonlinear Analysis and Applications (W. de Gruyter, Berlin–New York).
- [96] Drábek, P. and Pohozaev, S. (1997). Positive solutions for the  $p$ -Laplacian: application of the fibering method, *Proc. Roy. Soc. Edinburgh* **127A**, 703–726.
- [97] Drábek, P. and Simader, C.G. (1999). Nonlinear eigenvalue problems for quasilinear equations in unbounded domains, *Math. Nachr.* **203**, 5–30.
- [98] Druet, O., Hebey, E., and Robert, F. (2008). *Blow-up Theory for Elliptic PDE's in Riemannian Geometry* (Princeton University Press, Princeton).
- [99] Dupaigne, L., Ghergu, M., and Rădulescu, V. (2007). Lane–Emden–Fowler equations with convection and singular potential, *J. Math. Pures Appl.* **87**, 563–581.
- [100] Duvaut, G. and Lions, J.-L. (1982). *Inequalities in Mechanics and Physics* Springer-Verlag, Berlin-New York.
- [101] Efimov, N.V. and Stechkin, S.B. (1961). Approximate compactness and Chebyshev sets, *Soviet Math. Dokl.* **2**, 1226–1228.
- [102] Ekeland, I. (1974). On the variational principle, *J. Math. Anal. Appl.* **47**, 324–353.
- [103] Ekeland, I. (1990). *Convexity Methods in Hamiltonian Mechanics* (Springer Verlag, Berlin).
- [104] El Hamidi, A. (2004). Multiple solutions with changing sign energy to a nonlinear elliptic equation, *Commun. Pure Appl. Anal.* **3**, 253–265.
- [105] El Hamidi, A. (2004). Existence results to elliptic systems with nonstandard growth conditions, *J. Math. Anal. Appl.* **300**, 30–42.
- [106] Escobar, J. (1992). The Yamabe problem on manifolds with boundary, *J. Differ. Geom.* **35**, 21–84.
- [107] Escobar, J. (1992). Conformal deformation of Riemannian metric to a scalar flat metric with constant mean curvature on the boundary, *Ann. Math.* **136**, 1–50.
- [108] Escobar, J. (1996). Conformal metric with prescribed mean curvature on the boundary, *Calc. Var. Partial Differential Equations* **4**, 559–592.
- [109] Esteban, M. (1983). Nonlinear elliptic problems in strip-like domains: symmetry of positive vortex rings, *Nonlinear Analysis, TMA* **7**, 365–379.
- [110] Esteban, M. and Lions, P.-L. (1983). A compactness lemma, *Nonlinear Analysis, TMA* **7**, 381–385.

- [111] Fadel, E.R. and Hussein, S.Y. (1987). Relative cohomological index theories, *Adv. Math.* **64**, 1–31.
- [112] Fan, X.L. and Zhao, Y.Z. (2001). Linking and multiplicity results for the  $p$ -Laplacian on unbounded cylinders, *J. Math. Anal. Appl.* **260**, 479–489.
- [113] Faraci, F. and Kristály, A. (2007). One-dimensional scalar field equations involving an oscillatory nonlinear term, *Discrete Contin. Dyn. Syst.* **18**, 107–120.
- [114] Felmer, P. and Figueiredo, D.G. de (1994). On superquadratic elliptic systems, *Trans. Amer. Math. Soc.* **343**, 99–116.
- [115] Felmer, P., Manásevich, R., and Thélin, F. de (1992). Existence and uniqueness of positive solutions for certain quasilinear elliptic systems, *Comm. Partial Differential Equations* **17**, 2013–2029.
- [116] Figueiredo, D.G. de (1989). *The Ekeland Variational Principle with Applications and Detours* (Tata Institute of Fund. Research, Bombay).
- [117] Figueiredo, D.G. de (1998). Semilinear elliptic systems, *Nonlinear Functional Analysis and Applications to Differential Equations*, Trieste, 1997, 122–152. World Science Publ. River Edge, NJ.
- [118] Figueiredo, D.G. de, Gossez, J.-P., and Ubilla, P. (2003). Local superlinearity and sublinearity for indefinite semilinear elliptic problems, *J. Funct. Anal.* **199**, 452–467.
- [119] Figueiredo, D.G. de, Gossez, J.-P., and Ubilla, P. (2006). Multiplicity results for a family of semilinear elliptic problems under local superlinearity and sublinearity, *J. Eur. Math. Soc. (JEMS)* **8**, 269–286.
- [120] Figueiredo, D.G. de and Ruf, B. (2004). Elliptic systems with nonlinearities of arbitrary growth, *Mediterr. J. Math.* **1**, 417–431.
- [121] Filippucci, R., Pucci, P., and Rădulescu, V. (2008). Existence and non-existence results for quasilinear elliptic exterior problems with nonlinear boundary conditions, *Comm. Partial Differential Equations* **33**, 706–717.
- [122] Findley, W.L., Lai, J.S., and Onaran, K. (1976). *Creep and Relaxation of Nonlinear Viscoelastic Materials* (North Holland Publ. House, Amsterdam–New York–Oxford).
- [123] Floer, A. and Weinstein, A. (1986). Nonspreading wave packets for the cubic Schrödinger equation with a bounded potential, *J. Funct. Anal.* **69**, 397–408.
- [124] García-Azorero, J., Montefusco, E., and Peral, I. (2000). Bifurcation for the  $p$ -Laplacian in  $\mathbb{R}^N$ , *Adv. Diff. Equations* **5**, 435–464.
- [125] García-Azorero, J. and Peral, I. (1987). Existence and uniqueness for the  $p$ -Laplacian: nonlinear eigenvalues, *Comm. Partial Differential Equations* **12**, 1389–1430.
- [126] Gasinski, L. and Papageorgiu, N. (2005). *Nonsmooth Critical Point Theory and Nonlinear Boundary Value Problems* (Chapman & Hall/CRC, Boca Raton).
- [127] Georgiev, P. (2005). Parametric Borwein-Preiss variational principle and applications, *Proc. Amer. Math. Soc.* **133**, 3211–3225.

- [128] Ghergu, M. and Rădulescu, V. (2008). *Singular Elliptic Problems. Bifurcation and Asymptotic Analysis*. Oxford Lecture Series in Mathematics and its Applications, Vol. 37 (Oxford University Press, New York).
- [129] Ghoussoub, N. (1991). Location, multiplicity, and Morse indices of min-max critical points, *J. Reine Angew. Math.* **417**, 27–76.
- [130] Ghoussoub, N. and Preiss, D. (1989). A general mountain pass principle for locating and classifying critical points, *Ann. Inst. H. Poincaré, Anal. Non Linéaire* **6**, 321–330.
- [131] Gomes, J.M. and Sanchez, L. (2005). A variational approach to some boundary value problems in the half-line, *Z. Angew. Math. Phys.* **56**, 192–209.
- [132] Grognet, S. (2000). Théorème de Motzkin en courbure négative, *Geom. Dedicata* **79**, 219–227.
- [133] Grossinho, M.R. (1987). Some existence and bifurcation results for nonlinear elliptic problems in strip-like domains, *Ricerche Mat.* **36**, 127–138.
- [134] Grossinho, M.R. and Tersian, S. (2001). *An Introduction to Minimax Theorems and their Applications to Differential Equations* (Kluwer Academic Publishers, Dordrecht).
- [135] Grove, K. (1973). Condition (C) for the energy integral on certain path spaces and applications to the theory of geodesics, *J. Differential Geometry* **8**, 207–223.
- [136] Guillerme, J. (1994). Nash equilibrium for set-valued maps. *J. Math. Anal. Appl.* **187**, 705–715.
- [137] Hebey, E. (1996). *Sobolev Spaces on Riemannian Manifolds* (Springer Verlag, Heidelberg).
- [138] Hebey, E. (1999). *Nonlinear Analysis on Manifolds: Sobolev Spaces and Inequalities* (Courant Lecture Notes in Mathematics, New York).
- [139] Hebey, E., Variational methods and elliptic equations in Riemannian geometry, Notes from lectures given at ICTP, unpublished.
- [140] Hebey, E. and Vaugon, M. (1997). Sobolev spaces in the presence of symmetries, *J. Math. Pures Appl.* **76**, 859–881.
- [141] Hiriart-Urruty, J.-B. (1983). A short proof of the variational principle for approximate solutions of a minimization problem, *Amer. Math. Monthly* **90**, 206–207.
- [142] Hofer, H. (1982). Variational and topological methods in partially ordered Hilbert spaces, *Math. Annalen* **261**, 493–514.
- [143] Hulshof, J., Mitidieri, E., and Vorst, R. van der (1998). Strongly indefinite systems with critical Sobolev exponents, *Trans. Amer. Math. Soc.* **350**, 2349–2365.
- [144] Il'yasov, Y. and Runst, T. (2003). Positive solutions for indefinite inhomogeneous Neumann elliptic problems, *Electronic J. Diff. Equations* **57**, 1–21.
- [145] Ioffe, A.D. and Tikhomirov, V.M. (1997). Some remarks on variational principles, *Mathematical Notes* **61**, 248–253.
- [146] Jabri, Y. (2003). *The Mountain Pass Theorem. Variants, Generalizations and Some Applications*. Encyclopedia of Mathematics and its Applications, Vol. 95 (Cambridge University Press, Cambridge).

- [147] Jayne, J.E. and Rogers, C.A. (2002). *Selectors* (Princeton University Press, Princeton).
- [148] Jebelean, P. (2008). Infinitely many solutions for ordinary  $p$ -Laplacian systems with nonlinear boundary conditions, *Commun. Pure Appl. Anal.* **7**, 267–275.
- [149] Jebelean, P. and Moroşanu, G. (2005). Mountain pass type solutions for discontinuous perturbations of the vector  $p$ -Laplacian, *Nonlinear Funct. Anal. Appl.* **10**, 591–611.
- [150] Jebelean, P. and Moroşanu, G. (2006). Ordinary  $p$ -Laplacian systems with nonlinear boundary conditions, *J. Math. Anal. Appl.* **313**, 738–753.
- [151] Jones, C. and Küpper, T. (1986). On the infinitely many solutions of a semilinear elliptic equation, *SIAM J. Math. Anal.* **17**, 803–835.
- [152] Jost, J. (1997). *Nonpositivity Curvature: Geometric and Analytic Aspects* (Birkhäuser Verlag, Basel).
- [153] Kachanov, L.M. (1967). *The Theory of Creep* (National Lending Library for Science and Technology, Boston Spa, Yorkshire).
- [154] Kachanov, L.M. (1971). *Foundations of the Theory of Plasticity* (North Holland Publishing House, Amsterdam–London).
- [155] Kassay, G., Kolumbán, J., and Páles, Zs. (1999). On Nash stationary points, *Publ. Math. Debrecen* **54**, 267–279.
- [156] Kay, D.C. (1967). The Ptolemaic inequality in Hilbert geometries, *Pacific J. Math.* **21**, 293–301.
- [157] Kawohl, B. (1990). On a family of torsional creep problems, *J. Reine Angew. Math.* **410**, 1–22.
- [158] Kelly, P. and Straus, E. (1958). Curvature in Hilbert geometry, *Pacific J. Math.* **8**, 119–125.
- [159] Kobayashi, J. (2001). A principle of symmetric criticality for variational inequalities, *Mathematical aspects of modelling structure formation phenomena*, Gakuto Internat. Ser. Math. Sci. Appl. **17**, 94–106.
- [160] Kobayashi, J. and Ôtani, M. (2004). The principle of symmetric criticality for non-differentiable mappings, *J. Funct. Anal.* **214**, 428–449.
- [161] Kourougenis, N. and Papageorgiou, N. (2000). Nonsmooth critical point theory and nonlinear elliptic equations at resonance, *J. Austral. Math. Soc.* **69**, 245–271.
- [162] Kozma, L., Kristály, A., and Varga, Cs. (2004). Critical point theorems on Finsler manifolds, *Beiträge Algebra Geom.* **45**, 47–59.
- [163] Krasnosel’skii, M.A. (1964). *Topological Methods in the Theory of Nonlinear Integral Equations* (Pergamon Press Book, Macmillan, New York).
- [164] Krawcewicz, W. and Marzantowicz, W. (1990). Some remarks on the Lusternik-Schnirelman method for non-differentiable functionals invariant with respect to a finite group action, *Rocky Mountain J. Math.* **20**, 1041–1049.
- [165] Kristály, A. (2004). An existence result for gradient-type systems with a nondifferentiable term on unbounded strips, *J. Math. Anal. Appl.* **229**, 186–204.

- [166] Kristály, A. (2005). Existence of two non-trivial solutions for a class of quasilinear elliptic variational systems on stripe-like domains, *Proc. Edinb. Math. Soc.* **48**, 1–13.
- [167] Kristály, A. (2005). Infinitely many radial and non-radial solutions for a class of hemivariational inequalities, *Rocky Mountain J. Math.* **35**, 1173–1190.
- [168] Kristály, A. (2006). Infinitely many solutions for a differential inclusion problem in  $\mathbb{R}^N$ , *J. Differential Equations* **220**, 511–530.
- [169] Kristály, A. (2006). Existence of nonzero weak solutions for a class of elliptic variational inclusions systems in  $\mathbb{R}^N$ , *Nonlinear Analysis* **65**, 1578–1594.
- [170] Kristály, A. (2008). Detection of arbitrarily many solutions for perturbed elliptic problems involving oscillatory terms, *J. Differential Equations* **245**, 3849–3868.
- [171] Kristály, A. (2009). Asymptotically critical problems on higher-dimensional spheres, *Discrete Contin. Dyn. Syst.* **23**, 919–935.
- [172] Kristály, A. (2009). Metric projections on Finsler manifolds, preprint.
- [173] Kristály, A. and Kozma, L. (2006). Metric characterization of Berwald spaces of non-positive flag curvature, *Journal of Geometry and Physics* **56**, 1257–1270.
- [174] Kristály, A. and Marzantowicz, W. (2008). Multiplicity of symmetrically distinct sequences of solutions for a quasilinear problem in  $\mathbb{R}^N$ , *Nonlinear Differential Equations and Applications (NoDEA)* **15**, 209–226.
- [175] Kristály, A., Moroşanu, G., and Róth Á. (2008). Optimal placement of a deposit between markets: Riemann-Finsler geometrical approach, *J. Optim. Theory Appl.* **139**, 263–276.
- [176] Kristály, A., Moroşanu, G., and Tersian, S. (2007). Quasilinear elliptic problems in  $\mathbb{R}^N$  involving oscillatory nonlinearities, *J. Differential Equations* **235**, 366–375.
- [177] Kristály, A., Motreanu, V., and Varga, Cs. (2005). A minimax principle with a general Palais-Smale condition, *Commun. Appl. Anal.* **9**, 285–297.
- [178] Kristály, A. and Rădulescu, V. (2009). Sublinear eigenvalue problems on compact Riemannian manifolds with applications in Emden-Fowler equations, *Studia Mathematica* **191**, 237–246.
- [179] Kristály, A. and Varga, Cs. (2003). Set-valued versions of Ky Fan’s inequality with application to variational inclusion theory, *J. Math. Anal. Appl.* **282**, 8–20.
- [180] Kristály, A. and Varga, Cs. (2004). *An Introduction to critical Point Theory for Nonsmooth Functions* (Casa Cărţii de Ştiinţă, Cluj-Napoca).
- [181] Kristály, A. and Varga, Cs. (2007). Multiple solutions for elliptic problems with singular and sublinear potentials, *Proc. Amer. Math. Soc.* **135**, 2121–2126.
- [182] Kristály, A., Varga, Cs., and Kozma, L. (2004). The dispersing of geodesics in Berwald spaces of non-positive flag curvature, *Houston J. Math.* **30**, 413–420.

- [183] Kristály, A., Varga, Cs., and Varga, V. (2005). An eigenvalue problem for hemivariational inequalities with combined nonlinearities on an infinite strip, *Nonlinear Analysis* **63**, 260–272.
- [184] Kulpa, W. and Szymanski, A. (2004). Infimum principle, *Proc. Amer. Math. Soc.* **132**, 203–210.
- [185] Kuratowski, K. and Ryll-Nardzewski, C. (1965). A general theorem on selectors, *Bull. Acad. Pol. Sci.* **13**, 397–403.
- [186] Lao, S.Y. (1992). Nonlinear  $p$ -Laplacian problems on unbounded domains, *Proc. Amer. Math. Soc.* **115**, 1037–1045.
- [187] Ledyaev, Y.S. and Zhu, Q.J. (2007). Nonsmooth analysis on smooth manifolds, *Trans. Amer. Math. Soc.* **359**, 3687–3732.
- [188] Lee, M.J. and Parker, T.H. (1987). The Yamabe problem, *Bull. Amer. Math. Soc.* **17**, 37–91.
- [189] Li, S. and Liu, Z. (2002). Perturbations from symmetric elliptic boundary value problems, *J. Differential Equations* **185**, 271–280.
- [190] Li, S. and Wang, Z.-Q. (2002). Ljusternik-Schnirelman theory in partially ordered Hilbert spaces, *Trans. Amer. Math. Soc.* **354**, 3207–3227.
- [191] Li, Y.Y. and Li, A. (2006). A fully nonlinear version of the Yamabe problem on manifold with boundary, *J. Eur. Math. Soc.* **8**, 295–316.
- [192] Lieb, E.H. and Loss, M. (1997). *Analysis* Graduate Studies in Mathematics, Vol. 14 (American Mathematical Society, Providence, RI).
- [193] Lions, P.-L. (1982). Symétrie et compacité dans les espaces de Sobolev, *J. Funct. Anal.* **49**, 312–334.
- [194] Lions, P.-L. (1984). The concentration–compactness principle in the calculus of variations, *Ann. Inst. H. Poincaré, Anal. Non Linéaire* **1**, 109–145, 223–283.
- [195] Lisei, H., Varga, Cs., and Horváth, A. (2008). Multiplicity results for a class of quasilinear eigenvalue problems on unbounded domains, *Arch. Math.* **90**, 256–266.
- [196] Lisei, H., Moroşanu, G., and Varga, Cs. (2009). Multiplicity results for double eigenvalue problem involving the  $p$ -Laplacian, *Taiwanese Journal of Mathematics* **13**, 1095–1110.
- [197] Ljusternik, L. and Schnirelman, L. (1934). *Méthodes Topologiques dans les Problèmes Variationnels* (Hermann, Paris).
- [198] Loewen, P.D. and Wang, X. (2001). A generalized variational principle, *Canad. J. Math.* **53**, 1174–1193.
- [199] Luo, Q. (2003). Generic stability of Nash equilibria for set-valued mappings, *Acta Math. Sinica (Chin. Ser.)* **46**, 925–930.
- [200] Manásevich, R. and Mawhin, J. (1998). Periodic solutions for nonlinear systems with  $p$ -Laplacian like operators, *J. Differential Equations* **145**, 367–393.
- [201] Manásevich, R. and Mawhin, J. (2000). Boundary value problems for nonlinear perturbations of vector  $p$ -Laplacian like operators, *J. Korean Math. Soc.* **37**, 665–685.
- [202] Marano, S. and Motreanu, D. (2002). On a three critical points theorem for non-differentiable functions and applications to nonlinear boundary value problems, *Nonlinear Analysis* **48**, 37–52.

- [203] Marano, S. and Motreanu, D. (2002). Infinitely many critical points of non-differentiable functions and applications to a Neumann-type problem involving the  $p$ -Laplacian, *J. Differential Equations* **182**, 108–120.
- [204] Marcus, M. and Mizel, M. (1979). Every superposition operator mapping one Sobolev space into another is continuous, *J. Funct. Anal.* **33**, 217–229.
- [205] Marino, A. (1989). The calculus of variations and some semilinear variational inequalities of elliptic and parabolic type, *Partial Differential Equations and the Calculus of Variations. Volume II. Essays in Honor of Ennio De Giorgi*, Birkhäuser, Boston.
- [206] Marino, A. and Scolozzi, D. (1983). Geodetiche con ostacolo, *Boll. Un. Mat. Ital. B* **2**, 1–31.
- [207] Matsumoto, M. (1989). A slope of a mountain is a Finsler surface with respect to a time measure, *J. Math. Kyoto Univ.* **29**, 17–25.
- [208] Mawhin, J. (2000). Some boundary value problems for Hartman type perturbations of the ordinary vector  $p$ -Laplacian, *Nonlinear Anal.* **40**, 497–503.
- [209] Mawhin, J. and Willem, M. (1989). *Critical Point Theory and Hamiltonian Systems* (Springer, New York).
- [210] McClendon, J.F. (1983). Minimax and variational inequalities for compact spaces, *Proc. Amer. Math. Soc.* **89**, 717–721.
- [211] McLeod, J.B., Stuart, C.A., and Troy, W.C. (2003). Stability of standing waves for some nonlinear Schrödinger equations, *Differential Integral Equations* **16**, 1025–1038.
- [212] Mercuri, F. (1976). Closed geodesics on Finsler manifolds, *Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur.* **60**, 111–118.
- [213] Mezei, I. and Varga, Cs. (2008). Multiplicity results for a double eigenvalue quasilinear problem on unbounded domain, *Nonlinear Analysis TMA* **69**, 4099–4105.
- [214] Mitidieri, E. (1993). A Rellich type identity and applications, *Comm. Partial Differential Equations* **18**, 125–151.
- [215] Mitidieri, E. and Bozhkov, Y. (2003). Existence of multiple solutions for quasilinear systems via fibering method, *J. Differential Equations* **190**, 239–267.
- [216] Moalla, F. (1966). Sur quelques théoremes globaux en géometrie finslerienne, *Ann. Math. Pura Appl.* **73**, 319–365.
- [217] Montefusco, E. and Rădulescu, V. (2001). Nonlinear eigenvalue problems for quasilinear operators on unbounded domains, *Nonlinear Differential Equations Appl. (NoDEA)* **8**, 481–497.
- [218] Morgan, J. and Scalzo, V. (2007). Pseudocontinuous functions and existence of Nash equilibria, *J. Math. Econom.* **43**, 174–183.
- [219] Morrey, C.B. Jr. (1966). *Multiple Integrals in the Calculus of Variations* (Springer-Verlag, New York).
- [220] Morse, M. (1925). Relations between the critical points of a real function of  $n$  independent variables, *Trans. Amer. Math. Soc.* **27**, 345–396.
- [221] Morse, M. (1934). The calculus of variations in the large, *Amer. Math. Soc. Colloq. Publ.* **18**.

- [222] Moskovitz, D. and Dines, L. (1939). Convexity in a linear space with an inner product, *Duke Math. J.* **5**, 520–534.
- [223] Motreanu, D. and Panagiotopoulos, P.D. (1998). *Minimax Theorems and Qualitative Properties of the Solutions of Hemivariational Inequalities*, Nonconvex Optimization Applications, vol. 29 (Kluwer, Dordrecht).
- [224] Motreanu, D. and Rădulescu, V. (2003). *Variational and Nonvariational Methods in Nonlinear Analysis and Boundary Value Problems* (Kluwer Academic Publishers, Dordrecht).
- [225] Motreanu, D. and Varga, Cs. (1997). Some critical point results for locally Lipschitz functionals, *Comm. Appl. Nonlin. Anal.* **4**, 17–33.
- [226] Motreanu, D. and Varga, Cs. (1999). A nonsmooth equivariant minimax principle, *Comm. Appl. Anal.* **3**, 115–130.
- [227] Nash, J.F. (1950). Equilibrium points in  $n$ -person games, *Proc. Nat. Acad. Sci. U. S. A.* **36**, 48–49.
- [228] Nash, J.F. (1951). Non-cooperative games, *Annals Math.* **54**, 286–295.
- [229] Nessah, R. and Kerstens, K. Characterization of the existence of Nash equilibria with non-convex strategy sets, Document du travail LEM, 2008-19, preprint, [http://lem.cnrs.fr/Portals/2/actus/DP\\_200819.pdf](http://lem.cnrs.fr/Portals/2/actus/DP_200819.pdf)
- [230] Obata, M. (1971). The conjectures on conformal transformations of Riemannian manifolds, *J. Diff. Geom.* **6**, 247–258.
- [231] Obersnel, F. and Omari, P. (2006). Positive solutions of elliptic problems with locally oscillating nonlinearities, *J. Math. Anal. Appl.* **323**, 913–929.
- [232] Omari, P. and Zanolin, F. (1996). Infinitely many solutions of a quasilinear elliptic problem with an oscillatory potential, *Comm. Partial Differential Equations* **21**, 721–733.
- [233] O’Neill, B. (1983). *Semi-Riemannian Geometry. With Applications to Relativity* (Academic Press, New York).
- [234] Palais, R. (1966). Ljusternik–Schnirelmann theory on Banach manifolds, *Topology* **5**, 115–132.
- [235] Palais, R. (1979). The principle of symmetric criticality, *Comm. Math. Phys.* **69**, 19–30.
- [236] Palais, R. (1984). Applications of the Symmetric Criticality Principle in Mathematical Physics and Differential Geometry, *Proceedings of the 1981 Shanghai Symposium on Differential Geometry and Differential Equations* (Shanghai/Hefei 1981), 247-301, Science Press, Beijing.
- [237] Palais, R. and Smale, S. (1964). A generalized Morse theory, *Bull. Amer. Math. Soc.* **70**, 165–171.
- [238] Pedersen, F.P. (1952). On spaces with negative curvature, *Mat. Tidsskrift B* **1952**, 66-89.
- [239] Péllissier, M.C. and Reynaud, M.L. (1974). Étude d’un modèle mathématique d’écoulement de glacier, *C.R. Acad. Sci. Paris, Sér. I Math.* **279**, 531–534.
- [240] Penot, J.-P. (1986). The drop theorem, the petal theorem and Ekeland’s variational principle, *Nonlinear Anal.* **10**, 813–822.

- [241] Pflüger, K. (1998). Existence and multiplicity of solutions to a  $p$ -Laplacian equation with nonlinear boundary condition, *Electronic J. Differential Equations* **10**, 1–13.
- [242] Pflüger, K. (1998). Compact traces in weighted Sobolev spaces, *Analysis* **18**, 65–83.
- [243] Phelps, R.R. (1957). Convex sets and nearest points, *Proc. Amer. Math. Soc.* **8**, 790–797.
- [244] Phelps, R.R. (1974). Support cones in Banach spaces and their applications, *Adv. Math.* **13**, 1–19.
- [245] Pohozaev, S.I. (1979). On one approach to nonlinear equations, *Dokl. Akad. Nauk* **247**, 1327–1331.
- [246] Pohozaev, S.I. (1988). On a constructive methods in calculus of approach to nonlinear equations, *Dokl. Akad. Nauk* **298**, 1330–1333.
- [247] Pohozaev, S.I. (1990). On fibering method for the solutions of nonlinear boundary value problems, *Trudy Mat. Inst. Steklov* **192**, 146–163.
- [248] Pohozaev, S.I. (1999). The fibering method and its applications to nonlinear boundary value problems, *Rend. Istit. Mat. Univ. Trieste* **31**, 235–305.
- [249] Poincaré, H. (1890). Sur les équations aux dérivées partielles de la physique mathématique, *Amer. J. Math.* **12**, 211–294.
- [250] Pucci, P. and Serrin, J. (1984). Extensions of the mountain pass theorem, *J. Funct. Anal.* **59**, 185–210.
- [251] Pucci, P. and Serrin, J. (1985). A mountain pass theorem, *J. Diff. Equations* **60**, 142–149.
- [252] Pucci, P. and Serrin, J. (2007). *The Maximum Principle*. Progress in Nonlinear Differential Equations and their Applications, Vol. 73 (Birkhäuser Verlag, Basel).
- [253] Rabinowitz, P.H. (1986). *Minimax Methods in Critical Point Theory with Applications to Differential Equations*, CBMS Reg. Conference Series in Mathematics, **65** (American Mathematical Society, Providence, RI).
- [254] Rabinowitz, P.H. (1992). On a class of nonlinear Schrödinger equations, *Z. Angew. Math. Phys.* **43**, 270–291.
- [255] Rădulescu, V. (2008). *Qualitative Analysis of Nonlinear Elliptic Partial Differential Equations: Monotonicity, Analytic, and Variational Methods*. Contemporary Mathematics and Its Applications, Vol. 6 (Hindawi Publ. Corp., New York).
- [256] Rapcsák T. (1989). Minimum problems on differentiable manifolds, *Optimization* **20**, 3–13.
- [257] Rapcsák T. (1991). Geodesic convexity in nonlinear optimization, *J. Optim. Theory Appl.* **69**, 169–183.
- [258] Ricceri, B. (1999). A new method for the study of nonlinear eigenvalue problems, *C.R. Acad. Sci. Paris, Sér. I* **328**, 251–256.
- [259] Ricceri, B. (2000). On a three critical points theorem, *Arch. Math.* **75**, 220–226.
- [260] Ricceri, B. (2000). A general variational principle and some of its applications, *J. Comput. Appl. Math.* **113**, 401–410.

- [261] Ricceri, B. (2001). A further improvement of a minimax theorem of Borenshtein and Shul'man, *J. Nonlinear Convex Anal.* **2**, 279–283.
- [262] Ricceri, B. (2001). Infinitely many solutions of the Neumann problem for elliptic equations involving the  $p$ -Laplacian, *Bull. London Math. Soc.* **33**, 331–340.
- [263] Ricceri, B. (2002). Three solutions for a Neumann problem, *Topol. Meth. Nonlin. Anal.* **20**, 275–281.
- [264] Ricceri, B. (2004). Sublevel sets and global minima of coercive functionals and local minima of their perturbations, *J. Nonlin. Convex Anal.* **5**, 157–168.
- [265] Ricceri, B. (2005). A general multiplicity theorem for certain nonlinear equations in Hilbert spaces, *Proc. Amer. Math. Soc.* **133**, 3255–3261.
- [266] Ricceri, B. (2006). Minimax theorems for limits of parametrized functions having at most one local minimum lying in certain set, *Topol. Appl.*, **153**, 3308–3312.
- [267] Ricceri, B. A three critical points theorem revisited, *Nonlin. Anal.* **70**, 3084–3089.
- [268] Saint Raymond, J. (2000). On a minimax theorem, *Arch. Math.* **74**, 432–437.
- [269] Saint Raymond, J. (2002). On the multiplicity of the solutions of the equation  $-\Delta u = \lambda f(u)$ , *J. Differential Equations* **180**, 65–88.
- [270] Salvatore, A. (2008). Multiple solutions for elliptic systems with nonlinearities of arbitrary growth, *J. Differential Equations* **244**, 2529–2544.
- [271] Schoen, R. (1984). Conformal deformation of a Riemannian metric to constant scalar curvature, *J. Diff. Geom.* **20**, 479–495.
- [272] Schwartz, J.T. (1964). Generalizing the Lusternik-Schnirelman theory of critical points, *Comm. Pure Appl. Math.* **17**, 307–315.
- [273] Serre, J.-P. (1951). Homologie singulière des espaces fibrés, *Ann. Math.* **54**, 425–505.
- [274] Servadei, R. (2006). Existence results for semilinear elliptic variational inequalities with changing sign nonlinearities, *Nonlinear Differential Equations Appl. (NoDEA)* **13**, 311–335.
- [275] Shen, Z. (2003). Projectively flat Randers metrics with constant flag curvature, *Math. Ann.* **325**, 19–30.
- [276] Shen, Z. (2003). Projectively flat Finsler metrics of constant flag curvature, *Trans. Amer. Math. Soc.* **355**, 1713–1728.
- [277] Shen, Z. (2005). Finsler manifolds with nonpositive flag curvature and constant  $S$ -curvature, *Math. Z.* **249**, 625–639.
- [278] Shen, Z. Some open problems in Finsler geometry,  
<http://www.math.iupui.edu/~zshen/Research/papers/Problem.pdf>
- [279] Showalter, R.E. and Walkington, N.J. (1991). Diffusion of fluid in a fissured medium with microstructure, *SIAM J. Math. Anal.* **22**, 1702–1722.
- [280] Sirakov, B. (2000). Existence and multiplicity of solutions of semi-linear elliptic equations in  $\mathbb{R}^N$ , *Calc. Var. Partial Differential Equations* **11**, 119–142.

- [281] Socié-Méthou, E. (2004). Behaviour of distance functions in Hilbert-Finsler geometry, *Differential Geom. Appl.* **20**, 1–10.
- [282] Stegall, C. (1978). Optimization of functions on certain subsets of Banach spaces, *Math. Ann.* **236**, 171–176.
- [283] Strauss, W.A. (1977). Existence of solitary waves in higher dimensions, *Commun. Math. Phys.* **55**, 149–162.
- [284] Struwe, M. (1990). *Variational Methods*, (Springer-Verlag, Berlin).
- [285] Sullivan, F. (1981). A characterization of complete metric spaces, *Proc. Amer. Math. Soc.* **83**, 345–346.
- [286] Szulkin, A. (1986). Minimax methods for lower semicontinuous functions and applications to nonlinear boundary value problems, *Ann. Inst. H. Poincaré, Anal. Non Linéaire* **3**, 77–109.
- [287] Takahashi, W. (1991). Existence theorems generalizing fixed point theorems for multivalued mappings, *Fixed point theory and applications (Marseille, 1989)*, 397–406, Pitman Res. Notes Math. Ser., Vol. 252 (Longman Sci. Tech., Harlow).
- [288] Tala, J.E. and Marchi, E. (1996). Games with non-convex strategy sets, *Optimization* **37**, 177–181.
- [289] Trudinger, N. (1968). Remarks concerning the conformal deformation of riemannian structures on compact manifold, *Ann. Scuola Norm. Sup. Pisa* **22**, 265–274.
- [290] Tsar'kov, I.G. (2004). Nonunique solvability of certain differential equations and their connection with geometric approximation theory, *Math. Notes* **75**, 259–271.
- [291] Udriște, C. (1994). *Convex Functions and Optimization Methods on Riemannian Manifolds*, Mathematics and its Applications, vol. 297 (Kluwer Academic Publishers, Dordrecht).
- [292] Vázquez, J.-L. and Véron, L. (1991). Solutions positives d'équations elliptiques semi-linéaires sur des variétés riemanniennes compactes, *C.R. Acad. Sci. Paris, Sér. I Math.* **312**, 811–815.
- [293] Vélin, J. and Thélin, F. de (1993). Existence and nonexistence of non-trivial solutions for some nonlinear elliptic systems, *Rev. Mat. Univ. Complut. Madrid* **6**, 153–194.
- [294] Vorst, R. van der (1992). Variational identities and applications to differential systems, *Arch. Rational Mech. Anal.* **116**, 375–398.
- [295] Walter, R. (1976). Some analytical properties of geodesically convex sets, *Abh. Math. Sem. Univ. Hamburg* **45**, 263–282.
- [296] Westfall, R.S. (1980). *Never at Rest. A Biography of Isaac Newton* (Cambridge University Press, Cambridge).
- [297] Whitehead, J. H. C. (1932). Convex region in the geometry of paths, *Quarterly J. Math. Oxford* **3**, 33–42.
- [298] Willem, M. (1996). *Minimax Theorems* (Birkhäuser, Boston).
- [299] Yamabe, H. (1960). On a deformation of Riemannian structures on compact manifolds, *Osaka Math. J.* **12**, 21–37.
- [300] Yu, H. and Zhang, Z. (2007). Pure strategy equilibria in games with countable actions, *J. Math. Econom.* **43**, 192–200.

- [301] Zabreiko, P.P. and Krasnoselskii, M.A. (1971). The solvability of non-linear operator equations, *Funct. Anal. Appl.* **5**, 206–208.
- [302] Zeidler, E. (1985). *Nonlinear Functional Analysis and its Applications*, Vol. III (Springer Verlag, Berlin–Heidelberg–New York).
- [303] Zhou, Y. and Huang, Y. (2000). Existence of solutions for a class of elliptic variational inequality, *J. Math. Anal. Appl.* **250**, 187–195.
- [304] Ziad, A. (1999). Pure strategy Nash equilibria of non-zero-sum two-person games: non-convex case, *Econom. Lett.* **62**, 307–310.
- [305] Ziemer, W. (1989). *Weakly Differentiable Functions. Sobolev Spaces and Functions of Bounded Variation*. Graduate Texts in Mathematics, Vol. 120 (Springer, New York).
- [306] Zou, W. (2006). On finding sign-changing solutions, *J. Funct. Anal.* **234**, 364–419.

# Index

- (PS)-condition, 7
- (PS)<sub>c</sub>-condition, 7
- (PSZ)<sub>c</sub>-condition, 36
- ( $\nabla^2$ )<sub>ij</sub>, 367
- (epi)<sub>c</sub>, 60
- ( $g^{ij}$ ), 267
- ( $g_{ij}$ ), 267
- $BV(\Omega; \mathbb{R}^N)$ , 59
- $C^1$  local minimizer, 31
- $C^1$ -manifold, 99
- $C^{k,\alpha}(M)$ , 368
- $DF(u, c)$ , 382
- $D^\alpha u$ , 361
- $D^{k,p}(\mathbb{R}^N)$ , 364
- $F^{+1}(M)$ , 381
- $F^{-1}(M)$ , 381
- $H_0^1$  local minimizer, 31
- $H^1(\mathbb{R}^N)$ , 365
- $H^{m,p}(\Omega)$ , 362
- $H_{0,s}^1(\Omega)$ , 47
- $H_0^1(\Omega)$ , 366
- $H_G^1(\mathbb{R}^N)$ , 365
- $K$ -pseudo-convex, 336
- $K_c$ , 376
- $L^\infty(\Omega)$ , 360
- $L^p(\Omega)$ , 360
- $L_{loc}^p(\Omega)$ , 360
- $L^m(\Gamma; w_2)$ , 88
- $L^p(\Omega; w_1)$ , 88
- $N_S(x)$ , 359
- $TM$ , 15
- $TM^*$ , 16
- $T_K(u)$ , 382
- $T_p M$ , 16
- $W^{-m,p'}(\Omega)$ , 362
- $W^{m,p}(\Omega)$ , 362
- $W_{0,G}^{1,p}$ , 366
- $W_0^{m,p}(\Omega)$ , 362
- $X^*$ , 7
- $\Delta_{\mathbb{S}^n}$ , 280
- $\Delta_p u$ , 282
- $\Gamma$  function, 364
- $\Gamma_h$ , 17
- $\gamma(A)$ , 371
- $\mathcal{C}^p$ , 367
- $\nabla^k$ , 266
- $\nabla_p$ , 86
- $\omega_n$ , 364
- $\mathbb{R}$ , 379
- $\partial f(u)$ , 378
- $\pi(g)$ , 38
- $dv(g)$ , 367
- $f^0(u; v)$ , 378
- $f_\varepsilon^0(u; v)$ , 377
- $f^a$ , 379
- $f_c$ , 376
- $p$ -Laplace operator, 282
- $\text{Limsup}_{u' \rightarrow u} F(u')$ , 383
- $\text{cat}(M)$ , 17
- $\text{cat}_M(A)$ , 17, 370
- $\text{cuplong}(M)$ , 371
- $\text{dist}_S(x)$ , 359
- $\text{dist}_S^0(x; v)$ , 359
- $\text{epi}(f)$ , 355
- Čech homology, 296
- absolute neighborhood retract, 17
- action of a group, 48
- acyclic set, 296
- Aleksandrov curvature condition, 295
- almost critical point, 7
- altitude
  - positive, 13
  - zero, 15
- Arzelà–Ascoli theorem, 32
- Asplund space, 8
- asymptotically critical growth, 259
- averaging operator, 40

- axially symmetric function, 47
- backward proximal set, 312
- Banach manifold, 15
- Banach space, 3
- Banach-Finsler manifold, 16
  - complete, 16
- Berwald space, 289, 294, 295
- best approximation, 356
- best Sobolev constant, 364
- Bishop-Phelps theorem, 8
- bootstrap, 32
- Borwein-Preiss variational principle, 8
- boundary condition
  - Dirichlet, 264
- bounded set
  - backward, 288, 303
  - forward, 288
- Brézis-Lieb lemma, 361
- brachistochrone problem, ix
- Brouwer fixed point theorem, 296
- Busemann
  - non-positive curvature space, 291
  - NPC inequality, 291, 295
- Carathéodory function, 90, 100, 112
- Caristi fixed point theorem, 227
- Cauchy sequence, 95
- Cerami condition, 15
- chain rule, 73
- Chebyshev set, 356
- Chern connection, 285
- Christofel symbol, 289
- Christoffel symbols, 267
- Clarke directional derivative, 374
- Clarke subdifferential, 374
- closed symmetric manifold, 99
- cocycles, 371
- coercive functional, 24
- compact embedding, 363
- compact Riemann manifold, 266
- compact topological space, 3
- compatible, 364
- complete metric space, 43
- concave-convex nonlinearity, 197, 202
- concentration compactness principle, 124
- condition
  - Cerami, 15
  - Palais-Smale, 7, 17, 376
- conditionally critical point, 41
- cone, 356
  - contingent, 382
  - convex, 355
  - normal, 359
  - positive, 356
- conformal Laplacian, 268
- conformal metric, 267
- conjugate point, 287, 293
- constrained critical point, 99
- constrained minimization problem, 107
- contingent cone, 382
- contingent derivative, 382
- contingent point
  - Nash, 334
- continuous embedding, 363
- convex cone, 355
- convex function, 35, 37, 354
- convex set, 353
- core, 381
- cotangent bundle, 16
- covariant
  - derivative, 233, 285
  - tensor field, 233
- creep, 87
- critical exponent, 197
- critical hyperbola, 156
- critical point, 7, 101, 113, 118, 195
  - conditionally, 160
  - constrained, 99
  - lower, 378
  - nonstable, 12
  - of Degiovanni-type, 376
  - of Nash-type, 341
  - of Szulkin-type, 35
  - stable, 11
- critical value, 35
- curvature tensor, 285
- Daneš drop theorem, 43
- deformation, 370
- deformation lemma, 376
- derivative
  - Clarke generalized directional, 374
  - directional, 93, 359
  - Fréchet, 93
  - Gâteaux, 7, 22
  - generalized directional, 377
  - Radon-Nikodym, 70
- derivative rule of vector fields, 286
- dilation, 353
- direct method, 224
- directional derivative, 93, 359
- Dirichlet boundary value condition, 264
- Dirichlet integral, x
- distance function, 359
- divergence theorem, 88
- double eigenvalue problem, 24, 99, 100
- dual Finsler structure, 16
- duality mapping, 7
- duality pairing, 373
- Eberlein-Smulyan theorem, 148
- eigenspace, 280
- eigenvalue, 280, 282
- eigenvalue problem, 52

- double, 24
- quasilinear, 87
- Ekeland variational principle, 4, 58
- elliptic regularity, 369
- elliptic system, 125
- embedding, 89, 92
  - compact, 363
  - continuous, 363
- Emden-Fowler equation, 231, 263
- energy
  - kinetic, 123
  - potential, 123
  - total, 123
- energy functional, 12, 92
- epigraph, 355
- equation
  - Emden-Fowler, 231
  - Euler, 31
  - Laplace, x
  - Poisson, 158
  - scalar field, 173
  - Schrödinger, 123
- equilibrium point
  - Nash, 329
- essential value, 379
- Euclidean metric, 5
- Euler equation, 31
- Euler-Lagrange variational principle, 84
- exponential map, 286, 303
- Fatou lemma, 65, 361
- Fermat theorem, 7
- fibering
  - spherical, 44
- fibering constraint, 44
- fibering functional, 44
- fibering method, 123, 124
- fibering scheme, 41
- Finsler
  - manifold, 285, 292
  - metric, 300
- Finsler structure, 16
  - dual, 16
- Finsler-Banach manifold, 15
- Finsler-metric ball
  - backward, 287
  - forward, 287
- Finslerian-Poincaré disc, 306, 310
- first variation of arc length, 287
- fixed point set, 38
- fixed point set of a group action, 38
- flag curvature, 288, 289, 295
- flower petal theorem, 43
- fluid
  - dilatant, 86
  - Newtonian, 86
  - pseudo-plastic, 86
- fluid mechanics, 86
- force field, 123
- forward proximal set, 312
- Fréchet differentiability, 93, 94
- function
  - axially symmetric, 47
  - Carathéodory, 90, 100, 112
  - convex, 35, 37, 354
  - distance, 359
  - Gamma, 364
  - indicator, 47, 52–54
  - penalty, 8
  - quasi-concave, 27
- function of bounded variation, 59
- functional
  - coercive, 24
  - energy, 12
  - locally Lipschitz, 373
  - lower semi-continuous, 4
  - of Szulkin type, 35
  - perturbed, 5
  - recession, 71
  - upper semi-continuous, 50
- G-space, 290
- Gâteaux differentiability, 7, 21, 22
- Gamma function, 364
- gauge theory, 2
- generalized
  - Cauchy-Schwarz inequality, 289
  - directional derivative, 302
  - gradient, 301
  - length of a curve, 290
- generalized directional derivative, 377
- generalized gradient, 378
- genus, 99
- geodesic, 286, 293, 307, 368
- geodesic convex, 288
- geodesically complete
  - backward, 288
  - forward, 288
- geometrical analysis, 2
- glacial sliding, 86
- global elliptic regularity, 369
- gravitational field, 307
- group-invariant
  - function, 38
  - set, 38
- Hölder inequality, 107
- Hölder space, 368
- Haar measure, 40
- Hahn-Banach theorem, 374
- Hamiltonian system, 154
- Hardy inequality, 88
- Hartman-Stampacchia theorem, 45
- Hele-Shaw flow, 86
- hemivariational inequality, 59
- Hilbert

- distance, 292
  - metric, 291
- homology
  - Čech, 296
- Hopf lemma, 280
- Hopf maximum principle, 369
- hyper-surface, 84
- hyperbolic conservation law, 83
- hyperbolic space, 265
- image processing, 83
- indefinite nonlinearity, 199
- indicator function, 47, 52–54
- inequality
  - Hölder, 107
  - hemivariational, 59
  - interpolation, 132
  - minimax, 181
  - nonconvex, 45
  - Poincaré, 204, 363
  - variational, 45
  - weighted Hardy-type, 88
  - Young, 89
- integral length, 287, 291
- interpolation inequality, 132
- invariant
  - function, 264
  - set, 250
- inverse image, 381
- isometry group, 265
- Jacobi field, 286
- Kardar-Parisi-Zhang system, 125
- kinetic energy, 123
- KKM theorem, 296
- Krasnoselski genus, 19, 371
- Krasnoselskii–Zabreiko theorem, 43
- Kuratowski upper limit, 383
- Lagrange multiplier method, 355
- Lagrange multiplier theorem, 19
- Lagrange multipliers, 99
- Laplace equation, x
- Laplace–Beltrami operator, 267
- Laplace–Beltrami operator, 232, 244
- Lebesgue decomposition, 70
- Lebesgue dominated convergence theorem, 222
- Lebesgue measure, 204
- Lebesgue space, 360
  - weighted, 88
- Lebourg mean value theorem, 375
- lemma
  - Brézis–Lieb, 361
  - deformation, 376
  - Fatou, 65, 361
  - Hopf, 280
  - Phelps, 8
  - pseudogradient, 43
- level set, 14
- Levi-Civita connection, 266, 285
- Lipschitz perturbation, 11
- Ljusternik–Schnirelmann category, 370
- local minima, 117, 118
- local minimizer, 31
- local minimum, 37
- locally
  - geodesic space, 291
  - Lipschitz function, 299, 301
  - Minkowski space, 289
- locally Lipschitz functional, 373
- location property, 203
- lower critical point, 378
- lower semi-continuity, 5
- lower semi-continuous functional, 4
  - set-valued map, 381
- manifold
  - Banach, 15
  - closed symmetric, 99
  - Finsler–Banach, 15
  - Riemann, 280
- Markovian property, 172
- maximum principle, 34
  - Hopf, 369
- mean curvature, 84
- mean value theorem, 129
  - Lebourg, 375
- measurable functions, 90
- measure
  - Haar, 40
  - Lebesgue, 204
- method
  - direct, 224
  - fibering, 41
  - Lagrange multiplier, 355
  - sub and supersolutions, 34
  - topological, 195
- metric
  - Riemannian, 280, 281
- metric ball
  - backward, 290
  - forward, 290
- metric space, 373
  - complete, 43
- metric tensor, 267
- minimal geodesic, 287
- minimal surface, 84
- minimax inequality, 181
- minimization
  - compact case, 3
  - noncompact case, 3
- minimization principle
  - Takahashi, 43
- minimizer
  - local, 31

- strict, 8
- strong, 8
- minimizing sequence, 43
- minimum
  - local, 37
- Minkowski norm, 145
- Minkowski space, 289
- Minty-type variational inequality, 297
- Morse theory, 43
- mountain pass geometry, 121, 224
- mountain pass landscape, 14
- mountain pass point, 13
- mountain pass theorem, 13, 15, 123
- multiplicity, 2
- Nash contingent point, 334
- Nash critical point, 341
- Nash equilibrium point, 329
- Nash-Minty point, 330
- Nash-Stampacchia point, 330
- Nemytskii operator, 90, 91, 93, 100
- non-positive curvature, 290
- non-positively curved metric space, 290
- nonconvex inequality, 45
- nonlinear equation
  - normal solvability, 43
- nonlinear field equation, 87
- nonlinearity
  - oscillatory, 195
- nonstable critical point, 12
- normal cone, 359
- normalized Haar measure, 40
- one-parametric fibering method, 41, 158
- operator
  - $p$ -Laplace, 282
  - averaging, 40
  - Laplace-Beltrami, 232, 244
  - Nemytskii, 90, 91, 93, 100
  - quasilinear, 86
  - trace, 110
- operator equation
  - normal solvability, 43
- optimal path, 13
- orthogonal group, 245
- oscillation, 224
  - at infinity, 212
  - near the origin, 206
- oscillatory nonlinearity, 195
- pair
  - nontrivial, 162
  - trivial, 379
- Palais-Smale condition, 7, 17, 376
- parallel transport, 368
- path
  - optimal, 13
- penalty function, 8
- Penot flower petal theorem, 43
- perturbation
  - Lipschitz, 11
  - superlinear, 11
- perturbed functional, 5
- Phelps lemma, 8
- Plateau problem, 84
- Poincaré inequality, 204, 363
- point
  - mountain pass, 13
- Poisson bracket, 267
- Poisson equation, 158
- porous media, 86
- positive altitude, 13
- positive cone, 356
- potential, 123
  - sublinear, 136
  - superlinear, 128
- potential energy, 123
- power-law fluids, 86
- principle
  - concentration compactness, 124
  - symmetric criticality, 38
  - Takahashi, 43
  - uncertainty, 123
- principle of symmetric criticality, 2, 38
- product space, 126
- proper map, 48
- property
  - location, 203
  - Markovian, 172
- pseudogradient lemma, 43
- Pucci-Serrin theorem, 20
- quantum field theory, 2
- quantum mechanics, 87
- quasi-concave function, 27
- quasilinear eigenvalue problem, 87
- quasilinear operator, 86
- Radon-Nikodym derivative, 70
- Randers space, 306
- Rayleigh quotient, 97
- recession functional, 71
- reference vector, 286
- reflexive Banach space, 3
- Rellich-Kondrachov theorem, 368
- representation of a group, 38
- Ricceri variational principle, 37
- Riemann-Finsler geometry, 300
- Riemannian
  - manifold, 289
- Riemannian connection, 266
- Riemannian manifold, 280
  - complete, 231, 288
  - with boundary, 281
  - without boundary, 282
- Riemannian metric, 280, 281
- rule

- chain, 73
- derivative, 286
- Lagrange multipliers, 99
- saddle point theorem, 379
- scalar curvature, 267, 281
- scalar field
  - system, 173
- scalar field systems, 173
- Schauder regularity, 32
- Schrödinger equation, 123
- sectional curvature, 288, 295
- separable Asplund space, 8
- separable Banach space, 18
- sequentially weakly continuous, 113
- set
  - acyclic, 296
  - Chebyshev, 356
  - closed, 9
  - compact, 17
  - convex, 353
  - open, 24
  - upper contour, 355
  - weakly closed, 178
- set-valued analysis, 381
- set-valued map, 381
  - $K$ -pseudo-convex, 336
- shear stress, 86
- shock waves, 83
- sign-changing solution, 245, 264
- sleek, 383
- smooth bump, 8
- Smulyan test, 8
- Sobolev embedding theorem, 363
- Sobolev exponent
  - critical, 46
- Sobolev space, 360
  - weighted, 88
- solution
  - shift, 144
  - sign-changing, 264
  - strong, 159
  - weak, 87
- space
  - Asplund, 8
  - Banach, 3
  - functions of bounded variation, 59
  - Hölder, 368
  - Lebesgue, 360
  - metric, 373
  - product, 126
  - Sobolev, 360
  - tangent, 16
  - topological, 27
  - weighted Sobolev, 88
- spherical fibering, 44
- stable critical point, 11
- Stampacchia-type variational inequality, 297
- stationary variational inequality, 46
- Stegall variational principle, 8
- strict minimizer, 8
- strip-like domain, 46, 51
- strong extremum, 8
- strong minimizer, 8
- subdifferential, 35
- sublinear potential, 136
- subsolution, 33
- superconductivity, 2
- supercritical growth, 245
- superlinear case, 99
- superlinear perturbation, 11
- superlinear potential, 128
- supersolution, 33
- symbols
  - Christoffel, 267
- symmetric criticality
  - principle, 38
- symmetric space, 265
- system
  - Kardar-Parisi-Zhang, 125
  - non-variational, 154
  - of gradient type, 125
  - scalar field, 173
- Szulkin-type functional, 35
- Takahashi minimization principle, 43
- tangent bundle, 16
- tangent space, 16
- tangent vector, 359
- theorem
  - Ambrosetti–Rabinowitz, 43
  - Arzelà–Ascoli, 32
  - Bishop–Phelps, 8
  - Caristi, 227
  - Cartan–Hadamard, 289
  - divergence, 88
  - dominated convergence, 222, 361
  - drop, 43
  - Eberlein–Smulyan, 148
  - Fermat, 7
  - flower petal, 43
  - Hahn–Banach theorem, 374
  - Hartman–Stampacchia, 45
  - Hopf–Rinow, 288
  - Krasnoselskii–Zabreiko, 43
  - Lebourg mean value theorem, 375
  - mean value, 129
  - mountain pass, 43, 123
  - Pucci–Serrin, 20
  - Rellich–Kondrachov, 368
  - saddle point, 379
  - Sobolev embedding, 363
  - Weierstrass, 3
- thermodynamic variables, 353

- topological invariant, 15
- topological space, 27
  - compact, 3
- Torricelli point, 299
- torsional creep, 86
  - elastic, 86
  - plastic, 86
- total
  - s-backward cost, 301
  - s-forward cost, 301
  - backward cost, 299, 301
  - forward cost, 299, 301
  - variation, 70
- total energy, 123
- trace operator, 110
- transitive group action, 246
- trivial pair, 379
- truncation map, 32
- unbounded domain, 87
- uncertainty principle, 123
- unit outward normal, 91
- unit sphere
  - volume, 364
- upper contour set, 355
- upper limit
  - Kuratowski, 383
- upper semi-continuous, 116
  - functional, 50
  - set-valued map, 381
- value
  - critical, 35
  - essential, 379
- variational inequality, 45, 46, 52
- variational principle
  - Borwein-Preiss, 8
  - Ekeland, 4, 58
  - Euler-Lagrange, 84
  - Ricceri, 37
  - Stegall, 8
- vector
  - field, 264
  - space, 52
  - tangent, 359
- vertex, 356
- weak
  - partial derivative, 362
  - distributional derivative, 361
  - slope, 376
  - solution, 92
- weakly closed set, 178
- weakly convergent, 113, 118
- Weierstrass theorem, 3
- weight function, 88
- weighted Hardy-type inequality, 88
- weighted Lebesgue space, 88
- weighted Sobolev space, 87, 88
- Yamabe problem, 124, 232, 266
- Young inequality, 89
- zero altitude, 15